

1. All the following drugs are safe to give during pregnancy **EXCEPT:**
- A- Paracetamol
  - B- Insulin
  - C- Cephalosporins
  - D- Phenytoin XX
2. Which of the following statements is **false?**
- A- Heparin can be given during pregnancy
  - B- Highly ionized drugs cross the placenta slowly
  - C- Drugs with high lipid solubility cross the placenta poorly xx
  - D- Warfarin should be avoided during pregnancy
3. All the following drugs are contraindicated in pregnancy **EXCEPT:**
- A- Thalidomide
  - B- Warfarin
  - C- Tetracycline
  - D- Heparin xx
4. All the following drugs should be avoided during lactation **EXCEPT:**
- A- Aspirin
  - B- Tetracycline
  - C- Radioactive iodine
  - D- Amoxicilin xx
5. All the following statements about type A (Augmented) adverse drug reactions are true **EXCEPT:**
- A- Account for 80% of adverse drug reactions
  - B- Are due to excessive of pharmacological activity of drug
  - C- Dose-related
  - D- Unpredictable XX
6. Which of the following statement is **false?**
- A- Older people have an increased risk to develop adverse drug reactions
  - B- Genetic factors may predispose to development of adverse effects of drugs
  - C- Hepatic diseases increase the risk of adverse drug reactions
  - D- Heart diseases can cause adverse drug reactions xx
7. All the following statements about drug interactions are true **EXCEPT:**
- A- Drug interactions may occur outside the body
  - B- Drug interactions may occur in the intestine
  - C- Oral contraceptive is metabolized rapidly when given with phenytoin
  - D- Naloxone is given to treat morphine overdose is a harmful interaction xx
8. Which of the following statements about the antidotes is **false?**
- A- Acetylcysteine is given in paracetamol overdose
  - B- Vitamin K is given in warfarin overdose
  - C- Atropine is given in benzodiazepine overdose xx
  - D- Naloxone is given in opioids overdose

**9. All the following the statements are true EXCEPT:**

- A- Activated charcoal reduces absorption of drugs from the intestine
- B- Gastric lavage should be performed within 1 hour of poisoning
- C- Emesis is induced in unconscious patients **xx**
- D- Both emesis and lavage are contraindicated for corrosive poisons