

## General Microbiology Course Lecture 3 (Microbial Growth and Metabolism)

Dr. Mohammad Odaibat Department of Microbiology and Pathology Faculty of Medicine, Mutah University

# Lecture Outlines

- Some definitions.
- Microbial nutrition.
- Microbial growth and its requirements.
- Reproduction of microbes.
- Bacterial respiration.

## Definitions

- Metabolism: the sum of the chemical reactions in an organism.
- **Catabolism**: provides energy and building blocks for anabolism
- Anabolism: uses energy and building blocks to build large molecules



## Microbial nutrition

#### Types of microbial nutrition

#### Carbon source

- Heterotrophs must obtain carbon in an organic form made by other living organisms such as proteins, carbohydrates, lipids and nucleic acids
  - nutritionally dependent on other living things
- Autotrophs (self feeding): an organism that uses  $CO_2$ , an inorganic gas as its carbon source

## Microbial Growth:

Refers to an increase in **cell <u>number</u>**, not in cell size.

Bacteria grow and divide by **binary** fission, a rapid and relatively simple process. **Requirements for Growth** 

## Physical Requirements

## 1. <u>Temperature</u>

Microbes classified into several groups based on their preferred temperature ranges.

- A. <u>Psychrophiles</u> : "Cold-loving". Can grow at 0°C. Two groups:
  - True Psychrophiles

Optimum growth at 15 C or below. Found in very cold environments (North pole, ocean depths). Seldom cause disease or food spoilage.

**Psychrotrophs**: Optimum growth at 20 to 30 C. Responsible for most low temperature food spoilage.

## 1. <u>Temperature</u> :

B. <u>Mesophiles</u>: "Middle loving". Most bacteria.
Best growth between 25 to 40 €.

Optimum temperature commonly 37 C.

Many have adapted to live in the bodies of animals.

#### C. <u>Thermophiles</u> : "Heat loving".

ØOptimum growth between 50 to 60 ℃.

Adapted to live in sunlit soil, compost piles, and hot springs.

# Extreme <u>Thermophiles</u> (Hyperthermophiles): Optimum growth at 80° C or higher. Archaebacteria. Most live in volcanic and ocean vents.

## Growth Rates of Bacterial Groups at Different Temperatures



- 2. <u>pH</u>: Organisms can be classified as:
- A. <u>Acidophiles</u> : "Acid loving".

Grow at very low pH (0.1 to 5.4)

*Lactobacillus* produces lactic acid, tolerates mild acidity.

#### B. <u>Neutrophiles</u> :

Grow at pH 5.4 to 8.5.

Includes most human pathogens.

C. <u>Alkaliphiles</u> : "Alkali loving".

☑Grow at alkaline or high pH (7 to 12 or higher)☑*Vibrio cholerae* optimal pH 9.

- **3. Osmotic Pressure** : Cells are 80 to 90% water.
- ☑Halophiles \_: Require moderate to large salt concentrations. Ocean water contains 3.5% salt.
  ☑Most bacteria in oceans.
- **Extreme or Obligate Halophiles** : Require very high salt concentrations (20 to 30%).
  - Bacteria in Dead Sea.
- ☑ Facultative Halophiles : Do not require high salt concentrations for growth, but tolerate 2% salt or more.

1. <u>Carbon</u>: Makes up 50% of dry weight of cell.

Structural backbone of all organic compounds.

- ■Chemoheterotrophs : Obtain carbon from their energy source: lipids, proteins, and carbohydrates.
- Chemoautotrophs and Photoautotrophs : Obtain carbon from carbon dioxide.

- 2. Oxygen: bacteria are classified into
- 1- Aerobes: utilizes oxygen and can detoxify it
- **obligate aerobes**: cannot grow without oxygen
- **facultative anaerobes**: utilize oxygen but can also grow in its absence
- microaerophilic : requires only a small amount of oxygen
- 2- Anaerobes : do not utilize oxygen
- **obligate anaerobes**: lack the enzymes to detoxify oxygen so cannot survive in an oxygen environment
- **aerotolerance anaerobes**: do no utilize oxygen but can survive and grow in its presence

#### Categories of oxygen requirement



#### Anaerobic conditions



Q: Why do some bacteria are forced to live in oxygen free habitats?

- To avoid the toxic forms of oxygen (Oxygen free radicals)
- If a microbe is not capable of dealing with toxic oxygen, it is forced to live in oxygen free habitats.
- The oxygen free radicals including:
  - A. Superoxide Free Radicals  $(O_2^{-.})$
  - B. Hydrogen Peroxide  $(H_2O_2)$

The oxygen free radicals

1. <u>Hydrogen Peroxide</u>  $(H_2O_2)$ :

There are two different enzymes that break down H2O2:

A. Catalase: Breaks hydrogen peroxide into water and O<sub>2</sub> Common. Produced by humans, as well as many bacteria.

 $2 H_2O_2 \xrightarrow{\text{Catalase}} 2H_2O + O_2$ Hydrogen Gas peroxide Bubbles

B. Peroxidase: Converts hydrogen peroxide into water. Peroxidase  $H_2O_2+2H \longrightarrow H_2O_2$ Hydrogen peroxide

#### The oxygen free radicals

2- <u>Superoxide Free Radicals</u>  $(O_2^{-})$ : Extremely toxic and reactive form of oxygen. All organisms growing in atmospheric oxygen must produce an enzyme **superoxide dismutase** (SOD), to get rid of them. SOD is made by aerobes, facultative anaerobes, and aerotolerant anaerobes, **but not** by anaerobes or microaerophiles.

#### <u>Reaction</u> :

$$O_2^{-.} + O_2^{-.} + 2H^+ ----> H_2O_2^+ O_2$$

Superoxide free radicals Hydrogen peroxide

The oxygen free radicals



# Microbial Growth

#### Reproduction of microbes

- Microbial growth an increase in a population of microbes rather than an increase in size of an individual
- Result of microbial growth is discrete colony an aggregation of cells arising from single parent cell





# Microbial Growth

#### Reproduction of microbes Binary Division

- division exactly in half
- most common means of bacterial reproduction
  - forming two equal size progeny
  - genetically identical offspring
  - cells divide in a geometric progression doubling cell number



# Microbial Growth

## Bacteria grow in four stages

- lag phase "flat" period of adjustment, enlargement; little growth
- 2. **exponential or log phase** a period of maximum growth will continue as long as cells have adequate nutrients and a favorable environment
- 3. **stationary phase** rate of cell growth equals rate of cell death caused by depleted nutrients and  $O_2$ , excretion of organic acids and pollutants
- 4. **death phase** as limiting factors intensify, cells die exponentially in their own wastes

## **Bacterial physiology**

#### 4- Reproduction of microbes

#### The population Growth Curve



## **Bacterial Respiration**



#### Aerobic vs Anaerobic Respiration



#### Fermentation = Oxygen not present



## Fermentation





## Products of Fermentation

Without any form of respiration, glycolysis products, pyruvate and NADH, will accumulate. To keep making any more ATP by glycolysis, fermenting cells must convert NADH (red.) back to NAD+ (ox.) by passing its electrons to pyruvate. Reaction pathways that do this convert pyruvate to many other compounds, depending on the organism.



## **General Pathways**



# Thank you