

The digestive system II



The gastro- intestinal tract:

Composed of:

- Esophagus
- Stomach
- Small intestine
- Large intestine
- Anal canal



General features of the wall of the GIT

its wall is composed of 4 layers:

☐ Mucosa:

Epithelium

CT (Lamina propria, corium)

→ Muscularis mucosa (s. ms.)

☐ Submucosa: C.T.

☐ Musculosa : 2 layers of

smooth muscles (IC & OL)

■ Adventitia or serosa

The mucosa is a mucous membrane that lines the GI tract and secretes vessels mucus that lubricates and protects the GI tract. Lymphatic vessel The submucosa is a layer Nerve of connective tissue that contains blood vessels, lymph vessels, and nerves. The muscularis is made up of two layers of smooth muscle—one circular and one longitudinal. The serosa is a connective tissue covering that secretes a fluid to lubricate the outside of the GI tract.

Adventitia vs. serosa

Serosa: double layer membrane made of epithelium

One layer is attached to the organ called visceral layer, the other layer will be close to the body cavity & called partial layer. In between these two epithelial layer is fluid called serous for lubrication (reduce friction)

Serosa will wrap organs that set in a body cavity i.e abdominal cavity like GIT organs within the peritoneum i.e intraperitoneal organs (liver, stomach, spleen, 1st part pf duodenum, ileum, jejunum, transverse & sigmoid colon)

Adventitia: is not epithelial is loose CT that wraps organs that set outside the peritoneal cavity i.e. retroperitoneal and attach them to the abdominal cavity

pancreas, rest of duodenum, cecum, ascending & descending Colcon


The esophagus


- Muscular tube connects the pharynx with stomach, transport food
- Its wall consists of 4 layers:
- Mucosa

Submucosa:

Musculosa

Adventitia





Mucosa

Epithelium: Non-keratinized stratified squamous epith.
Lamina propria: B.V., nerves, lymphatics (!Cardiac orifice)
Muscularis mucosa: smooth ms.

- Submucosa: loose C.T. contains BV, lymphatics, <u>Meissner's</u>
 plexus of nerves & esophageal mucous glands
- Musculosa: IC &OL (<u>OL</u>: upper 1/3 Striated *, middle 1/3 mixed & lower 1/3 smooth ms.) NB: swallowing start with controllable motion but finishes with involuntary peristalsis
- Adventitia: covers most of the esophagus except the most distal portion which is located in the abdominal cavity is covered by serosa


Layers of the wall of the esophagus Stratified Squamous Epithelium Mucosa Lymphoid Tissue Lamina Propria Muscularis Mucosa Blood Vessels & Submucosa Glands Circular Muscle Neural Plexus Musculosa Longitudinal Muscle **Adventitia**


Prof Dr H Elmazar

7

Changes at gastro- esophageal junction

- 1. The stratified Squamous \rightarrow simple columnar epithelium
- The lamina propria of stomach is wide & contains gastric glands (branched tubular)
- 3. The esophageal glands in the submucosa of esophagus stops in that of stomach
- 4. The musculosa becomes more thick in stomach due to the appearance of inner oblique layer






Layers of wall of stomach

Opening of Gastric Glands Lamina Propria Muscularis Mucosa Submucose Oblique Muscle Layer Circular Muscle Layer Longitudinal Muscle Layer 🎇 Serosa Frank Boumphrey M.D. 2009

Layers of wall of esophagus




Prof Dr H Elmazar

9

The stomach

- The most dilated part of the GIT
- The mucosa in empty stomach forms longitudinal folds called gastric rugae
- It acidifies & converts the food → chyme



- The mucosa of stomach contains gastric glands (cardiac, fundic, pyloric)
- These glands secrete gastric juice which contains:
- > Acid: HCl
- Mucus
- > enzymes: pepsinogen, lipase

The stomach


The stomach is subdivided into 4 regions:

1. The cardiac region


2. The fundus

3. The body

4. The pyloric region




The fundus & body of the stomach



1- The mucosa:

- epithelium: simple columnar cells, these cells secrete neutral mucus for lubrication & protection*
- lamina propria: contains <u>gastric glands</u> & C.T. fills the spaces between the glands. It also contains B.V., lymphatics, nerves

The gastric (fundic) glands




Muscularis mucosa: layer of smooth muscles arranged as (IC & OL)


inner circular & outer longitudinal

Gastric glands (fundus)

- simple branched tubular.
- o ccupy the entire thickness of the mucosa.
- They open onto the surface epithelium through gastric pits.
- through the pits the mucus, HCl & gastric enzymes reach the lumen of the stomach




- <u>Each gland</u> is formed of 3 parts: isthmus, neck & base
- 6 types of cells line the fundic glands:
- 1- Surface mucous cells (Foveolar cells): cover the surface & line the gastric pits & isthmus. Their apical cytoplasm contains mucin granules.
- They sec. <u>neutral mucus</u> for protection (Gastric mucosal barrier)
- 2- Mucous neck cell: present in neck of gastric glands, low columnar cells e foamy cytoplasm. They secrete <u>acidic mucus</u>



Prof Dr H Elmazar

Gastric mucosal barrier




- 1- epithelial cell lining. Cells in the epithelium of the stomach are bound by tight junctions
- 2- A special mucus covering, secreted by surface epithelial cells. This insoluble mucus forms a protective gel-like coating over the entire surface of the gastric mucosa.
- 3- Bicarbonate ions, secreted by the surface epithelial cells. The bicarbonate ions act to neutralize harsh acids that find access to


cells Prof Dr H Elm


3- <u>stem cells</u>: present in <u>neck region</u>, low columnar. They differentiate to other gastric cells

4- Parietal (oxyntic) cells:

- triangular in shape e <u>acidophilic</u>
 cytoplasm & rounded central nucleus.
 present mainly in the upper half of the glands fewer in the base
- <u>E/M</u>: their apical surfaces show branching Intracellular canaliculi that open at the apex.
- ↑ mitochondria, ↑SER, NO sec. granules
- They secret HCl & intrinsic factor(glycoprotein)
 needed for vit. B12 absorption









Oxyntic cell secretes HCl & intrinsic factor

Formation of HCL






Showing tubulovesicular system in active vs resting parietal cell

Pernicious Anemia

- · Pernicious anemia is caused by a lack of intrinsic factor
- Intrinsic factor is a protein made in the stomach. It helps your body absorb vitamin B12, necessary for normal RBC production; RBCs are larger






One of the signs of pernicious anemia is red tongue with smooth surface (Beefy tongue)


Prof Dr H Elmazar

21

5-Peptic (Chief, Zymogenic) cells: mainly at the base of gastric glands. columnar cells e basal rounded nuclei.


- The basal cytoplasm is basophilic due to 个rER, while the apical part contains 个个 zymogen granules
- E/M : protein secreting cells
- These cells secrete pepsinogen & G. lipase







6- Entero-endocrine cells :

- present in the base of the glands.
- Hormone secreting cells
- (diffuse neuroendocrine system)
- Their secretions accumulates in the basal part to be released to the B.V.
- They secrete:
- ✓ Gastrin
- ✓ Enteroglucagon
- ✓ Serotonine
- ✓ Somatostatin(D cells)



Prof Dr H Elmazar





2- The submucosa: loose C.T. with B.V., lymphatics, meissner's plexus of nerves

3- The musculosa: formed of 3 layers of smooth ms.

Inner oblique - middle circular - outer longitudinal.

Auerbach's plexus is present between middle & outer layers

4- The Serosa: is the peritoneal covering, is formed simple squamous mesothelium & loose C.T. It contains B.V., lymphatics, & nerves

The difference between fundus & pylorus


Fundus

- Thick mucosa
- Pits are narrow & short
- F. Glands are simple branched tubular & long
- occupy most of mucosal thickness
- Lined e 6 types of cells
- Corium: lymphocytic infiltration
- Musculosa: thinner formed of <u>3 layers</u> of ms. (IO, MC,OL)

<u>Pylorus</u>


- Thin mucosa
- Pits are wide & long
- P. Glands are coiled branched tubular & short
- Occupy ½ of mucosal thickness
- Lined e mucous secreting cells
 No oxyntic, No peptic cells
- Lymphocytic infiltration & lymph nodules
- Thicker, formed of 2 layers of muscles. Thick IC to form the p. sphincter & OL

Difference between fundic & pyloric glands




Changes at gastro duodenal junction

- intestinal villi start to project from mucosa
- Intestinal crypts replace pyloric glands in the corium of duodenum
- Surface columnar cells with brush border. Goblet cells appear between cells
- Muscularis mucosa: pass unchanged
- Brunner's glands appear in duodenal submucosa
- Musculosa is thinner in the duodenum
- Serosa pass unchanged



Prof Dr H Elmazar





Gastro duodenal junction

Prof Dr H Elmazar 28

Wall of intestine

Wall of stomach





Thank you

