



Chemical Foundations



The Fundamental SI Units

<u>Physical Quantity</u>	<u>Name of Unit</u>	<u>Abbreviation</u>
Mass	kilogram	kg
Length	meter	m
Time	second	S
Temperature	kelvin	K
Electric current	ampere	А
Amount of substance	mole	mol.
Luminous intensity	candela	cd



Prefixes Used in the SI System

• Prefixes are used to change the size of the unit.

 Table 1.2
 Prefixes Used in the SI System (The most commonly encountered are shown in blue.)

Prefix	Symbol	Meaning	Exponential Notation*
exa	E	1,000,000,000,000,000,000	1018
peta	Р	1,000,000,000,000,000	1015
tera	т	1,000,000,000,000	1012
giga	G	1,000,000,000	10 ⁹
mega	M	1,000,000	106
kilo	k	1,000	10 ³
hecto	h	100	10 ²
deka	da	10	101
		1	100

© Cengage Learning. All Rights Reserved.



Prefixes Used in the SI System

Table 1.2 Prefixes Used in the SI System (The most commonly encountered are shown in blue.)

Prefix	Symbol	Meaning	Exponential Notation*
deci	d	0.1	10-1
centi	С	0.01	10-2
milli	m	0.001	10-3
micro	μ	0.000001	10-6
nano	n	0.00000001	10-9
pico	р	0.0000000001	10-12
femto	f	0.00000000000001	10-15
atto	а	0.0000000000000000000000000000000000000	10-18

*See Appendix 1.1 if you need a review of exponential notation.

Congage Learning. All Rights Reserved



Exponential Notation (scientific notation)

- Example 5000. : 5.000x10³ 4 S.F
- $5000 : 5x10^3$
- $381000000000000 = 3.81 \times 10^{17}$
- $0.00000000914 = 9.14 \times 10^{-10}$
- 5.00X10³ 3 S.F
 - 300. written as 3.00×10^2
 - Contains three significant figures.
- Two Advantages
 - Number of significant figures can be easily indicated.




Exponential Notation and Significant Figures:

90000	1 S.F	
90000 .	5 S.F	
9.0000x10 ⁴	5 S.F	
■ 9.0x10 ⁴	2 S.F	
■ 9.00X10 ⁴	3 S.F	
6510000000000000	6.51x10 ¹⁵	3 S.F
• 0.0000000000002710	2.710 x10 ⁻¹⁷	4 S.F
■ () × 10×		

Section 1.5 Significant Figures and Calculations

Measurement of Volume Using a Buret

- The volume is read at the bottom of the liquid curve (meniscus).
- Meniscus of the liquid occurs at about 20.15 mL.
 - Certain digits: 20.15
 - Uncertain digit: 20.15





- 1. Nonzero integers always count as significant figures.
 - 3456 has 4 sig figs (significant figures).


300000



- 2. There are three classes of zeros.
- a. <u>Leading zeros</u> are zeros that precede all the nonzero digits. These do not count as significant figures.
 - 0.0000048 has 2 sig figs.



- b. <u>Captive zeros</u> are zeros between nonzero digits. These always count as significant figures.
 - 16.07 has 4 sig figs.
 - 0.00807 3 s.f.
 - 2.00002001 9 s. f.



- <u>Trailing</u> zeros are zeros at the right end of the number. They are significant only if the number contains a decimal point.
 - 9.300 has 4 sig figs.
 - 150 has 2 sig figs.
 - 23100000 3 s.f.
 - -----
 - 410070000 5 s.f.
 - 4.10070000 9 s.f.
 - 500 3cf

Section 1.4 Uncertainty in Measurement



Precision and Accuracy

Accuracy


• Nearness of the measurements to the true value.

Precision

• Nearness of the measurements to each other.

Section 1.4 Uncertainty in Measurement


Precision versus Accuracy



Neither accurate nor precise. © Cengage Learning. All Rights Reserved.



Precise but not accurate.



Accurate and precise.






Three Systems for Measuring Temperature

- Fahrenheit
- Celsius
- Kelvin



The Three Major Temperature Scales





Converting Between Scales

$$T_{\rm K} = T_{\rm C} + 273.15$$
 $T_{\rm C} = T_{\rm K} - 273.15$

$$T_{\rm C} = \left(T_{\rm F} - 32^{\circ} {\rm F}\right) \frac{5^{\circ} {\rm C}}{9^{\circ} {\rm F}} \qquad T_{\rm F} = T_{\rm C} \times \frac{9^{\circ} {\rm F}}{5^{\circ} {\rm C}} + 32^{\circ} {\rm F}$$



Example

What is the F equivalent of 35 °C?
 35 °C x 9/5 + 32 = 95 °F

- What is the equivalent of 151 °F in K?
 First convert into °C then to K.
 - °C: (151 − 32) x 5/9 = 66.1 °C
 - K: 66.1 + 273.15 = 339.3 K





At what temperature does $^{\circ}C = ^{\circ}F$?



EXERCISE!

- Since °C equals °F, they both should be the same value (designated as variable x).
- Use one of the conversion equations such as:

$$T_{\rm C} = \left(T_{\rm F} - 32^{\circ}{\rm F}\right) \frac{5^{\circ}{\rm C}}{9^{\circ}{\rm F}}$$

• Substitute in the value of x for both $T_{\rm C}$ and $T_{\rm F}$. Solve for x.

EXERCISE!

$$T_{\rm C} = \left(T_{\rm F} - 32^{\circ}{\rm F}\right) \frac{5^{\circ}{\rm C}}{9^{\circ}{\rm F}}$$

$$x = \left(x - 32^{\circ}F\right)\frac{5^{\circ}C}{9^{\circ}F}$$

$$x = -40$$

So
$$-40^{\circ}\text{C} = -40^{\circ}\text{F}$$

