


# Meiosis

Ass. Prof Dr. Heba Hassan Abd El-Gawad



# Cell Cycle (Replicating)




# Cell Division


## I-Mitosis: in somatic cells

→ giving **two** daughter cells with the **same** number of **chromosomes** & **same** amount of **DNA** as parent cell.


## II-Meiosis: in germ cells

→ giving **four** daughter cells with **half** number of **chromosomes** & **half** amount of **DNA** (gametes).









DNA can be found in two forms:

- **Chromatin**: Hetero+Eu (interphase nucleus)



- **Chromosome**: tightly wound, condensed form (during division)

**1n** = amount of DNA in **23s** chromosomes.

**2n** = amount of DNA in **23d or 46 = 23pair s** chromosomes.

**4n** = amount of DNA in **46d** chromosomes.

# Meiosis!



**Definition** :It is a special form of cell division in which the chromosomes number is reduced from diploid ( $2n$ ) to haploid ( $1n$ ).

The DNA content of the original diploid cell is duplicated during the S-phase of the cell cycle before starting meiosis. The cell at the start of meiosis contains diploid number ( $46d$ ) & double amount of DNA ( $4n$ ).

Gives rise to four daughter cells: haploid number ( $23s$ ) & half the amount of DNA ( $1n$ ).

It occurs only in the germinal cells of the testis & ovary to produce gametes; (ova & spermatozoa) so, called gametogenesis.

## Phases of Meiosis:

It consists of two divisions:

1. Meiosis I: The cell with **46d** chromosomes divides giving rise to two daughter cells each has **23 d**-chromosomes and the amount of DNA is **2n**. Each chromosome is formed of 2 chromatids.


2. Meiosis II: In this division there is **no s-phase**. The **two** chromatids of each chromosome are separated, as in mitosis, leading to the formation of **two daughter cells each has 23s-chromosome** (haploid number of chromosomes and the amount of DNA is **1n**).



# STAGES OF MEIOSIS:


## MEIOSIS I

1. Prophase I
2. Metaphase I
3. Anaphase I
4. Telophase I



## MEIOSIS II

5. Prophase II
6. Metaphase II
7. Anaphase II
8. Telophase II



# Meiosis I اغلب الأسئلة هنا

## Prophase I:



There are 5 stages, each named based on the morphology of the chromosomes.

- 1- **Leptotene** (thin thread)
- 2- **Zygotene** (paired thread)
- 3- **Pachytene** (thick thread)
- 4- **Diplotene** (two thread)
- 5- **Diakinesis** (moving through)



## 1-Leptotene (thin thread):

- The **46 d-chromosomes** are visible & appear as long strands.
- Each consists of two chromatids attached by centromere.




## 2-Zygotene (paired threads):



- During this stage, **homologous** chromosomes (maternal & paternal) begin to synapse (**unite**) with each other by **synaptonemal complex** through a mechanism called (zipper like pairing).
- The paired homologous chromosomes are called **bivalents**. Each bivalent is formed of four chromatids (tetrad).



### 3. Pachytene (thick threads):


- The chromosomes become shorter & thicker.
- Crossing over between non-sister homologous chromatids takes place through chiasmata (site of exchange of genetic material).





#### 4. Diplotene (two threads):

- Longitudinal **separation** & more condensation of the two homologous chromosomes **except at the chiasmata**.
- The crossed area or chiasmata makes attachments between the chromosomes holding each bivalent (tetrad) together until metaphase.



## 5. Diakinesis (moving through):


- Maximum condensation of chromosomes (the d-chromosome becomes thicker & shorter).

## 5. Diakinesis (moving through):

-Maximum condensation of chromosomes (the d-chromosome becomes thicker & shorter).

-Disappearance of the nucleolus & nuclear envelope



-The chromosomes become **free** in the cytoplasm & the cell proceed to metaphase.





# Metaphase I:




- Shortest phase
- The homologous chromosomes (bivalents) or tetrads are arranged in equatorial plane.





sister chromatids fuse and function as one. Microtubules can attach to only one side of each centromere.

# Anaphase I:

- Homologous chromosomes migrate away from each other, going to opposite poles.
- Each chromosome still consists of two chromatids.



# Telophase I:



- The chromosomes reach the opposing poles, **nuclei are reformed.**
- Cytokinesis occurs giving rise to two daughter cells each cell contains **23d** chromosomes (**haploid number** of chromosomes)
- The amount of DNA is  $(2n)$  as each chromosome consists of two chromatids.

# II- Meiosis II

- It is like mitosis without prior DNA replication (**no S stage**, **no interphase**)

telophase & cytokinesis





prophase II

metaphase II

anaphase II

telophase II




Cleavage furrow



**MEIOSIS II**

Sister chromatids separate

Haploid daughter cells forming



**Two haploid cells form; chromosomes are still double**

**During another round of cell division, the sister chromatids finally separate; four haploid daughter cells result, containing single chromosomes**



## II- Meiosis II

- The **chromosomes arranged in equatorial plane** & kinetochores attach to microtubules of mitotic spindle followed by **migration** of the chromatids to opposite **poles**
- **Cytokinesis** divides each of the two cells giving a total **of four daughter cells**.
- Each of the four cells contains 23 s-chromosomes (haploid number of chromosomes) & (1n) amount of DNA.

## Major Phases of Meiosis:

- Meiosis 1: (S stage of interphase I)
- Meiosis 2: (NO S stage interphase 2)

46 s-chromosomes + 2n DNA

*S- Phase of cell cycle*



46 d-chromosomes + 4n DNA

**Meiosis I**



23 d-chromosomes + 2n DNA    Two daughter cells

**Meiosis II**



23 s-chromosomes + 1n DNA, Four germ cells, which may be ova or sperm



# Mitosis Vs meiosis?

	<b>Mitosis</b>	<b>Meiosis</b>
<b>Cell type</b>	Somatic	Gametes
<b># of divisions</b>	1	2
<b>S- phase</b>	Present	No s-phase between the two division
<b>Chromosome #</b>	Same as parent	Half of parent
<b># of daughter cells</b>	2	4
<b>Genetically identical?</b>	Yes	No
<b>Crossing over</b>	Absent	In prophase I
<b>When</b>	Throughout life	At sexual maturity
<b>Aim</b>	Growth and repair	Sexual reproduction

# Structure of human chromosome

**Chemically:** Chromosome consists of a continuous molecule of DNA double helix associated with proteins; histones and non-histones.

**Microscopically:** Each metaphase chromosome (d-chromosome) consists of two sister chromatids. Each one of the sister chromatid is formed of the followings:

- Short upper arm (p).
- Long lower arm (q).
- Centromere is a primary constriction connects the two chromatids together.
- Telomeres are the terminal regions at both ends of the chromosome.

Telomere protects chromosome ends from degradation and prevent end to end fusion with other chromosomes



**Metaphase chromosome  
formed of 2 sister chromatids**



**← Short arm (p)**

**Centromere →**

**← Long arm (q)**

**Telomeres**

احفظ  
المكان والفائدة



# Number:

## 1- Somatic cell:

In humans, the normal cell nucleus contains 46 chromosomes or 23 pairs (diploid number). One of each chromosome is paternal in origin (from father) and the other is maternal in origin (from mother). These pairs are divided into:

- 22 pairs called autosomes.
- One pair called sex chromosome.

A male has 44 autosomes and XY sex chromosomes. A female has 44 autosomes and XX sex chromosomes. In female, one X chromosome is an inactive which is called Barr body or sex chromatin.

~~1~~  
**2- Germ cell (gamete cells)** (ova and sperms) contains 23 s-chromosomes (haploid number). All the ova contain one type of sex chromosome (X), while half of the sperms contains X and the other half contains Y chromosomes.

### Sex chromatin or Barr body

**Definition:** It is dark staining heterochromatic mass lying against the inner surface of the nuclear membrane of interphase female cells only.

Barr body is believed to represent the inactivated X chromosome. All of the X chromosomes except one in a cell are inactivated early in development.

**Number:** The number of barr bodies in the female cell is equal to the number of X chromosomes present in the cell minus one. A female with chromosome number 47, XXX has two barr bodies, and a male with 49, XXXXY has three Barr bodies.

# Chromosome Classification

There are seven groups of chromosomes

الأرقام حسب

A: 1-3,

B: 4-5,

C: 6-12+X,

D: 13-15

E: 16-18,

F: 19-20

G: 21-22+Y

Chromosomes are classified according to

الأقصر هو  
Chromosome no. 22

## 1. Length of chromosome:



- Chromosomes are divided into 7 groups (A-F) according to their length:

- Group A contains chromosome numbers 1- 3 (the longest one).
- Group D contains acrocentric chromosomes, number 13-15 (medium-sized),
- Group G contains acrocentric chromosomes, (21 and 22) and Y chromosome (the shortest chromosomes).

## 2. Position of centromere

- Chromosomes are classified according to centromere position into three groups:

### a. Metacentric chromosomes:

- The centromere lies near the middle of the chromosome where  $p = q$ , such as number 1, 3, 16, 19 and 20.


### b. Submetacentric chromosome:

- The centromere lies closer to one end than the other.
- P arm is shorter than q arm such as number 2, 4, 12, 17, 18 and X chromosomes.


### c. Acrocentric chromosome:

- The short arm p is very small, and the centromere lies near one end such as chromosome number 13, 14, 15, 21, 22 and Y chromosome.
- These chromosomes except the Y have small chromatin masses known as satellites attached to their short arms by narrow stalk and contain genes for rRNA formation.






**Metacentric  
Chromosome  
Chromosome 3**



**Submetacentric  
Chromosome  
Chromosome 17**





**Acrocentric  
Chromosome  
Chromosome 21**

# Karyotyping & Chromosome banding

- **Karyotyping**: Study of the number, type & arrangement of chromosome.
- **Karyotype**: systemic arrangement of the chromosomal set of the cell
- Depends on length of chromosomes & position of centromere.
- Arrangement first autosomes & lastly the sex chromosomes (XX female or XY male)

Blood sample is taken



Incubate  
3-4 days




Add chemical to  
stop mitosis in  
metaphase




treated with hypotonic  
solution



Swollen cells are fixed,  
dropped in a glass  
slide, dried and stained



Examine with the  
microscope &  
photograph the  
chromosomes



Cut out the chromosomes  
and arrange into a karyotype

يُحصل أثناء الـ metaphase


## Steps of preparation:

- Culture of living cells in suitable medium
- Add **phytohemagglutinin** to stimulate division
- Add **colchicine** to arrest cells in metaphase
- Add **hypotonic** salt solution to swell the cells
- Spread on glass slides & Staining
- Photographing
- Cut chromosome individually
- Arrange by matching every two pairs in groups
- Seven groups (**A, B, C, D, E, F, G**) according to length and position of centromere
- Now by software of computer

رُحِمَ عليهنَّ بالمحاظرة الثالثة  
من العمر

Proteins  
glycoylex


# Normal male karyotyping ( 22+XY)





# Normal female karyotyping ( 22+XX)





Down  
female

---

1. Normal male: XY  
 2. Normal female: XX




3. Down syndrome: XXY  
 4. Klinefelter syndrome: XXY



5. Normal male: XY  
 6. Normal female: XX



7. Down syndrome: XXY  
 8. Klinefelter syndrome: XXY




9. Normal male: XY  
 10. Normal female: XX




# Down male

# Turner syndrome



## Chromosome banding:

Banding patterns are chromosomal patterns of **bright and dark transverse bands**. These bands identify where genes are located on a chromosome. The bright and dark bands are visible when the chromosome is stained with a chemical solution and examined under a microscope.



# Chromosome Banding Pattern

Chromosome banding patterns are available in various forms, such as:

**G-banding or Giemsa Staining,**  
**C-banding,**  
**Q-banding,**  
**R-banding**

## G-banding or Giemsa Staining

This is the **most basic chromosomal banding technique**. Because the whole complement of chromosomes is photographed, it can be used to detect genetic disorders.

**Giemsa = thiazine-eosin compound.**



## Positive G-bands

The **darkly** stained bands are positive G-bands. These areas are hydrophobic and facilitate the precipitation of the thiazine-eosin compound. They constitute the late replicating **heterochromatin**.

## Negative G-bands

Negative G-bands are **light-stained bands**. These are less condensed early replicating euchromatin. Base pairs from GC are abundant in these areas. These areas are less hydrophobic and less conducive to the precipitation of **thiazine-eosin**.

# Normal male karyotyping ( 22+XY)

