

## 6.1 Surface markings of the main non-alimentary tract abdominal organs

Structure	Position
Liver	Upper border: fifth right intercostal space on full expiration Lower border: at the costal margin in the mid-clavicular line on full inspiration
Spleen	Underlies left ribs 9–11, posterior to the mid-axillary line
Gallbladder	At the intersection of the right lateral vertical plane and the costal margin, i.e. tip of the ninth costal cartilage
Pancreas	Neck of the pancreas lies at the level of L1; head lies below and right; tail lies above and left
Kidneys	Upper pole lies deep to the 12th rib posteriorly, 7 cm from the midline; the right is 2–3 cm lower than the left



## 8.26 Gastrointestinal (GI) 'alarm features'

- Persistent vomiting
- Dysphagia
- Fever
- Weight loss
- GI bleeding
- Anaemia
- Painless, watery, high-volume diarrhoea
- Nocturnal symptoms disturbing sleep

### 6.3 Non-alimentary causes of abdominal pain

Disorder	Clinical features
Myocardial infarction	Epigastric pain without tenderness <i>Angor animi</i> (feeling of impending death) Hypotension Cardiac arrhythmias
Dissecting aortic aneurysm	Tearing interscapular pain <i>Angor animi</i> Hypotension Asymmetry of femoral pulses
Acute vertebral collapse	Lateralised pain restricting movement Tenderness overlying involved vertebra
Cord compression	Pain on percussion of thoracic spine Hyperaesthesia at affected dermatome with sensory loss below Spinal cord signs
Pleurisy	Lateralised pain on coughing Chest signs, e.g. pleural rub
Herpes zoster	Hyperaesthesia in dermatomal distribution Vesicular eruption
Diabetic ketoacidosis	Cramp-like pain Vomiting Air hunger Tachycardia Ketotic breath
Salpingitis or tubal pregnancy	Suprapubic and iliac fossa pain, localised tenderness Nausea, vomiting Fever
Torsion of testis/ovary	Lower abdominal pain Nausea, vomiting Localised tenderness



## 1

## 8.5 Diagnosing abdominal pain

	Disorder			
	Peptic ulcer	<del>Biliary colic</del>	Acute pancreatitis	Renal colic
Site	Epigastrium	Epigastrium/right hypochondrium	Epigastrium/left hypochondrium	Loin
Onset	Gradual	Rapidly increasing	Sudden	Rapidly increasing
Character	Gnawing	Constant	Constant	<del>Constant</del>
Radiation	Into back	Below right scapula	Into back	Into genitalia and inner thigh
<b>Timing</b>				
Frequency/periodicity	Remission for weeks/months	Able to enumerate attacks	Able to enumerate attacks	Usually a discrete episode
Special times	Nocturnal and especially when hungry	<del>Unpredictable</del>	After heavy drinking	Following periods of dehydration
Duration	$\frac{1}{2}$ –2 hours	<del>4–24 hours</del>	>24 hours	4–24 hours
Exacerbating factors	Stress, spicy foods, alcohol, non-steroidal anti-inflammatory drugs (NSAIDs)	Unable to eat during bouts	Alcohol Unable to eat during bouts	
Relieving factors	Food, antacids, vomiting		Eased by <u>sitting upright</u>	
Severity	Mild to moderate	Severe	Severe	Severe

## 6.4 Typical clinical features in patients with an 'acute abdomen'

Condition	History	Examination
Acute appendicitis	Nausea, vomiting, central abdominal pain that later shifts to right iliac fossa	Fever, tenderness, guarding or palpable mass in right iliac fossa, pelvic peritonitis on rectal examination
Perforated peptic ulcer with acute peritonitis	Vomiting at onset associated with severe acute-onset abdominal pain, previous history of dyspepsia, ulcer disease, non-steroidal anti-inflammatory drugs or glucocorticoid therapy	Shallow breathing with minimal abdominal wall movement, abdominal tenderness and guarding, board-like rigidity, abdominal distension and absent bowel sounds
Acute pancreatitis	Anorexia, nausea, vomiting, constant severe epigastric pain, previous alcohol abuse/cholelithiasis	Fever, periumbilical or loin bruising, epigastric tenderness, variable guarding, reduced or absent bowel sounds
Ruptured aortic aneurysm	Sudden onset of severe, tearing back/loin/abdominal pain, hypotension and past history of vascular disease and/or high blood pressure	Shock and hypotension, pulsatile, tender, abdominal mass, asymmetrical femoral pulses
Acute mesenteric ischaemia	Anorexia, nausea, vomiting, bloody diarrhoea, constant abdominal pain, previous history of vascular disease and/or high blood pressure	Atrial fibrillation, heart failure, asymmetrical peripheral pulses, absent bowel sounds, variable tenderness and guarding
Intestinal obstruction	Colicky central abdominal pain, nausea, vomiting and constipation	Surgical scars, hernias, mass, distension, visible peristalsis, increased bowel sounds
Ruptured ectopic pregnancy	Premenopausal female, delayed or missed menstrual period, hypotension, unilateral iliac fossa pain, pleuritic shoulder-tip pain, 'prune juice'-like vaginal discharge	Suprapubic tenderness, periumbilical bruising, pain and tenderness on vaginal examination (cervical excitation), swelling/fullness in fornix on vaginal examination
Pelvic inflammatory disease	Sexually active young female, previous history of sexually transmitted infection, recent gynaecological procedure, pregnancy or use of intrauterine contraceptive device, irregular menstruation, dyspareunia, lower or central abdominal pain, backache, pleuritic right upper quadrant pain (Fitz-Hugh–Curtis syndrome)	Fever, vaginal discharge, pelvic peritonitis causing tenderness on rectal examination, right upper quadrant tenderness (perihepatitis), pain/tenderness on vaginal examination (cervical excitation), swelling/fullness in fornix on vaginal examination



# Jaundice Related Symptoms

- Appetite and weight change
- Abdominal pain, altered bowel habit
- Gastrointestinal bleeding
- Pruritus, dark urine, rigors
- Drug and alcohol history
- Past medical history (pancreatitis, biliary surgery)
- Previous jaundice or hepatitis
- Blood transfusions (hepatitis B or C)
- Family history, e.g. congenital spherocytosis, haemochromatosis
- Sexual and contact history (hepatitis B or C)
- Travel history and immunisations (hepatitis A)
- Skin tattooing (hepatitis B or C)

## 6.6 Common causes of jaundice

### Increased bilirubin production

- Haemolysis (unconjugated hyperbilirubinaemia)

### Impaired bilirubin excretion

- Congenital:
  - Gilbert's syndrome (unconjugated)
- Hepatocellular:
  - Viral hepatitis
  - Cirrhosis
  - Drugs
  - Autoimmune hepatitis
- Intrahepatic cholestasis:
  - Drugs
  - Primary biliary cirrhosis
- Extrahepatic cholestasis:
  - Gallstones
  - Cancer: pancreas, cholangiocarcinoma

## 6.7 Urine and stool analysis in jaundice

	Urine		Stools	
	Colour	Bilirubin	Urobilinogen	Colour
Unconjugated	Normal	–	++++	Normal
Hepatocellular	Dark	++	++	Normal
Obstructive	Dark	++++	–	Pale

## 6.8 Examples of drug-induced gastrointestinal conditions

Symptom	Drug
Weight gain	Oral glucocorticoids
Dyspepsia and gastrointestinal bleeding	Aspirin Non-steroidal anti-inflammatory drugs
Nausea	Many drugs, including selective serotonin reuptake inhibitor antidepressants
Diarrhoea (pseudomembranous colitis)	Antibiotics Proton pump inhibitors
Constipation	Opioids
Jaundice: hepatitis	Paracetamol (overdose) Pyrazinamide Rifampicin Isoniazid
Jaundice: cholestatic	Flucloxacillin Chlorpromazine Co-amoxiclav
Liver fibrosis	Methotrexate



## 6.9 Specific signs in the 'acute abdomen'

Sign	Disease associations	Examination
Murphy's	Acute cholecystitis: Sensitivity 50–97% Specificity 50–80%	As the patient takes a deep breath in, gently palpate in the right upper quadrant of the abdomen; the acutely inflamed gallbladder contacts the examining fingers, evoking pain with the arrest of inspiration
Rovsing's	Acute appendicitis: Sensitivity 20–70% Specificity 40–96%	Palpation in the left iliac fossa produces pain in the right iliac fossa
Iliopsoas	Retroileal appendicitis, iliopsoas abscess, perinephric abscess	Ask the patient to flex their thigh against the resistance of your hand; a painful response indicates an inflammatory process involving the right psoas muscle
Grey Turner's and Cullen's	Haemorrhagic pancreatitis, aortic rupture and ruptured ectopic pregnancy (see <a href="#">Fig. 6.25</a> )	Bleeding into the falciform ligament; bruising develops around the umbilicus (Cullen) or in the loins (Grey Turner)

### Epigastric mass

- Gastric cancer
- Pancreatic cancer
- Aortic aneurysm

### Hepatomegaly


- Palpable liver not always enlarged
- Always percuss upper border
- Palpable gallbladder

### Generalised distension

- Fat (obesity)
- Fluid (ascites)
- Flatus (obstruction/ileus)
- Faeces (constipation)
- Fetus (pregnancy)

### Right iliac fossa mass

- Caecal cancer
- Crohn's disease
- Appendix abscess



### Left upper quadrant mass


- ? Spleen:
  - Edge
  - Can't get above it
  - Moves towards right iliac fossa on inspiration
  - Dull percussion note to 9th–11th ribs mid-axillary line
  - Notch
- ? Kidney:
  - Rounded
  - Can get above it
  - Moves inferiorly on inspiration
  - Resonant to percussion above it
  - Ballotable

### Tender to palpation

- ? Peritonitis:
  - Guarding
  - Rebound
  - Absent bowel sounds
  - Rigidity
- ? Obstruction:
  - Distended
  - Tinkling bowel sounds
  - Visible peristalsis

### Left iliac fossa mass

- Sigmoid colon cancer
- Constipation
- Diverticular mass



## 8.7 Causes of dysphagia

### Oral

- Tonsillitis, glandular fever, pharyngitis, peritonsillar abscess
- Painful mouth ulcers

### Neurological

- Bulbar or pseudobulbar palsy
- Cerebrovascular accident

### Neuromuscular

- Achalasia
- Pharyngeal pouch
- Myasthenia gravis
- Oesophageal dysmotility

### Mechanical

- Oesophageal cancer
- Peptic oesophagitis
- Other benign strictures, e.g. after prolonged nasogastric intubation
- Extrinsic compression, e.g. lung cancer
- Systemic sclerosis





## 8.11 Causes of abdominal distension

<b>Factor</b>	<b>Consider</b>
<b>Fat</b>	Obesity
<b>Flatus</b>	Pseudo-obstruction, obstruction
<b>Faeces</b>	Subacute obstruction, constipation
<b>Fluid</b>	Ascites, tumours (especially ovarian), distended bladder
<b>Fetus</b>	Check date of the last menstrual period
<b>Functional</b>	Bloating, often associated with irritable bowel syndrome



# Causes of rectal bleeding

- Haemorrhoids
- Anal fissure
- Colorectal polyps
- Colorectal cancer
- Inflammatory bowel disease
- Ischaemic colitis
- Complicated diverticular disease
- Vascular malformation

## 6.11 Grading of hepatic encephalopathy (West Haven)

Stage	State of consciousness
0	No change in personality or behaviour No asterixis (flapping tremor)
1	Impaired concentration and attention span Sleep disturbance, slurred speech Euphoria or depression Asterixis present
2	Lethargy, drowsiness, apathy or aggression Disorientation, inappropriate behaviour, slurred speech
3	Confusion and disorientation, bizarre behaviour Drowsiness or stupor Asterixis usually absent
4	Comatose with no response to voice commands Minimal or absent response to painful stimuli

*Reproduced from Conn HO, Leevy CM, Vlahcevic ZR, et al. Comparison of lactulose and neomycin in the treatment of chronic portal-systemic encephalopathy. A double blind controlled trial. Gastroenterology 1977; 72(4):573, with permission from Elsevier Inc.*

## 6.12 Differentiating a palpable spleen from the left kidney

Distinguishing feature	Spleen	Kidney
Mass is smooth and regular in shape	More likely	Polycystic kidneys are bilateral irregular masses
Mass descends in inspiration	Yes, travels superficially and diagonally	Yes, moves deeply and vertically
Ability to feel deep to the mass	Yes	No
Palpable notch on the medial surface	Yes	No
Bilateral masses palpable	No	Sometimes, e.g. polycystic kidneys
Percussion resonant over the mass	No	Sometimes
Mass extends beyond the midline	Sometimes	No (except with horseshoe kidney)



## 6.10 Causes of hepatomegaly

### Chronic parenchymal liver disease

- Alcoholic liver disease
- Hepatic steatosis
- Autoimmune hepatitis
- Viral hepatitis
- Primary biliary cirrhosis

### Malignancy

- Primary hepatocellular cancer
- Secondary metastatic cancer

### Right heart failure

### Haematological disorders

- Lymphoma
- Leukaemia
- Myelofibrosis
- Polycythaemia

### Rarities

- Amyloidosis
- Sarcoidosis
- Budd–Chiari syndrome
- Glycogen storage disorders

## 6.13 Causes of splenomegaly

### Haematological disorders

- Lymphoma and lymphatic leukaemias
- Myeloproliferative diseases, polycythaemia rubra vera and myelofibrosis
- Haemolytic anaemia, congenital spherocytosis

### Portal hypertension

### Infections

- Glandular fever
- Malaria, kala-azar (leishmaniasis)
- Bacterial endocarditis
- Brucellosis, tuberculosis, salmonellosis

### Rheumatological conditions

- Rheumatoid arthritis (Felty's syndrome)
- Systemic lupus erythematosus

### Rarities

- Sarcoidosis
- Amyloidosis
- Glycogen storage disorders

## 6.14 Causes of ascites

Diagnosis	Comment
<b>Common</b> Hepatic cirrhosis with portal hypertension Intra-abdominal malignancy with peritoneal spread	Transudate Exudate, cytology may be positive
<b>Uncommon</b> Hepatic vein occlusion (Budd–Chiari syndrome) Constrictive pericarditis and right heart failure Hypoproteinaemia (nephrotic syndrome, protein-losing enteropathy) Tuberculous peritonitis Pancreatitis, pancreatic duct disruption	Transudate in acute phase Check jugular venous pressure and listen for pericardial rub Transudate Low glucose content Very high amylase content



## 6.16 Causes of abnormal stool appearance

<b>Stool appearance</b>	<b>Cause</b>
Abnormally pale	Biliary obstruction
Pale and greasy	Steatorrhoea
Black and tarry (melaena)	Bleeding from the upper gastrointestinal tract
Grey/black	Oral iron or bismuth therapy
Silvery	Steatorrhoea plus upper gastrointestinal bleeding, e.g. pancreatic cancer
Fresh blood in or on stool	Large bowel, rectal or anal bleeding
Stool mixed with pus	Infective colitis or inflammatory bowel disease
Rice-water stool (watery with mucus and cell debris)	Cholera

## Right

## Left

Gallstones  
Stomach Ulcer  
Pancreatitis

Stomach Ulcer  
Heartburn/ Indigestion  
Pancreatitis, Gallstones  
Epigastric hernia

Stomach  
Ulcer  
Duodenal  
Ulcer  
Biliary Colic  
Pancreatitis

Kidney stones  
Urine Infection  
Constipation  
Lumbar hernia

Pancreatitis  
Early Appendicitis  
Stomach Ulcer  
Inflammatory Bowel  
Small bowel  
Umbilical hernia

Kidney Stones  
Diverticular Disease  
Constipation  
Inflammatory bowel  
disease

Appendicitis  
Constipation  
Pelvic Pain (Gynae)  
Groin Pain  
(Inguinal Hernia)


Urine infection  
Appendicitis  
Diverticular disease  
Inflammatory bowel  
Pelvic pain (Gynae)

Diverticular Disease  
Pelvic pain (Gynae)  
Groin Pain  
(Inguinal Hernia)

# Serum-ascites albumin gradient (SAAG)

	SAAG (g/dL)	
	$\geq 1.1$	$< 1.1$
Total protein (g/dL)		
$< 2.5$	Cirrhosis Acute liver failure	Nephrotic syndrome
$\geq 2.5$	CHF Constrictive pericarditis Budd-Chiari syndrome Veno-occlusive disease	Peritoneal carcinomatosis TB peritonitis Pancreatic ascites Chylous ascites







## 9.2 Features of bladder outlet obstruction due to prostatic hyperplasia

- Slow flow
- Hesitancy
- Incomplete emptying (the need to pass urine again within a few minutes of micturition)
- Dribbling after micturition
- Frequency and nocturia (due to incomplete bladder emptying)
- A palpable bladder



## 9.6 Causes of urinary incontinence

- Pelvic floor weakness following childbirth
- Pelvic surgery or radiotherapy
- Detrusor overactivity
- Bladder outlet obstruction
- Urinary tract infection
- Degenerative brain diseases and stroke
- Neurological diseases, e.g. multiple sclerosis
- Spinal cord damage





## 9.11 Urinary incontinence: points to cover in the history

- Age at onset and frequency of wetting
- Occurrence during sleep (enuresis)
- Any other urinary symptoms
- Provocative factors, e.g. coughing, sneezing, exercising
- Past medical, obstetric and surgical histories
- Number of pads used. Are they damp, wet or soaked?
- Impact on daily living



## 9.7 Abnormalities of urine colour

### Orange-brown

- Conjugated bilirubin
- Rhubarb, senna
- Concentrated normal urine, e.g. very low fluid intake
- Drugs: sulfasalazine

### Red-brown

- Blood, myoglobin, free haemoglobin, porphyrins
- Beetroot, blackberries
- Drugs: rifampicin, rifabutin, clofazimine, entacapone

### Brown-black

- Conjugated bilirubin
- Drugs: L-dopa, metronidazole, nitrofurantoin, chloroquine, primaquine
- Homogentisic acid (in alkaptonuria or ochronosis)

### Blue-green

- Drugs/dyes, e.g. propofol, fluorescein, triamterene



## 9.9 Causes of proteinuria

### Renal disease

- Glomerulonephritis
- Diabetes mellitus
- Amyloidosis
- Systemic lupus erythematosus
- Drugs, e.g. gold, penicillamine
- Malignancy, e.g. myeloma
- Infection

### Non-renal disease

- Fever
- Severe exertion
- Severe hypertension
- Burns
- Heart failure
- Orthostatic proteinuria\*

\*Occurs when a patient is upright but not lying down; the first morning sample will not show proteinuria.



## 9.10 Causes of transient proteinuria

- Cold exposure
- Vigorous exercise
- Febrile illness
- Abdominal surgery
- Heart failure



## 1

## 9.12 Some hereditary and congenital conditions affecting the kidneys and urinary tract

Name	Principal findings	Commonly associated abnormalities	Most common form of inheritance
Adult polycystic kidney disease	Bilateral enlarged kidneys, sometimes massive, with nodular surface	Liver cysts Intracranial berry aneurysms Mitral or aortic valve abnormalities	Autosomal dominant
Alport's syndrome	Haematuria, proteinuria, renal failure	Nerve deafness Lens and retinal abnormalities	X-linked dominant
Medullary sponge kidney	Tubular dilatation; renal stones	Other congenital abnormalities, e.g. hemihypertrophy, cardiac valve abnormalities, Marfan's syndrome	Congenital, rarely familial
Nail-patella syndrome	Proteinuria Renal failure (30%)	Nail dysplasia, patellar dysplasia or aplasia	Autosomal dominant
Cystinosis	Tubular dysfunction; renal failure	Rickets, growth retardation, retinal depigmentation and visual impairment	Autosomal recessive
Tuberous sclerosis complex	Renal cysts Renal angioliomata	Seizures, mental retardation, facial angiofibromata, retinal lesions	Autosomal dominant
Prune-belly syndrome	Dilated bladder and urinary tract; urinary infection and renal failure	Absent abdominal wall musculature	Sporadic mutation

## 12.2 Causes of acute kidney injury

### Prerenal

- Hypovolaemia (e.g. blood loss, diarrhoea, vomiting, diuresis, inadequate oral intake)
- Relative hypovolaemia (e.g. heart failure, nephrotic syndrome)
- Sepsis
- Drugs (e.g. antihypertensives, diuretics, non-steroidal anti-inflammatory drugs)
- Renal artery stenosis or occlusion
- Hepatorenal syndrome

### Intrarenal

- Glomerular disease (e.g. systemic vasculitis, systemic lupus erythematosus, immunoglobulin A nephropathy)
- Interstitial nephritis (drug-induced)
- Acute tubular necrosis/injury (may follow a prerenal cause)
- Multiple myeloma
- Rhabdomyolysis
- Intrarenal crystal deposition (e.g. urate nephropathy, ethylene glycol poisoning)
- Thrombotic microangiopathy (e.g. haemolytic uraemic syndrome, scleroderma renal crisis)
- Accelerated-phase hypertension
- Cholesterol emboli

### Postrenal

- Renal stones (in papilla, ureter or bladder)
- Papillary necrosis
- Ureteric or bladder transitional cell carcinoma
- Intra-abdominal or pelvic malignancy (e.g. cervical carcinoma)
- Retroperitoneal fibrosis
- Blood clot
- Bladder outflow obstruction (e.g. prostatic enlargement)
- Neurogenic bladder
- Urethral stricture
- Posterior urethral valves
- Iatrogenic (e.g. ureteric damage at surgery, blocked urethral catheter)