

Epidemiology

Archive

Lecture 8

archive

Corrected By :

Mohammed
aldahamsha

1. The most common mode of transmission in Brucellosis is?

- a. Ingestion mode of transmission
- b. Air borne
- c. Contact transmission
- d. Absorption from mucus membrane
- e. Inculcation mode of transmission

answer: c

2. Brucellosis can be transmitted by all of the following modes, EXCEPT?

- a. Contact with infected placenta
- b. ingestion of raw vegetables from infected farms
- c. Ingestion of unboiled milk
- d. inhalation of infected dust or aerosol
- e. Perinatal transmission

answer: e

3. The reservoir for human Brucellosis is?

a. goat

The reservoir for human Brucellosis is?

- a. Human
- b. Cattle
- c. Rodents
- d. Cats
- e. vegetation

answer: b

4. FOR THE BRUCEILLOSIS'S ERADICATION AND CONTROL, ALL THE FOLLOWING MEASUREMENTS ARE RECOMMENDED EXCEPT ?

- a. Slaughtering the infected animals
- b. Compulsory systematic yearly vaccination of animals for 7 to 10 years
- c. Provision of a clean sanitary environment for animals
- d. Vaccination of human
- e. Sanitary disposal of urine and faeces

answer: d