



GIT – Anatomy

Archive

Lecture(8)

small intestine

Done by :

Aya Ayman

Anatomy-Lecture(8)

1. The following relations of the third part of the duodenum are true except?

- A. Posteriorly: The left ureter
- B. Anteriorly: The root of the mesentery of the small intestine
- C. Posteriorly: The inferior vena cava and the aorta
- D. Superiorly: The head of the pancreas
- E. Anteriorly: The superior mesenteric vessels

ANSWER: A

2. The correct statement regarding small intestines is:

- A. Jejunum has more plicae circularis than duodenum or ileum
- B. Peyer's patches are mainly found in ileum along antimesenteric border
- C. Tenia coli are longitudinal structures on wall of small intestines
- D. The entire small intestines have mesentery unlike large intestines
- E. Ileum wall is thicker than jejunum wall

Answer: B

Anatomy-Lecture(8)

3. Which of the following is correct regarding third part of the duodenum :

- A- Cross by superior mesenteric vessels.
- B- Runs upward and to the left to the duodenojejunal flexure.
- C- Related posteriorly to the right hilum of the kidney.
- D- Begins at the pylorus.
- E- Related superiorly to the epiploic foramen.

Answer: A

4. All of the following is a true difference between jejunum and ileum:

- A- Ileum form only one or two arcades.
- B- Jejunum is thicker walled and wider bored than ileum.
- C- The jejunal mesentery is attached below and to the right of the aorta.
- D- Ileum lies in the upper part of peritoneal cavity while jejunum in the lower part.
- E- The plicae circularis absent in the middle part of ileum.

Answer: B