




The Nervous Tissue



Nerve = Neuro....

Part II




Neuroglia

- They are supporting cells of the nervous system
- Branching cells that bind the neurons together and with blood vessels
- They can be demonstrated by
 - 1- immunohistochemical stains using antibodies against glial fibrillary acid protein
 - 2- Gold / silver impregnation technique




2



1

Types of neuroglia

1- Neuroglia in PNS




Satellite cells

Found around nerve cell bodies in Ganglia

Schwann cells

Found around axons of peripheral nerves

Types of neuroglia



2- Neuroglia in CNS

Asrtrocytes

Microglia

Oligodendrocytes

Ependymal cells


Types of Neuroglia

Central Nervous System


Ependymal cells




Oligodendrocytes



Astrocytes



Microglia




Peripheral Nervous System

Satellite cells



Schwann cells



1- Astrocytes = Macroglia

Large, star shaped cells, have multiple process each ends by foot like expansion on the surface of the blood vessels



Function:

1-Support: processes provide structural support for neurons


2- Nutrition& ion levels: through connection with B.V.

3- Metabolic function: excess neurotransmitters formed by the neurons are cleared either **by uptake by astrocytes** or **by degradation by specific enzyme** to maintain brain function & prevent overstimulation


4- Formation of blood brain barrier



Blood brain Barrier



Blood brain barrier



Blood neural barrier

Types of astrocytes

```
graph TD; A[Types of astrocytes] --> B[Protoplasmic<br/>In gray matter of<br/>brain & spinal cord]; A --> C[Fibrous<br/>in white matter]
```

Protoplasmic

In gray matter of
brain & spinal cord

Fibrous



in white matter

2- Microglia

- Small, oval cells have processes . The cell body & the processes have minute spines
- They are originate from blood monocytes i.e. member of the mononuclear phagocyte system = mesodermal in origin
- All other glial cells are ectodermal in origin

- Function:



Phagocytosis of bacteria, apoptotic and malignant cells



3- oligodendrocytes



- Small cells with few short processes (4-6) that wrap around axons in CNS (white matter where majority of axonal myelination occur) forming myelin sheath
- The processes can myelinate multiple axons (unlike schwann cells that myelinate single axon)
- Function:

Formation of myelin sheath in CNS




4- Ependymal cells

- Epithelial – like Cuboidal cell
- The apical surface have microvilli & few cilia
- Their basal surfaces have infoldings without basement membrane needed for ion transport
- Line the brain ventricles & central canal of the spinal cord



Function:

- Formation of cerebrospinal fluid
- Cilia help in circulation of CSF
- Microvilli help in absorption



Neuroglial Cell Types & Function




www.biologyexams4u.com

Central Nervous system


Peripheral Nervous system

Astrocytes



Maintain blood brain barrier
-controlling the levels of neurotransmitter around synapses,
-regulate ion, and providing metabolic support.

Ependymal cells




Line spinal cord & ventricles of the brain.
-involved in producing cerebrospinal fluid (CSF).

Oligodendrocytes




Myelinate CNS axons, provide structural framework

Microglia




Brain's immune cells
-Remove dead cells and pathogens by phagocytosis

Satellite cells



Surround neuron cell bodies in ganglia. Regulate neurotransmitter levels

Schwann cells



Myelinate neurons in PNS. maintenance and regeneration of neurons after injury

Comparison between neuron & neuroglia

Neuron	Neuroglia
Large	Small
Transmit nerve impulse	Do not transmit nerve impulse
Not able to divide	Able to divide
Form synapse	Do not form synapse

Chromatolysis


Changes occur in the neuron as a result of injury or damage to the axon

- Degeneration and permanent loss of neuron will lead to atrophy of the innervated muscle
- The feature of chromatolysis include:

1- Swelling of the neuron : the cell body becomes enlarged as a result of breakdown of r-ER & ribosomes which will cause disruption in protein synthesis

2- Loss of Nissl bodies: the Nissl bodies disperses throughout the cytoplasm leading to loss of its characteristic staining

3- Relocation of the nucleus: the nucleus move to the periphery i.e. eccentric



Regeneration of nerve fiber particular in PNS

The process involves the following:


1- Wallerian degeneration: After injury the distal part of the axon (part away fro the cell body) degenerates, clearing the way for regeneration

2- Schwan cell activation: Schwan cells proliferate and form a regeneration pathway for the axon to follow

3- Axon growth: the axon grows back toward its target, typically guided by Schwan cells and extracellular matrix proteins

4- Re-Innervation: once axon reaches its target (ms., sensory receptor) it can reestablish synaptic connections

Wallerian degeneration is essential for repair I PNS but is not effective in CNS due to several factors like lack of support from glial cells



Regeneration of nerve fiber

Thank you

