

CENTRAL NERVOUS SYSTEM

The Meninges of the Brain and Dural Venous Sinuses

Dr. Aiman Qais Afar
Surgical Anatomist



College of Medicine / University of Mutah

Monday 9 December 2024

The Meninges of the Brain

The brain in the skull and the spinal cord in the vertebral column are surrounded by three protective membranes, or meninges:



The dura mater, the arachnoid mater, and the pia mater.



Dura Mater

- ❖ The dura mater of the brain is conventionally described as **two layers**: the **endosteal layer** and the **meningeal layer**
- ❖ These are closely united except along certain lines, where they separate to form **venous sinuses**.

venous sinuses




- The **endosteal layer** is nothing more than **the periosteum** covering the inner surface of the skull bones.
- ❖ At **the foramen magnum**, it does not become continuous with the dura mater of the spinal cord.

❖ Around the **margins of all the foramina in the skull**, it becomes continuous with the periosteum on the outside of the skull bones.

❖ At the sutures, it is continuous with the sutural ligaments

MENINGES OF BRAIN: OVERVIEW

3 layers, like spinal cord:
Dura Mater – tough mother;
Arachnoid = spiderlike;
Pia Mater = tender mother;
- arrangement different: **NO EPIDURAL SPACE**



Dura Mater

Dr. Aiman Qais Afar
Monday 9 December 2024

5


□ The meningeal layer

❖ is the **dura mater proper**. It is a **dense, strong fibrous membrane** covering the brain

❖ is continuous through the foramen magnum with the dura mater of the spinal cord.


❖ It provides **tubular sheaths for the cranial nerves** as the latter pass through the foramina in the skull.

❖ Outside the skull, the sheaths fuse with the **epineurium of the nerves**



Dura Mater


- ❖ The meningeal layer sends inward **four septa**, which divide the cranial cavity into freely communicating spaces that lodge the subdivisions of the brain.
- ❖ The function of these **septa is to restrict the displacement of the brain** associated with **acceleration** and **deceleration**, when the head is moved.



❑ 1. The falx cerebri is a **sickle-shaped fold of dura mater** that lies in the midline between the **two cerebral hemispheres**

✓ Its narrow anterior end is attached to **the internal frontal crest and the crista galli.**



✓ Its broad posterior part blends in the midline with the **upper surface of the tentorium cerebelli.**



✓ The **superior sagittal sinus** runs in its upper fixed margin, **the inferior sagittal sinus** runs in its lower concave free margin, and **the straight sinus** runs along its attachment **to the tentorium cerebelli**

□ 2. The tentorium cerebelli is a **crescent-shaped fold of dura mater** that roofs over the posterior cranial fossa

□ It covers the **upper surface of the cerebellum** and **supports the occipital lobes of the cerebral hemispheres.**





□ In the anterior edge **there is a gap, the tentorial notch**, for the passage of **the midbrain** which produces an **inner free border** and an **outer attached or fixed border.**

Dura Mater

❑ The **fixed border** is attached to **the posterior clinoid processes**, the superior borders of the petrous bones, and **the margins of the grooves for the transverse sinuses on the occipital bone**.

❑ The **free border** runs forward at its two ends, crosses the attached border, and is affixed to **the anterior clinoid process** on each side.

✓ At the point where the two borders cross, the **third and fourth cranial nerves** pass forward to enter the lateral wall of **the cavernous sinus**






The **falx cerebri** and the **falx cerebelli** are attached to the upper and lower surfaces of the **tentorium**, respectively.

The **straight sinus** runs along its attachment to the **falx cerebri**,

The **superior petrosal sinus** runs along its attachment to the **petrous bone**, and



The **transverse sinus** runs along its attachment to the **occipital bone**



Dura Mater

3. The **falx cerebelli**, a **small, sickle-shaped fold of dura mater** attached to the internal occipital crest, projects forward between **the two cerebellar hemispheres**


❖ Its posterior fixed margin contains **the occipital sinus.**



Dura Mater



□ 4. The diaphragma sellae is a **small, circular fold of dura mater** that forms the roof for the sella turcica

A small opening in its center allows passage of the stalk of **the hypophysis cerebri**



Dural Venous Sinuses

- ✓ The venous sinuses of the cranial cavity are situated **between the layers of the dura mater**
- ✓ Their main function is to receive blood from the brain through **the cerebral veins** and **the cerebrospinal fluid** from **the subarachnoid space** through the **arachnoid villi**
- ✓ The blood in the dural sinuses ultimately drains into **the internal jugular veins** in the neck.





Dural Venous Sinuses

✓ The **dural sinuses** are lined by **endothelium**, and their walls are thick but **devoid of muscular tissue**.

✓ They have **no valves**.

✓ **Emissary veins**, which are also valveless, connect the dural venous sinuses with **the diploic veins** of the skull and with **the veins of the scalp**




Dural Venous Sinuses

Dr. Aiman Qais Afar
Monday 9 December 2024

15

- The superior sagittal sinus occupies the upper fixed border of the falx cerebri (It begins anteriorly at the foramen cecum, where it occasionally receives a vein from the nasal cavity). It runs posteriorly, grooving the vault of the skull;
 - ✓ Its course receives the superior cerebral veins
- At the internal occipital protuberance, it is dilated to form the confluence of the sinuses





Here, the superior sagittal sinus usually becomes continuous with the right transverse sinus; it is connected to the opposite transverse sinus and receives the occipital sinus.

Dural Venous Sinuses

Dr. Aiman Qais Afar
Monday 9 December 2024

16



- ❑ The inferior sagittal sinus occupies the free lower margin of the falx cerebri
- ❑ It runs backward and joins the great cerebral vein at the free margin of the tentorium cerebelli to form the straight sinus
- ❑ It receives a few cerebral veins from the medial surface of the cerebral hemispheres.



Dural Venous Sinuses

❑ The straight sinus occupies the line of junction of the falx cerebri with the tentorium cerebelli

❑ It is formed by the union of the inferior sagittal sinus with the great cerebral vein. It ends by turning to the left (sometimes to the right) to form the transverse sinus.





Dural Venous Sinuses

Dr. Aiman Qais Afar
Monday 9 December 2024

18

- The **transverse sinuses** are paired structures that begin at **the internal occipital protuberance**
- The **right sinus** is usually continuous with **the superior sagittal sinus**, and the **left** is continuous with **the straight sinus**.





- Each sinus occupies the attached margin of **the tentorium cerebelli**, grooving **the occipital bone** and the posteroinferior angle of the parietal bone.

Dural Venous Sinuses

The transverse sinuses receive the superior petrosal sinuses, the inferior cerebral and cerebellar veins, and the diploic veins.

They end by turning downward as the sigmoid sinuses





Dural Venous Sinuses

Dr. Aiman Qais Afar 20
Monday 9 December 2024

The superior and inferior petrosal sinuses are small sinuses situated on the superior and inferior borders of **the petrous part of the temporal bone** on each side of the skull



Each superior sinus drains the cavernous sinus into the transverse sinus, and each inferior sinus drains the cavernous sinus into the internal jugular vein



Dural Venous Sinuses

Dr. Aiman Qais Afar
Monday 9 December 2024


- ❑ The sigmoid sinuses are a direct continuation of the transverse sinuses.
- ✓ Each sinus turns **downward and medially** and grooves **the mastoid part of the temporal bone**
- ✓ The sinus then turns **forward** and then **inferiorly** through the posterior part of the **jugular foramen** to become continuous with the **superior bulb of the internal jugular vein**.



Dural Venous Sinuses

The occipital sinus is a small sinus occupying the **attached margin of the falx cerebelli**.

It commences near **the foramen magnum**, where it communicates with the **vertebral veins** and drains **into the confluence of sinuses**




Dural Venous Sinuses

23

Dr. Aiman Qais Afar
Monday 9 December 2024

- ❑ The cavernous sinuses are situated **in the middle cranial fossa** on each side of the body of the sphenoid bone
- ❖ Numerous trabeculae cross their interior, giving them a spongy appearance, hence the name.
- ❖ Each sinus extends from **the superior orbital fissure** in front to **the apex of the petrous part of the temporal bone** behind.




Arachnoid Mater

✓ The **arachnoid mater** is a delicate, impermeable membrane covering the brain and lying between the **pia mater internally** and the **dura mater externally**

✓ It is separated from the **dura** by a potential space, the **subdural space**, filled by a **film of fluid**;


✓ it is separated from the **pia** by the **subarachnoid space**, which is filled with **cerebrospinal fluid**.



✓ The outer and inner surfaces of the arachnoid are covered with flattened mesothelial cells

❖ The arachnoid bridges over the sulci on the surface of the brain, and in certain situations, **the arachnoid** and **pia** are widely separated to form **the subarachnoid cisternae**.

❖ **The cisterna cerebellomedullaris** lies between the inferior surface of the cerebellum and the roof of the fourth ventricle. (**cisterna magna**)



❖ **The cisterna interpeduncularis** lies between the two cerebral peduncles.

❖ All the cisternae are in free communication with one another and with the remainder of the subarachnoid space


In certain areas, the arachnoid projects into **the venous sinuses** to form **arachnoid villi**.

The **arachnoid villi** are most numerous along **the superior sagittal sinus**.

Aggregations of arachnoid villi are referred to as **arachnoid granulations**

Arachnoid villi serve as sites where the cerebrospinal fluid diffuses into the bloodstream.

The **arachnoid** is connected to the **pia mater** across the fluid-filled **subarachnoid space** by **delicate strands of fibrous tissue**.




Arachnoid Mater

✓ Structures passing to and from the brain to the skull or its foramina must pass through **the subarachnoid space**.

✓ All the **cerebral arteries** and **veins** lie in the space, as do the **cranial nerves**

✓ The arachnoid fuses with the epineurium of the nerves at their point of exit from the skull



✓ **The subarachnoid space** extends around **the optic nerve** as far as the eyeball.


Arachnoid Mater

- ❖ The cerebrospinal fluid is produced by the choroid plexuses within the lateral, third, and fourth ventricles of the brain.
- ❖ It escapes from the ventricular system of the brain through the three foramina in the roof of the fourth ventricle and so enters the subarachnoid space.



- It now circulates both upward over the surfaces of the cerebral hemispheres and downward around the spinal cord.

- The spinal subarachnoid space extends down as far as the second sacral vertebra



- Eventually, the fluid enters the bloodstream by passing into the arachnoid villi and diffusing through their walls.



- ❖ The pia mater **is a vascular membrane** covered by flattened mesothelial cells. It closely invests the brain, covering the gyri and descending into the deepest sulci
- ❖ It extends out over the cranial nerves and fuses with their epineurium.
The cerebral arteries entering the substance of the brain carry a sheath of pia with them.



- ❖ The pia mater forms **the tela choroidea** of the roof of the third and fourth ventricles of the brain,
- ❖ and it fuses with the ependyma to form **the choroid plexuses** in the **lateral, third, and fourth ventricles** of the brain.



Dr. Aiman Qais Afar
Surgical Anatomist
Monday 9 December 2024

