

shape
color
size
painful?
itchy?
burning?

associated?
derma syst.

Healing?

Complication?

Try Skin Lesions

جع نايم ::
color

Flat

< 1cm

macule



well circumscribed

> 1cm

patch



جع فuid

Fluid collection

< 1cm

vesicle



> 1cm

bulla

Pus

pustule

Solid

< 1cm

papule



> 1cm

nodule



Wheal

transient smooth
papule or plaque



Petechiae

flat bleeding under skin
< 2mm don't discolor on pressing.



Purpura

red, pink, purple patches
> 2mm also under m.m.

جع دار و حادة
Plaque



> 1cm
flat topped lesion.

rubioxa = Measles

المحة

diagnosed Clinically (para)

3Cs #
cough
coryza (runny nose)
conjunctivitis


VitA 5days
↓ symptoms
↓ early comp.
↓ rate of remote comp.

Comp → pneumonia

- Giant cell pneumonitis
- Super infection Staph
- SSPE


(bad)
(1 year disease)

Significant
measles
German measles
vaccine




German / 3rd

Rubella



Chicken pox

جراحي




- highly contagious
- air borne.
- viremia → latent phase


If the mother has Chicken Pox in last week of preg or one week after delivery \rightarrow varicella zoster virus

Neonatal Chicken pox


✓ Ig
phacelia ?
other symptoms
eye brain GI




Exanthematous Diseases



Scarlet fever



Roseola infantum



HHV 6 / 7 care.

5-15

A MC (post natal)
B


6 months - 2 years ← age.

Infection results in immunotolerance.

- 3 days high grade fever
- bulging ant. Fontanelle

- Main cause of febrile seizure.
- simple.
- Normal CNS.

Erythema infectiosum



IP → healthy → benign
hemolytic anemia
↓ aplastic crisis
(Fevers / lethargy / malaise / headache / GI / RSV)
fetuses & Hb
Transfusion

spreads → RS secretions
vertical.