

Glycogen Metabolism


Glycogen

- Large, branched polysaccharide, available, storage form of glucose ([Glc]↓- degradation, [Glc]↑- synthesis)


Functions:

- Liver (5% = 90g) → blood glucose conc. maintenance
- Muscle (0.7% = 245g) → source of ATP
- Enzymes for glycogen biosynthesis and degradation are permanently and firmly bound in glycogen granules


**Glycogen granules
in hepatocytes**



Two basic types of glycosidic bonds in glycogen




α -1,4 - glycosidic bond




α -1,6 - glycosidic bond

Glycogen structure

- Glucose units linked by α -1,4 glycosidic bonds (linear molecule), while in branching points α -1,6 bonds (~10:1)
- Non-reducing ends - DEGRADATION!!!
- ONLY 1 reducing end, but permanently bound to Glycogenin - self- glucosylating



- Glycogen is more branched structure than amylopectin
- More soluble and more easy to degrade (nonreducing ends!!!)
- Starch is consisted of:
 - Amylose - linear molecule, α -1,4 glycosidic bonds
 - Amylopectin – α -1,4 and α -1,6 glycosidic bonds
- Cellulose – β -1,4 glycosidic bonds
- Humans lack **β - glucosidase** for cellulose degradation



Glycogenesis versus Glycogenolysis

- Different reaction pathways and Hormonal regulation
- Regulate glucose blood **concentration (liver)**
- Provide glucose **reserve for muscle work**



Glycogenesis

- It takes place in virtually all animal tissues, but especially prominent in the **liver and skeletal muscles**
- **3 enzymes:**
 - 1. Glycogenin** – self-glucosylating enzyme
 - Synthesis of a primer- first 8 glucose molecules
 - 2. Glycogen synthase**
 - Further extension the primer by adding Glc molecules
 - Formation of α -1,4 glycosidic bonds
 - Substrate for the synthesis is UDP-glucose
 - 3. Branching enzyme** [glycosyl(4 \rightarrow 6)-transferase]
 - Formation of α -1,6glycosidic bonds

1- After entering the cell, glucose is phosphorylated by the activity of hexokinase I and II (glucokinase) forming glucose 6-phosphate



2- Glucose 6-phosphate isomerization (reversible reaction) into glucose 1-P by phosphoglucomutase

- When higher amount of glucose 6-phosphate is present in the cell, the equilibrium of the reactions is shifted to the left, towards the formation of glucose 1-phosphate





3- **UDP glucose formation** (uridine diphosphate -glucose)

- UDP-glucose is formed by the activity of **UDP-glucose pyrophosphorylase**
- The synthesis of UDP-glucose is driven by the essentially irreversible hydrolysis of pyrophosphate catalyzed by pyrophosphatase (many biosynthetic reactions are driven by the hydrolysis of pyrophosphate)



- UDP-glucose is activated form of glucose
- Anomeric carbon of a sugar is activated by attachment to a nucleotide through a **phosphate ester** linkage
- Sugar nucleotides are the substrates for polymerization of monosaccharides into disaccharides, glycogen, starch, cellulose, and more complex extracellular polysaccharides




Sucrose: UDP-Glc + Fru-6-P



Lactose: UDP-Glc + UDP-Gal


Glucuronides: UDP-Glc \longrightarrow UDP-GA (UDP- glucuronate)

- **Glycogen synthase** cannot synthesize glycogen **de novo**
- **Glycogenin** starts glycogen synthesis
- Functions: bonding of 1st molecule UDP-glucose (with UDP release), and oligomerisation of the following 7 molecules of glucose
- Glycogen synthase can act (by adding Glu units) only upon existing oligosaccharride chain containing at least **8 glucose** residues and reducing end of glycogen is permanently bond to glycogenin (self-glucosylating enzyme)




glycogen synthase transfers the glucose residue of UDP-glucose to the glycogen non-reducing end to make a new **PRIMER** ($\alpha 1,4$) linkage






Glycogen branching - formation of α -1,6 bond

- Branching enzyme [glycosyl-(4 \rightarrow 6) transferase] transfer of an oligosaccharide chain and formation of a new α -1,6 glycosidic bond, forming a new branch point.



- Some athletes consume large amounts of carbohydrates after training (carbohydrate loading) \rightarrow rapid glycogen synthesis and faster recovery
- The consumption of high-glycemic carbohydrates soon after exercise can maximize and sustain the rate of glycogen synthesis to help speed glycogen restoration.

Overall glycogenesis reactions



- If the starting substrate is Glu 6-P, 1 ATP is spent to store 1 Glu molecule (for UTP regeneration)
- If the starting substrate is glucose, 2 ATPs are needed (for Glu phosphorylation and UTP regeneration)

Glycogenolysis

- 3 enzymes involved:

1. **Glycogen-phosphorylase** hydrolyses α -1,4 bonds forming glucose-1-phosphate




- Co-enzyme is **PLP** derived from pyridoxine (vitamin B6)

2. **Debranching enzyme** which has 2 activities

a) Transferase - transfer of 3 glucose residues


b) Glucosidase - hydrolysis of α -1,6 glycosidic bond

3. **Phosphoglucomutase** transfers glucose-1-phosphate into glucose-6-phosphate




1) PHOSPHOROLYSIS

P_i ↓ glycogen phosphorylase






Glycogenolysis



- phosphorolytic breaking of α -(1,4)-glycosidic bond, except for **4 glucose residues** away from the branching site


- formation of **glucose 1-phosphate**



- transference of **3 glucose residues** from one branch and formation of α -(1,4)-glycosidic bond on the other branch

- hydrolysis of α -(1,6)-glycosidic bond with **glucose** formation

- products:
glucose 1-phosphate and **glucose**
in ratio **10 : 1**




Glycogen
(n residues)


α -1,6-Glucosidase
(Debranching enzyme)

\downarrow H_2O

2.b) HYDROLYSIS




Glucose



Glycogen
($n - 1$ residues)

3. phosphoglucomutase

- isomerisation of **glucose 1-P** into **glucose 6-P**



- **phosphoglucomutase (phosphoenzyme!)** catalyses the reaction in the direction of **glucose 6-P** formation, since the **glucose 1-P** concentration in the cell is much higher than of **glucose 6-P**

Glucose 6-phosphate


- different roles:

1. **Muscle, brain**- fuel

for aerobic and
anaerobic
metabolism
(**pyruvate**,
lactate)


2. **Liver, kidneys** -

transformation of
G 6-P into **glucose**
for other tissues by
glucose
6-phosphatase
(other tissues do
not have glucose
6-phosphatase!)



Glucose 6-phosphatase hydrolysis glucose 6-phosphate

- **liver, kidneys** - elevation of glucose blood concentration
- occurs in the lumen of **endoplasmatic reticulum (ER)** - separated from cytosol (glycolysis!)



- genetic defects in either **glucose 6-phosphatase** or **T1 transporter** lead to serious derangement of glycogen metabolism, resulting in **type Ia glycogen storage disease**

Glycogenolysis (summary)

- Rhoosphorolytic cleavage of glycogen from the non-reducing end
- Released glucose is phosphorylated and thus ready to join the metabolism without ATP cost
- Glucose 1-phosphate cannot diffuse out of the cell
- Tissues which primarily use glucose as the energy source, do not contain glucose 6-phosphatase, but use G-6-P as fuel for glycolysis

Regulation of Glycogen Synthesis and Degradation

Importance of maintaining blood glucose levels.

- Glycogen storage form in liver and muscle.

- In liver:

Glycogen synthesis during periods well fed state.

Glycogen degradation during periods of fasting.

- In skeletal muscle:

Glycogen degradation occurs during active exercise, activated by increase AMP and calcium calmodulin

Synthesis begins as soon as the muscle is at rest.

- Regulation of glycogen synthesis and degradation is accomplished on two levels:


- Glycogen synthase and phosphorylase are: allosterically controlled

- Glycogen synthase and phosphorylase are: hormonally regulated.

- The regulation of glycogen synthesis and degradation is extremely complex, involving many enzymes: protein kinases and phosphatases

A. Covalent modification:

- Glycogen synthase is the key enzyme, present in two form:
Glycogen synthase a (active form) which is dephosphorylated.
Glycogen synthase b (inactive form) which is phosphorylated.



B. Induction and repression of the key enzyme:

- In well fed state: induce insulin synthesis for the key enzyme (induction) so, glycogenesis is stimulated.
- In fasting: decrease insulin leading to decrease synthesis of the key enzyme (repression) and hence glycogenesis is inhibited.

C. Allosteric regulation

Glycogen synthase is:

- allosterically activated by glucose-6-P.
- allosterically inhibited by glycogen molecule.

Regulation of Glycogenolysis:

Phosphorylase is the key enzyme

A. Covalent modification:


- It is present in two forms:

Phosphorylase “a” which is phosphorylated active form.

Phosphorylase “b” which is dephosphorylated inactive form

B- Induction and repression of phosphorylase enzyme.

- In well fed state : induce insulin which leads to decrease synthesis of key enzyme (repression) so glycogenolysis is inhibited.
- Fasting decrease insulin which increase synthesis of key enzyme (induction) so glycogenolysis is stimulated.



C. Allosteric regulation:

Muscle phosphorylase is:

- Allosterically activated by AMP which is increase during muscular exercise.
- Allosterically inhibited by ATP and G-6-P

TABLE 1 Glycogen Storage Diseases of Humans

| <i>Type (name)</i> | <i>Enzyme affected</i> | <i>Primary organ affected</i> | <i>Symptoms</i> |
|--------------------------------|---|------------------------------------|--|
| Type 0 | Glycogen synthase | Liver | Low blood glucose, high ketone bodies, early death |
| Type Ia (von Gierke's) | Glucose 6-phosphatase | Liver | Enlarged liver, kidney failure |
| Type Ib | Microsomal glucose 6-phosphate translocase | Liver | As in Ia; also high susceptibility to bacterial infections |
| Type Ic | Microsomal P _i transporter | Liver | As in Ia |
| Type II (Pompe's) | Lysosomal glucosidase | Skeletal and cardiac muscle | Infantile form: death by age 2; juvenile form: muscle defects (myopathy); adult form: as in muscular dystrophy |
| Type IIIa (Cori's or Forbes's) | Debranching enzyme | Liver, skeletal and cardiac muscle | Enlarged liver in infants; myopathy |
| Type IIIb | Liver debranching enzyme (muscle enzyme normal) | Liver | Enlarged liver in infants |
| Type IV (Andersen's) | Branching enzyme | Liver, skeletal muscle | Enlarged liver and spleen, myoglobin in urine |
| Type V (McArdle's) | Muscle phosphorylase | Skeletal muscle | Exercise-induced cramps and pain; myoglobin in urine |
| Type VI (Hers's) | Liver phosphorylase | Liver | Enlarged liver |
| Type VII (Tarui's) | Muscle PFK-1 | Muscle, erythrocytes | As in V; also hemolytic anemia |
| Type VIb, VIII, or IX | Phosphorylase kinase | Liver, leukocytes, muscle | Enlarged liver |
| Type XI (Fanconi-Bickel) | Glucose transporter (GLUT2) | Liver | Failure to thrive, enlarged liver, rickets, kidney dysfunction |