

# *Muscle Tissues*

Quit

# MUSCLE

Specialized for **contraction** allow movement

The cells are called **fibres (myocyte)** because of their length

sarcoplasm = protoplasm




sarcolemma = cell membrane

sarcoplasmic reticulum = smooth surfaced EPR

sarcomere = functional unit

sarcosomes = mitochondria

**Types:**



Main menu

Slide menu



Quit



Muscle cells can be placed into three categories:

a. **Smooth Involuntary Muscle**

- i. found in hollow visceral organs such as the gut, uterus and blood vessels
- ii. associated with various exocrine glands.


b. **Striated Involuntary Muscle** - found in the heart (cardiac muscle)

c. **Striated Voluntary Muscle** - makes up the skeletal muscles of the body



## Skeletal Muscle

- known as **striated** or **voluntary muscle**, comprises some 40-50% of the body mass in adults
- long fibres, the average length of skeletal muscle cells in humans is about 3 cm (sartorius muscle up to 30 cm, stapedius muscle only about 1 mm). Their diameters vary from 10 to 100µm.



Main menu

Slide menu



Quit

-According to the **myoglobin** content there are:

## Red fibres (Type I fibres)

- Red muscle fibres are comparatively **thin**. Contain lots of **myoglobin**
- Many mitochondria
- **Slow twitching** (contract slower) – **tire slower**
- Found in **limbs**, **long** muscles of the **back** (long, slow contraction for erect posture). Red muscles are needed when **sustained** production of force is necessary

## White fibres (Type II fibres)

- are **thicker**, **Less** myoglobin
- **Less** mitochondria
- **Fast twitching**, contraction is fast – **tire quickly**
- Found in **extraocular** muscles, **digits** (for rapid and precise movement)
- Fast twitch fibers can be further categorized into **Type IIa** and **Type IIb** fibers.



RED MUSCLE

MIXED MUSCLE

WHITE MUSCLE


high mitochondrial content

medium mitochondrial content



low mitochondrial content



**Most muscles have all types in varying ratios**



- During embryonic development **mesodermal cells** differentiate into uninuclear **myoblasts**, which elongate and **fuse** together to form **myotubes**, which further develop into the mature muscle fibers or **myofibers**. These myofibers are the basic units of skeletal muscle
- Mature skeletal muscle cells **can't divide**




- A further cell-type, known as **satellite cells (myosatellite cells)**, may be found adjacent to the sarcolemma. These are elongated, poorly-differentiated cells that are very difficult to discern in typical preparations, but become active by **exercise** or during **repair** and **regeneration** processes after muscle injury.



# Structure of skeletal muscle:

## *Light Microscopy*

- Many nuclei - 35/mm
- Nuclei are oval - situated peripheral
- Dark and light bands lie across the fiber
- **No branching**



## Skeletal muscle




Main menu

Slide menu



Quit





Main menu

Slide menu



Quit

This is skeletal muscle. The ← ← show the peripheral nuclei of a skeletal muscle fiber. Notice the cross striations and that the fibers don't have any connections.



Main menu


Slide menu




Quit

This is a cross section through skeletal muscle. The   indicate the peripheral nuclei of skeletal muscle fibers.


**Skeletal Muscle H&E**




**Skeletal Muscle H&E  
iris diaphragm open**




**Skeletal muscle H&E**



(Ross and Romrell, 1989).



This is a drawing showing how a number of myofibrils make up a muscle fiber and how a number of fibers make up a **muscle fasciculus** (bundle). A number of these bundles make up a muscle. Notice the A,I and H bands and Z disc (line) across the myofibril.




Main menu

Slide menu




Quit

(Ross and Romrell, 1989).



This drawing shows how the myofilaments (actin + myosin) make up a myofibril. It also shows the different bands across the fibril. Drawings 1,2,3,4 show cross sections through different parts of the fibril.



Main menu

Slide menu



Quit

The average length of a **sarcomere** (functional unit) is about 2.5  $\mu\text{m}$  (contracted  $\sim 1.5 \mu\text{m}$ , stretched  $\sim 3 \mu\text{m}$ ).

**I-band** - actin filaments,

**A-band** - myosin filaments which may overlap with actin filaments,


**H-band** - zone of myosin filaments only (no overlap with actin filaments) within the A-band,

**Z-line** - zone of apposition of actin filaments belonging to two neighbouring sarcomeres (mediated by a protein called alpha-actinin),

**M-line** - band of connections between myosin filaments (mediated by proteins, e.g. myomesin, M-protein).

rat skeletal muscle

0.5 micron




## Contraction:

A - band stays the same



I - band, H - bands become narrower

Myosin heads ratchet on the actin molecule

### Relaxed



### Contracted state



Main menu


Slide menu

Muscle  
Animation



Quit

Notice how the I band changes during contraction. The next 2 slides will show how the I band changes during contraction. To see it go forwards and backwards.




Main menu

Slide menu



Quit



Main menu

Slide menu





Quit

# Electron Microscopy

Two types of myofilaments

## Actin

- The actin molecule has 3 components:
  - actin monomers
  - tropomyosin - 7 actin molecules long
  - troponin



Main menu

Slide menu




Quit

## Electron Microscopy


Two types of myofilaments

### Actin

- actin monomers form 2 threads that spiral
- tropomyosin - lie in the groove of the spiral
- troponin - attach every 40 nm
- one end attach to the Z line
- other end goes to the middle of the sarcomere
- Z line consists of  $\alpha$  actinin



(Junqueira et al, 1986).



Main menu



Slide menu



Quit

## **Myosin:**

- 15 nm  $\phi$
- 1,6  $\mu\text{m}$  long
- The molecule has a head and a tail
- tails are parallel
- heads project in a spiral
- in the middle is a thickening



Main menu



Slide menu



Quit

# Titin (Connectin)

- titin is the **third** most abundant protein in muscle after myosin and actin
- its length ~27,000 to ~33,000 amino acids (depending on the splice isoform), titin is the **largest** known protein in mammals (greater than 1  $\mu\text{m}$  in length)
- located between the **myosin thick** filament and the **Z line**
- **extends** from the **Z-line** to the **M-line**.
- **Two** titin molecules extends from each half of thick filaments to Z line == **four** titin molecules for each thick filaments and Z line
- important in the **contraction**, functions as a molecular **spring** / **elastic** properties,
- (1) to **stabilize** the thick filament, (2) **center** it between the thin filaments, (3) prevent **overstretching** of the sarcomere, and (4) to **recoil** the sarcomere like a spring after it is stretched
- keep the filaments of the contractile apparatus in **alignment** and to the **passive stretch** resistance of muscle fibres.




## Sarcolemma:


- 9 nm thick
- invaginate to form T-tubule
- myofibrils - attach to the sarcolemma

## Sarcoplasmic Reticulum:

- specialized smooth EPR, regulates muscle contraction
- *Consists of T-tubules, terminal cisternae and sarcotubules*
- It is speculated that there are gap junctions between the T-tubule and terminal cisterna
- An impulse is carried into the fiber by the T-tubule from where it goes to the rest of the sarcoplasmic reticulum



(Ross and Romrell, 1989).




Main menu

Slide menu



Quit

(Ross and Romrell, 1989).




**Identify:**

T tubule,  
Terminal cisternae and  
Sarcotubules

The level at which the  
T tubule lies

Sarco tubules





Main menu

Slide menu





Quit



## Nerves: (motor)

The axon of a motor neuron branches and ends in **motor end plates (myoneural junction)** on the fiber

The excitatory transmitter at the motor end plate is **acetylcholine**




Main menu


Slide menu



Quit




Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.




**Nerves:** sensory

- Specialized fusiform sensory organ called **spindles** ( function as **stretch receptors**) form sensory receptors in muscles telling the brain how far the muscle has stretched




A number of small specialised **intrafusal muscle fibres** (nuclear bag fibres and nuclear chain fibres) are surrounded by a capsule of connective tissue.

**Stretch receptor**



(Ross and Romrell, 1989).





Main menu


Slide menu



Quit



This is the motor end plate. Slide 1 shows a low magnification. The ↓↓ indicate 2 motor end plates. The → in slide 2 shows where the myelin sheath ends. Slide 3 shows a single motor end plate.



Main menu

Slide menu





Quit

## Connective tissue coverings of the muscle

- Endomysium around fibres, perimysium around bundles (fascicle) and epimysium around the whole muscle
- blood vessels and nerves lie in these connective tissue coverings.
- The CT goes over into a tendon or aponeurosis which attaches to the periosteum

(Junqueira et al, 1986).




Main menu

Slide menu






Quit



## Tendon/ Ligament / Fasciae

- Tendon connect muscle to bone
- Ligaments join bone to bone
- Fasciae connect muscles to muscles and soft tissues.

All made of collagen, present in different orientations



## Tendon




- **dense regular** connective tissue fascicles encased in **dense irregular** connective tissue **sheaths**.

- Normal healthy tendons are composed mostly of **parallel** arrays of **collagen** fibers closely packed together// **fibroblasts** between fibers



- The dry mass = **86% collagen** (98% type I ), 2% elastin, 1–5% proteoglycans, and 0.2% inorganic

- **Aponeurosis**, are layers of flat **broad tendons**. They have a shiny, whitish-silvery color, are histologically similar to tendons, with limited blood supply e.g.

- Anterior abdominal aponeuroses
- Posterior lumbar aponeuroses



# Tendon structure



# Ligament

- band of **dense regular connective** tissue bundles made of collagenous fibers
- protected by **dense irregular connective** tissue sheaths.
- Ligaments connect bones to other bones to form joints // ligaments **limit** the mobility of articulations // or prevent certain movements
- Have **more** elastic fibres and more **ground** substances than tendon
- More **weaving** pattern and more **random** than tendon
- non-parallel collagen arrangement, aligned in direction of **imposed** stress

