




The lymphatic system (Part I)



Professor Dr Hala El-mazar



The lymphatic system



Lymphatic
vessels

Lymphatic
tissues & organs

Immunity: is body's ability to resist or eliminate potentially harmful foreign materials or abnormal cells

- Includes the following activities:
 - Defense against invading pathogens (viruses & bacteria)
 - Removal of 'worn-out' cells (e.g., old RBCs) & tissue debris (e.g., from injury or disease)
 - Identification & destruction of abnormal or mutant cells (primary defense against cancer)
 - Rejection of 'foreign' cells (e.g., organ transplant)
 - Other responses:
 - **Allergies - response to normally harmless substances**
 - **Autoimmune diseases**

The immune system

The immune system has 2 components:

- **The innate immune system** : non-specific, acts rapidly & has no immunological memory

its contents are:

physical : skin barrier , chemical : Complement proteins C1 –C9, acid in stomach, cellular : mast cells, eosinophils, neutrophils, macrophages, & natural killer cells

- **The adaptive immune system**: specific, last long, able to distinguish self from non-self, has memory, specificity & diversity


Its contents are : T & B lymphocytes & APCs

They communicate with each other through signaling molecules called cytokines & cell surface markers

The Adaptive immune system functions to defend the body by:

- Humoral immunity B cells → Against antigens → production of antibodies
- Cell mediated immunity T cells → Against tumor, transplant cells, virus infected cells & microorganisms

immune response



The structure of the Lymphatic tissue

The basic structure of the lymphatic tissue is mainly

lymphocytes (T & B), other cells also found such as

plasma cells & macrophages.

Pluripotent hemopoietic stem cell



Lymphoid progenitor cell



Lymphoblast




Prolymphocytes: have one of three different fates:



- Remain in **bone marrow** & give **B lymphocytes**
- Migrate to **thymus** and give **T lymphocytes**
- Give rise to **NK cells** which enter **blood directly**

Bone marrow



The lymphatic tissue

The lymphatic tissue present in 2 forms:


- Diffuse lymphatic tissue

No capsule present, scattered cells
Found in CT of almost all organs



- Nodular lymphatic tissue


No capsule present
Oval-shaped masses
Found single or in groups




The diffuse and /or nodular forms found in the lymphatic

organs:

- Bone marrow : diffuse form only
- Thymus: diffuse only
- Lymph node
- Tonsils
- Spleen
- MALT mucosa associated lymphatic tissue





LAYERS OF GASTROINTESTINAL WALL




Diffuse lymphatic tissue

- Lymphocytes in lamina propria & submucosa of many organs (RS, GIT, UT, RT)
- Also called mucosa associated lymphatic tissue (MALT)
- Appear as scattered dark stained nuclei within C.T.





**Nodular
Lymphatic tissue
(follicle)**



Primary
No germinal
center

Secondary
With germinal
center

**Germinal
center
(GC)**

**Mantle zone
(M)**


**Peripheral
zone
(P)**

Primary Lymphatic nodules

- Non capsulated collection of lymphocyte
- Found in all lymphoid organs EXCEPT **thymus** & **bone marrow**.

Primary nodule: has no germinal center **Only small B lymphocytes (not activated)**

Primary
lymphatic
nodule





The Secondary lymphatic nodule

contains :



1- Pale germinal center: B lymphocytes actively divide as a result of Ag stimulation , plasma cells & dendritic cells

2- Mantle zone (corona): formed by dense population of resting & memory B lymphocytes (**Mantel cell lymphoma**)

3- Peripheral zone: small B lymphocytes



Activation of B cells & development of germinal center:



Lymphatic organs

**Primary
(central)
organs**

**Secondary
(peripheral)
organs**

Thymus

**Bone
marrow**

**Lymph
nodes**

Spleen

Tonsils

MALT

Primary Lymphoid Organs

- B & T lymphocytes arise from same stem cell in bone marrow
- are **initial** “education centers” of the immune system
- In these organs, lymphocytes (**T /thymus, B/bone marrow**) differentiate into **immunocompetent cells**



(i.e. they can recognize “self” vs. “non-self”)
- This differentiation is said to be **antigen-independent**
- **The lymphocytes then enter the blood & lymph to reside in the 2nry lymphatic organs**

Secondary Lymphoid Organs

- The lymph nodes, MALT, tonsils, spleen
 - Are **secondary** “education centers” of the immune system, where **most immune response occurs**
 - In these organs, immuno-competent lymphocytes differentiate into **immune effectors & memory cells**
- (The activation and proliferation is **antigen-dependent**)


These lymphocytes then carry out their functions

Lymph nodes




- Principal **2ry lymphoid organs** of the body


- Found along lymphatic vessels



- Oval or bean shaped /encapsulated organs

- Have convex surface where afferent lymphatic's enter the node & concave surface where efferent lymphatic's, arteries &veins exit the node







A- Stroma

Capsule: may contain smooth ms., thickened at hilum

Septa (Trabeulae): extend from capsule and divide cortex into compartments

Reticular network: of reticular fibers form the background of the organ to support the parenchyma





B- Parenchyma

Is divided into 3 parts:

cortex,

paracortex ,


medulla

1- Cortex: outer zone under the capsule contains:

➤ **A- lymphatic nodules (1ry & 2ry)**

1ry: mainly B cells, APCs, reticular cells


2ry: activated B cells, macrophages, Plasma cells



➤ **B- lymphatic sinuses (subcapsular & cortical):** are spaces contains : **lymph**, B Lymphocytes, macrophages, few T-lymphocytes)

2- Paracortex:


- between the cortex and medulla
- Is called the **Thymus dependent zone** of







the lymph node, contains T cells that have migrated from the thymus [**T lymphocytes + High endothelial venules (HEV)**]

High endothelial venules(HEV): is a post- capillary venule

- is the point of entry of T cells from blood to lymph node
- its endothelial lining is unusual
- is cuboidal to facilitate movement of T cells into LN







3- Medulla: contains

➤ Medullary cords:

- * Cords of aggregated cells
- * Contains: B lymphocytes, Plasma cells , macrophages

➤ Medullary sinuses:

- Dilates spaces, continuous e cortical sinuses, & contains lymph, B cells, macrophages, they join at hilum → efferent lymph vessels




Medullary cords

Medullary sinus

Flow of lymph:

Flows from Afferent lymphatic (valves) → lymph node


- → **subcapsular sinus**
(contains B lymphocytes, macrophages & dendritic cells)
- → **cortical sinuses**
(contains B cells)
- → **paracortex**
(contains T cells)
- → **medullary sinuses**
(B cells & plasma cells)
- → **hilum** → Efferent lymphatic



Functions of lymph nodes:


- 1- **Filtration** of lymph from microorganisms & particles before it reaches the general circulation.
- 2- **Promote interaction** of the circulating antigens in lymph with lymphocytes to initiate immune response (*antigen – dependent differentiation*)
- 3- **Activation, proliferation of B lymphocytes** and antibody production.
- 4- **Activation T lymphocytes into cytotoxic T cells**

Lymph and blood supply of Lymph Node



Tonsils

Masses of Lymphoid tissue **at entrance** of digestive and respiratory **under oral or respiratory epithelium** produce lymphocytes to guard against infections




Characteristics of its lymphoid tissue:

- Covered by epithelium.
- Not situated along course of lymphatic vessels



Tonsils



Palatine
Non keratinized
stratified
squamous epi



Pharyngeal
Pseudo-
stratified
Col. Ciliated



Lingual
Non keratinized
stratified
Squamous

Palatine tonsils

2 tonsils located in the oral part of pharynx.

- **Crypt**: Epithelial invaginations into the tonsil substance lined by surface epithelium.
- **Stratified squamous epith**: Covers the free surface of the tonsil and lines the **crypts**.
- **Lymphoid tissue**: diffuse + nodular lymphatic tissue. May contain germinal centers.






Palatine Tonsile


Pharyngeal tonsil

- **Single** mass of lymphoid T. in nasopharynx
- Covered by pseudo-st. columnar ciliated e goblet cells (**respiratory epithelium**)
- It has **No crypts**, underlying capsule is thin




Lingual tonsil



- The posterior 1/3 human tongue
- Covered e non - k. stratified squamous epith.
- **Contains crypts**, mucus glands at the root of tongue drain through several ducts into the crypts
- Tensile contains lymphoid nodules + diffuse lymphocytes.



Palatine Tonsil w/ SS non-K (arrow)



Pharyngeal Tonsil w/ Resp epith (arrow = nodule)



Thank you

