

# ***THORACIC DUCT & THE SPLEEN***

***Dr. Aiman Q. Afar***  
***Surgical Anatomist***

***College of Medicine / University of Mutah***  
***2024-2025***



***Wednesday 26 March 2025***

# Thoracic Duct

- It is the largest lymphatic vessels in the body.

**\*\* Beginning:** from the upper end of **the cisterna chyli**. (Lies anterior to bodies of L1,L2 vertebrae between the right crus of the diaphragm and the aorta)



**\*\* Shape:** It is a thin-walled vessel which **has a beaded appearance** due to presence of many valves.



# Thoracic Duct

## \*\* Course and relations:

1- It enters the thorax through **the aortic opening** of the diaphragm between **the aorta** (on the left) and **azygos vein** (on the right).




# Thoracic Duct

**\*\* Course and relations:**

**2-** In the posterior mediastinum, it ascends between **the aorta** (on the left) and **azygos vein** (on the right).

❖ Behind **right border of esophagus**.

❖ In front of the **vertebral column**, posterior intercostal arteries, and hemiazygos veins.



# Thoracic Duct


**\*\* Course and relations:**

**3- At the level of T5:**

- It crosses the median plane from right to left behind the esophagus.

**4- In the superior mediastinum:**

- It ascends behind left border of esophagus.





# Thoracic Duct

## \*\* Course and relations:

5- In the neck, at the level of the C6, it curved behind the carotid sheath.

6- Finally, it descends to end into the junction of the left subclavian and left internal jugular veins.

N.B; At the end of the duct there are 2 valves to prevent regurgitation of the blood.




# Thoracic Duct

## \*\* Tributaries:

1) **Cisterna chili**, receives lymphatic from the lower part of the body through;

**A- Intestinal lymph trunk**  
(from abdomen and pelvis).

**B- Right and left lumbar lymph trunks** (from 2 lower limbs).




# Thoracic Duct

## \*\* Tributaries:

2) Left broncho-mediastinal lymph trunk: drains the **left 1/2 of the thoracic cavity**.

3) Left subclavian trunk: drains **the left upper limb**.




4) Left jugular lymph trunk: drains **the left 1/2 of the head and neck**.



# Thoracic Duct

## \*\* Tributaries:

- 5) Efferent from the posterior mediastinal lymph nodes.
- 6) Efferent from the posterior intercostal lymph nodes.



# Spleen

- It is a **lymphatic organ** connected to the vascular system.

**\*\* Position:** It lies in the **left hypochondrium**


N.B.; - always remember the odd numbers **1, 3, 5, 7, 9, and 11.**

**1** inch thick, **3** inch broad, **5** inch long, **7 ounces (200 gm)** weight & lies between **9 & 11** ribs.

**\*\* Functions:**

1- Storage of blood.


2- Destruction of old RBCs.



# Spleen

## \*\* Surface anatomy:

- ❖ The long axis of the spleen lies along the long axis of the 10th rib.
- ❖ Its upper border is parallel to the superior border of the 9th rib.
- ❖ Its lower border is parallel to the inferior border of the 11th rib.
- ❖ The anterior end normally lies just behind the left midaxillary line.
- ❖ The posterior end lies **one and half inches** lateral to the 10th thoracic spine.




**N.B; the normal spleen is not palpable.**

- If the spleen is **felt below the costal margin**, it is **enlarged at least 3 times** of its normal size.

**\*\* How to place the spleen in the correct anatomical position**

1- Hold the spleen in your left hand with **its convex surface** applied to **the palm**, the **round posterior end** towards **the wrist**, the **broad anterior end** towards the **tips of fingers** and the **notched upper border** applied **to the thumb**.

2- Put your hand behind the left midaxillary line with an angle 45 degrees with the horizontal.



# Spleen

13



Dr. Aiman Al Maathidy  
Wednesday 26 March 2025

## \*\* Features of the spleen

The spleen has;

### A- 2 Ends

- 1- **Posterior end (tapering)** directed upwards, backwards and medially.
- 2- **Anterior end (broad)** directed downwards, forwards and laterally.



## B- 2 Borders

1- Upper border: sharp.



✓ It shows **one or more notches** near its anterior (lateral) end.

✓ It meets the anterior end in the angle of the spleen.

*N.B; Notching of the upper border is an indication of foetal lobulation.*

2- Lower border: thick and round.

## C- 2 Surfaces: diaphragmatic and visceral



# Spleen



- **Relations of the spleen**

- 1- **Diaphragmatic surface:** the outer smooth and convex surface.

- ❖ It is related to the diaphragm which separates it from:

- ✓ The **9th, 10th and 11th ribs** and the intercostal structures.

- ✓ **Left pleura and the left lung.**




# Spleen

**2- Visceral surface:** the inner surface of the spleen.

- ✓ It contains the hilum of the spleen.
- ✓ It is irregular as it carries 4 impressions

**A. Gastric impression:** between the upper border and the hilum.

- ✓ It is related to the posterior surface of the stomach.




**B. Renal Impression:** between the hilum and lower borders. It is related to **the anterior surface of the left kidney.**

**N.B;** **intermediate margin** is the ridge between gastric and renal impressions.



**C. Colic impression:** lies close to the anterior (lateral) end of the spleen. It is related to **the left colic flexure.**

**D. Pancreatic impression:** below the lateral end of the hilum. It is related to **the tail of pancreas.**



**E. Hilum of the spleen:** between the gastric and renal impression.

- ❖ It transmits 1- Terminal branches of the **splenic artery (5-6 branches)**.
- 2- Tributaries of **the splenic vein**.
- 3- Autonomic **nerves and lymphatic**.
- ❖ It gives attachment to **2 ligaments; lienorenal and gastro-splenic**.



# Spleen


## \*\* Ligaments (Peritoneal folds) of the spleen:

❖ The spleen is almost completely surrounded by peritoneum except at the hilum.

1- **Gastrosplenic ligament:** between hilum of the spleen and the stomach.

## \* Contents:



- 1- Short gastric vessels.
- 2- Left gastro-epiploic vessels.
- 3- Sympathetic plexus around the arteries.
- 4- Pancreatico-splenic lymph nodes.
- 5- Extraperitoneal fatty tissue.





**2- Lienorenal ligament, between the hilum of the spleen and left kidney.**

**\* Contents**

- 1- Tail of pancreas.
- 2- Splenic vessels.
- 3- Sympathetic plexus around the splenic artery.
- 4- Pancreatico-splenic lymph nodes.
- 5- Extraperitoneal fatty tissue.



**3- Phrenico-colic ligament:** It extends from the **left colic flexure** to the **diaphragm**. It supports the lateral end of the spleen. As a result of splenomegaly (enlarged spleen) extends towards the umbilicus.




# Spleen

- Blood supply

- \*\* Arterial supply:**

- Splenic artery** is the largest branches of the **coeliac trunk**. It runs as a **tortuous course** along the **upper border of pancreas**.

- It enters **the lienorenal ligament** to reach the hilum and end by dividing into **5-6 branches**. They are end-arteries.



# Spleen

## \*\* Venous drainage:



□ Splenic vein runs a straight course behind the pancreas.

- It ends by joining the superior mesenteric vein to form the portal vein.

## \* Lymphatic drainage:

- To the pancreatico-splenic lymph nodes.

- Red bulb of the spleen has no lymphatic.





**Dr. Aiman Al Maathidy**  
**Wednesday 26 March 2025**