

antineoplastic drug 2

2025-4-16

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

* microtubul inhibitors :-

Vinca alkaloid

Cell cycle Specific

* bind to tubular protein (tubulin)

↳ lead to

depolymerization

* prevent formation of mitotic spindle

- 16) Vinca alkaloids exert antitumor activity by?
- a. Activating topoisomerase II to cause breaks in DNA strands.
 - b. Crosslinking DNA strands.
 - c. Inhibiting DNA mediated RNA synthesis.
 - d. Inhibiting polymerization of tubulin to form intracellular microtubules.
 - e. None of the selection.

Taxanes

✓ polymerization

but prevent separation of chromosomes

use in :-

Breast, Ovarian Cancer

paclitaxel

Docetaxel

- 6) One of these paired cytotoxic agents have similar mechanism of action?
- a. Vincristine and paclitaxel.
 - b. Etoposide and paclitaxel.
 - c. Docetaxel and paclitaxel.
 - d. Docetaxel and etoposide.
 - e. Vincristine and etoposide.
- Ans : c

Vincristine.

vinblastine

not myelotoxic

toxic to Bone marrow (myelotoxic)

neurotoxic


least neurotoxic

Toxicity of BM (myelotoxic)

و مانع ذو قابل

* Topoisomerase inhibitors :-

- * inhibit (Topo..) to their function مطابق لـ Topo II
- * cause DNA breaks DNA
- * Cell Cycle specific



لها دوادل يكعون في تيكيث منع
بوقرقة وجه العريض ورعاة على الماء
وتحت خفيف العلاج

وتحت خفيف العلاج

anticancer drug ادوية الـ
L → IV يعطى

* Steroid hormones and their antagonists



Tamoxifen

* كلو مجم

estrogen antagonist

* Selective estrogen-receptor modulator

* weak estrogenic activity → (1/100 effect)
less than 1%
غير قابل ٪ ٥

inhibition tumor growth of Breast Cancer

* first line in estrogen-positive breast cancer

orally

* AE: Skin rash, vaginal bleeding
and discharge, hypercalcemia
وغيرها من الآثار الجانبية
وهي تؤثر على الغدد الصماء
Endometrial cancer, thromboembolism

* Monoclonal AB

AB against specific tumor Ag

Trastuzumab

* for epidermal

growth factor receptor

HER-2/neu

* approved IV for treatment

metastatic breast Cancer

↳ in patient whose tumor

Overexpress HER/neu

AEs

Congestive heart failure (rare)

fever, chills

Bevacizumab

* antiangiogenesis

For
vascular endothelial
growth factor (VEGF)

* approved IV → first line

drug against metastatic
Colorectal Cancer

Rituximab

For CD20⁺ B cell

* For
non-Hodgkin
lymphoma

Others :-

* platinum coordination complexes :-

platin ← مذکور بالسلایدات اعماقاً دوّرة ينتحروا بـ

platinum ← قُوْسَنْهُ فـ

platinum bind to DNA and forms CROSS links between
neighboring guanines

AE :-

nephrotoxicity.

Ototoxicity.

neurotoxicity

كما ، drug - drug interaction

Tyrosine Kinase inhibitors

Imatinib

inhibit bcr-abl tyrosin kinase

prevent the phosphorylation of tyrosin

inhibit subsequent step that lead to cell proliferation

* orally for

Chronic myelogenous leukemia

AE:

fluid retention • Edema

hepatotoxicity,

neutropenia ← esp.

Gefitinib

* for - epidermal growth factor receptor

* treatment of non-small cell lung cancer

* orally

AE:

* acne-like skin rashes

diarrhea • nausea

can manageable

- 3) Which one the following is a Tyrosine kinase inhibitor?
- a. Cisplatin.
 - b. Tamoxifen.
 - c. Irinotecan.
 - d. Gefitinib. XXX
 - e. Paclitaxel.

Wish

* كلام الدكتور

لي بي عصرين لدواه وهين حهم

الـ AE الي كز عليهم الدكتور وهمان (المذكورين في التحنيط)

اسمهاء الأدوية كلها تابعها



Done by : NOOR

أدعولي ☺