

وسهلا



أهلا

يُمنع أخذ السلايدات بدون
إذن المحرر واي اجراء
يخالف ذلك يقع تحت طائلة
المسؤولية القانونية
جميع المعلومات للاستخدام
التعليمي فقط

الأستاذ الدكتور يوسف حسين

كلية الطب - جامعة مؤتة - الأردن

دكتورة من جامعة كولونيا المانيا

اليوتيوب د. يوسف حسين

Prof. Dr. Youssef Hussein Anatomy - YouTube

الواتس 00201224904207


بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

Anatomy of Ureter

[Prof. Dr. Youssef Hussein Anatomy - YouTube](#)

- Rt Kidney → below Transpyloric
- Lt Kidney → above Transpyloric

- The ureters are 2 muscular tubes.
- They convey urine from the kidneys to the urinary bladder.
 - **** Begin**, from hilum of the kidney at transpyloric plane (L 1)
 - **** Termination**, opening into the posterosuperior angle of the urinary bladder.
 - **** Length**; It is about 25 cm long



Abdominal part

Anterior relations

Superior mesenteric vessel
in root of mesentery (cut)

Right gonadal vessel


Posterior relations

● Psoas major M

Genitofemoral N

↳ reach external genitalia
to upper part of medial side of the thigh
lumbar plexus
branches on P.M Ms.

Prof. Dr. Youssef Hussein Anatomy - YouTube



Ureter * بداية → Imp

posterior as the kidney in post.
abdominal wall.

Anterior relations

Left gonadal vessel

Sigmoid colon

↳ Store the stool
so, in Constipation
• Lt Ureter (عرق يقطع)

↓
[renal colic] التهاب الكلى
• دائما باتجاه السعال

* Varicocele in male also
mostly in the Lt side → Next Lec

Urinary bladder باتجاه الـ Forward medially بعد الـ Post

• Pelvic Part of the Ureter

- It **enters** pelvis by crossing bifurcation of common iliac artery.
- It **descends** on lateral wall of the pelvis along the internal iliac artery.
- Opposite **ischial spine**, it curves anteromedially to the angle of the urinary bladder.
 → postero superior angle

• Intramural part

- It runs oblique through urinary bladder wall **for 2 cm** before opening act as a valve like mechanism to prevent regurgitation of urine.
 ↓ as the Urinary bladder has No anatomical Sphincter.



* هيكل بطني

الـ Ureter محبوس

داخل جدار الـ bladder


و يبرز عن الكلية

بعد الـ Ureter

Common iliac artery

Internal iliac artery

Urinary bladder



• ** Relations of the pelvic part

- It crosses obturator nerve and vessel.
- **A- In the male**, It is crossed by vas deferens.
- **B- in female**,
- It forms posterior boundary of the **ovarian fossa**.
- Then runs on **lateral aspect of the cervix and upper part of the vagina** to reach the bladder.
- Here, it is crossed by uterine artery (**water under bridge**). * So in Cervixectomy keep care of Ureter.

* Relation of Ureter From Post → Ant :-



• Normal constrictions of ureter


- Where stones may be impacted

- 1- Pelviureteric junction.
- 2- Crossing the bifurcation of common iliac artery at the pelvic brim.
- 3- At ischial spine (curvature of ureter).
- ④ Intramural part (inside the wall of urinary bladder). *Most Common*

~^

* Order From narrower riskier place :-

4
3
1
2




IVP

Intravenous pyelogram

- Ureter descends downwards and slightly medially opposite the tips of transverse processes of the lumbar vertebrae.
- Then it descends in front of the ***sacroiliac joint.

Prof. Dr. Youssef Hussein Anatomy - YouTube

Blood supply of the ureter



- The ureter is identified at operation by

- a- it appears as thick muscular tube with longitudinal blood vessels.

- b- It shows peristalsis and gives urine on aspiration

- Nerve supply

- Sympathetic from T 11 to L 1.

- Parasympathetic S 2, 3, 4.

- Renal pain is referred to the groin and external genitalia which are supplied by genitofemoral nerve (L1&2).




↳ referred pain

dr_youssefhussein@yahoo.com

Anatomy of Urinary bladder

Prof. Dr. Youssef Hussein Anatomy - YouTube

- * Embry Urinary bladder → in pelvic
- * Full Urinary bladder → pelvic + abdominal cavity



**** Position:**

1- During **childhood**, it is an abdominal organ because the pelvis is narrow.


2- At **puberty**, lies in pelvic cavity.

- When the bladder is **distended**, it raises above the upper border of the symphysis pubis and becomes behind the **anterior abdominal wall**.

**** Function and capacity:**

- It is a muscular reservoir to the urine.
- The average capacity of the bladder is 400 - 500 cc.

Prof. Dr. Youssef Hussein Anatomy - YouTube




**** Peritoneal covering**

- **In female** only the **superior surface** is covered by peritoneum.
- The reflection of the peritoneum from the uterus to the superior surface of the bladder forming the **uterovesical pouch**.


Prof. Dr. Youssef Hussein Anatomy - YouTube

* Genital system of male bet- Urinary ant & GIT post
& Female (hind gut)



**** Peritoneal covering**

- **In male** the peritoneum covers the superior surface and upper part of the base.
- The reflection of the peritoneum from the rectum to the upper part of the base forming **rectovesical pouch**.




- **Relations of Superior surface In female:**

- 1) Anterior surface of uterus
- 2) Loop of ileum

- **Posterior relations**

- 1) Cervix of the uterus.
- 2) Anterior wall of the vagina.
- 3) Rectum




- **Relations of Superior surface In male:**

- 1) Sigmoid colon
- 2) Loop of ileum

- **Posterior relations**

- 1) Rectum
- 2) Rectovesical pouch

- **Posterior Relations of urinary bladder In male**



dr_youssefhussein@yahoo.com

Median umbilical ligament

Apex of Urinary bladder

Symphysis pubis

Neck of Urinary bladder

Urethra


Urinary bladder

Prostate gland

Female size *
pelvic fascia

Prof. Dr. Youssef Hussein Anatomy - YouTube

- Neck In female, rested on pelvic fascia.




• Ligaments of urinary bladder

Median umbilical ligament (obliterated urachus) if patent urine discharge from umbilicus

Two medial umbilical ligaments obliterated umbilical arteries if patent blood discharge from umbilicus

Urinary bladder



- Two lateral ligaments from the neck to the pelvic fascia

[Prof. Dr. Youssef Hussein Anatomy - YouTube](#)

Puboprostatic ligament: between body of pubis and prostate in male)

Pubovesical ligament: between body of pubis and neck of the bladder in female).

*** Arterial supply of urinary bladder:**

- 1- **Superior vesical arteries** from internal iliac artery.
- 2- **Inferior vesical artery** (**male**) or **vaginal** artery (**female**) from internal iliac artery.


*** Venous drainage:** venous plexus; drain into the internal iliac vein.

**** Lymphatic drainage:**

- 1- Mostly into the external iliac lymph nodes.
- 2- Partly Into the internal iliac, and common iliac lymph nodes.

**** Nerve supply:**

- a- **Sympathetic** from T11, 12 and L 1, 2 segments of the spinal cord.
- b- **Parasympathetic** from S 2, 3, 4. *micturition*



• Micturition (Urination)

- Is initiated by stimulating **stretch receptors in the detrusor muscle** in the bladder wall by increasing volume of urine.
- Can be assisted by contraction of the abdominal muscles, which increases the intraabdominal and pelvic pressures.

■ Involves the following processes:

1. **Sympathetic [GVE]** induce relaxation of the bladder wall and constrict the internal urethral sphincter, **inhibiting emptying**.
2. Impulses (**GVA**) arise from stretch receptors in the bladder wall and enter the spinal cord (S2–S4) then to the micturition center in the **brain**.
3. **Parasympathetic (GVE)** to the bladder musculature induce contraction of the detrusor muscle and relaxation of the internal urethral sphincter, **enhancing the micturition**.
4. **General somatic efferent (GSE)** fibers in the pudendal nerve cause voluntary relaxation of the external urethral sphincter, and the bladder begins to void.
5. **At the end of micturition**, external urethral sphincter contracts, and bulbospongiosus muscles expel the last few drops of urine from the urethra.

Clinical Notes

• Reflex Bladder – Spinal Cord Transection Above T12

- The afferent signals from the bladder wall are unable to reach the brain, and the patient will have **no awareness** of bladder filling.
- Thus, the bladder automatically empties as it fills

• Flaccid Bladder – Spinal Cord Transection Below T12

- The parasympathetic outflow to bladder is damaged. Paralysis of detrusor muscle.
- The spinal reflex does not function.
- The bladder will become abnormally distended until **overflow incontinence** occurs.

Urine Retention

- **Causes:** the most common cause is prostate enlargement (BPH) In males, a stone or large blood clot.
- **Acute retention** is a medical emergency, The patient needs urinary catheter.
- **Chronic retention** leads to accumulation of residual urine in the bladder. It is often complicated by infections and formation of bladder stones

https://www.youtube.com/channel/UCVSNqbibj9UWYaJdd_cn0PQ

يُمنع أخذ السلايدات بدون
إذن المحرر واي اجراء
يخالف ذلك يقع تحت طائلة
المسؤولية القانونية
جميع المعلومات للاستخدام
التعليمي فقط

اليو تيوب د. يوسف حسين



Thank You

Question

الواتس 00201224904207

<https://www.youtube.com/@ProfDrYoussefHusseinAnatomy/playlists>