



Candidiasis and Trichomoniasis

2024-2025

Dr. Sulaiman Mahmoud Bani Abdel-Rahman

MBBS - Mutah University


MSc Medical Microbiology – University of Manchester

PhD Medical Virology - University of Manchester



Candidiasis - Introduction

- Candida infections encompass a broad spectrum of clinical syndromes caused by yeasts of the *Candida* genus, most notably *C. albicans*. *Candida* can develop as:
 - Budding yeasts.
 - Pseudohyphae (image next slide): filament-like structures, resulting from the elongation of budding yeast, with constrictions.
 - Hyphae: filamentous structures separated by septae (rather than constrictions).




Candidiasis - Introduction

NOT STD,
don't need to treat
partner

These fungi are common commensals,
residing harmlessly on human mucosal
surfaces, skin, and within the
gastrointestinal and genitourinary tracts.

However, disruptions in the local microbial
flora, damage to mucosal integrity, or
weakened immunity can allow **Candida** to
proliferate and **become pathogenic**.

normal
flora




Mucocutaneous candidiasis

Skin, mucosa, and nails

Invasive candidiasis

Candidemia and organ infiltration



Clinical manifestations

- Vulvovaginal candidiasis
- Cutaneous candidiasis (intertrigo, balanitis)
- Oropharyngeal candidiasis (thrush)
- Esophageal candidiasis
- Invasive candidiasis (disseminated infection)
- Candiduria
- Chronic mucocutaneous candidiasis



Vulvovaginal candidiasis - Introduction

- Vulvovaginal candidiasis (VVC), commonly known as a yeast infection, is an inflammatory vulvar and vaginal infection caused by overgrowth of fungal *Candida* species, most commonly *Candida albicans*.
- Although *Candida* spp. are part of the normal vaginal flora, overgrowth can lead to symptomatic infection.



Vulvovaginal candidiasis - Pathophysiology

- *Candida* fungal spp. are part of the normal vaginal flora and exist in balance with the predominant bacteria of the vaginal microbiome, gram-positive *Lactobacillus* spp.
- Alterations in immune function or in the balance of the vaginal microbiome allow for excessive *Candida* proliferation.
- Symptomatic infection develops when pseudohyphae invade the vaginal epithelium, leading to inflammation.



Vulvovaginal candidiasis - Risk factors

- **Antibiotic use** reduces the number of *Lactobacilli*, thereby disrupting the normal flora and allowing for unchecked *Candida* growth.
- **High estrogen levels** (eg, pregnancy, postmenopausal estrogen therapy) increase glycogen content (used by *Candida* spp.) within vaginal epithelial cells.
- **Diabetes mellitus** (and treatment with sodium-glucose cotransporter-2 inhibitors) results in elevated glucose, which facilitates adhesion of *Candida* to epithelial cells.
- **Immunosuppression** (ie, HIV/AIDS, corticosteroid use) impairs regulation of *Candida* spp. proliferation.



Vulvovaginal candidiasis - Clinical presentation


imp
for
case
Q.




- VVC commonly presents with:
 - **Vulvovaginal pruritus** that can lead to excoriations.
 - **Vaginal discharge**, which is classically described as *** thick, white, cottage cheese–like discharge**; however, this can vary in presentation to include scant, thin, or watery discharge.
 - **Vulvovaginal inflammation** with erythema and edema of the vulva and vaginal mucosa, leading to dysuria and dyspareunia.

Pain during
sexual
intercourse

حالة = تفكك في الجلد





Candidal vulvovaginitis

Vaginal cavity of a woman presenting with vaginal pruritus and white ~~odorless~~ vaginal discharge

White, cottage cheese-like discharge can be seen adhering loosely to the vaginal walls, which have patchy areas of inflammation.



Vulvovaginal candidiasis - Diagnosis and laboratory evaluation


- Diagnosis of VVC is often **made clinically with laboratory testing**. Physical examination often shows thick, white discharge with vulvar and vaginal erythema (figure next slide). Findings are most often confirmed using office-based tests that provide immediate results:
 - Microscopy with potassium hydroxide (KOH) wet mount preparation: The KOH lyses epithelial cells to reveal budding yeast, pseudohyphae, and hyphae (image).
 - Vaginal pH: Normal pH ranges from 4.0 to 4.5. !
 - Vaginal cultures can ~~also~~ be obtained; however, because *Candida* spp. are part of the normal vaginal flora, a positive culture does not always indicate infection.




 Overlay

Candida albicans

- Photomicrograph of fungus germination (KOH preparation, 600x)
- Mother cells (blastospores (budding yeast); green overlay) produce a hyphal outgrowth of germ tubes (budding; dashed red outline).
- Budding is suggestive of active candida infection.






Candida albicans

- Incubating the sample in human serum at 37 C for 2-3 hours (ie, **germ tube test**) leads to the formation of **true hyphae**, projections from the yeast with **no constrictions**.
- A positive germ tube test is diagnostic of *C albicans* and distinguishes it from other *Candida* species (eg, *C tropicalis*, *C glabrata*).



Candida




Pseudohyphae seen in *Candida* vaginitis, which would present with thick, white vaginal discharge and associated vaginal inflammation (eg, vaginal erythema, pruritus).



~~spread~~ → yeast
invasion → pseudo convert


Candida



- The morphology of *Candida* includes branching pseudohyphae with blastospores (budding yeast).
- Pseudohyphae, an important diagnostic clue, form when yeasts do not separate after budding, creating a sausage-like chain of elongated yeasts joined together end-to-end with constrictions.
- Incubating the sample in human serum at 37 C for 2-3 hours (ie, **germ tube test**) leads to the formation of **true hyphae**, projections from the yeast with **no constrictions**. A positive germ tube test is diagnostic of *C albicans* and distinguishes it from other *Candida* species (eg, *C tropicalis*, *C glabrata*)
- Generally, growth at 37°C at neutral pH favours pseudohyphae, whereas **hyphae are induced by the presence of serum**.




Candida



©UWorld

Candida



©UWorld



فحص ، علاج

Treatment

The mainstay of VVC treatment is with oral or topical antifungals:

- **Fluconazole**, which is the most commonly used oral treatment and generally requires only a single dose for complete symptom resolution → decrease ergosterol synthesis and disrupt fungal cellular membrane formation.
- Intravaginal topical antifungals (eg, clotrimazole, miconazole)

Vulvovaginal candidiasis	
Risk factors	• Diabetes mellitus • Immunosuppression • Pregnancy • Antibiotic use
Examination	 • Thick, cottage cheese-like discharge • Vaginal inflammation
Laboratory findings	 • Normal pH (3.8-4.5) • Pseudohyphae
Treatment	• Fluconazole



Cutaneous candidiasis (intertrigo, balanitis)

- Intertrigo consists of erythematous ("beefy red") plaques within skin folds (inguinal, inframammary, axillary).
- It is common in patients with obesity, diabetes mellitus, and immunosuppressive conditions.
- Diagnosis is typically clinical, although unclear cases can be confirmed with KOH scraping showing pseudohyphae.



not
in exam



Cutaneous candidiasis (intertrigo, balanitis)

- Balanitis is inflammation of the glans penis, often in uncircumcised boys or men.
- Thick, white discharge may be present.
- Patients with recurrent infection should be screened for diabetes mellitus.



not
in exam





Trichomonas vaginalis



Introduction

- Trichomoniasis is a common sexually transmitted infection (STI) that affects both men and women and is caused by *Trichomonas vaginalis*, a motile, flagellated protozoan.
- The majority of patients are asymptomatic, but trichomoniasis can cause vaginitis and cervicitis in women and urethritis in men.



Pathogenesis and risk factors

- *T vaginalis* is a sexually transmitted flagellated protozoan that primarily infects the squamous epithelium of the urogenital tract (eg, vagina, urethra, cervix, prostate). The adherence of *T vaginalis* to the epithelial cells causes direct cytotoxic cell damage and an inflammatory response, contributing to the symptoms of vaginitis in women and urethritis in men.
- Risk factors for trichomoniasis include high-risk sexual behaviors (similar to other STIs) and include a history of other STIs, unprotected sexual intercourse, and multiple sexual partners.



Clinical presentation

- Trichomoniasis is most often asymptomatic. However, if symptoms are present, they differ slightly between men and women.




Clinical presentation - Women



- Symptoms in women are due to cervicitis, vaginitis, and urethritis and may include:
 -  **Malodorous, frothy, greenish-yellow vaginal discharge**
 - Vulvovaginal pruritus
 - Dysuria, urinary frequency (due to infection of the urethra)
 - Dyspareunia
 - Postcoital bleeding (due to capillary hemorrhages on the cervix and vagina [ie, **"strawberry" cervix**])

bleeding after sexual intercourse



Clinical presentation - Men


- Symptoms in men are due to urethritis and may include:
 - **Clear-mucopurulent urethral discharge**
 - Dysuria
 - Prostatitis, epididymitis (rare)




Diagnosis and laboratory evaluation

- **Nucleic acid amplification test (NAAT)** is the preferred method (gold standard) of diagnosis in both men and women due its high sensitivity and specificity.
 - Swab of vaginal fluid or first-void urine testing can detect the DNA or RNA of *T vaginalis*.
- In women, the diagnosis can also be made via **wet mount microscopy**, which reveals motile, ovoid, flagellated organisms (image).
 - However, the sensitivity is lower (~50%-70%), so patients with a negative wet mount microscopy still require NAAT.


vaginal swap
+
normal
saline



- *T vaginalis* appears as motile, flagellated trophozoites;
- A wet mount of trichomoniasis would show pear-shaped organisms (flagellated trophozoites).



Video - *T vaginalis*



Management



- Treatment for trichomoniasis is required for the patient and all sexual partners, regardless of whether symptoms are present. Treatment options include:
 - **Women**
 - Metronidazole: 500 mg orally twice daily for 7 days OR
 - Tinidazole: 2 g orally in a single dose
 - **Men**
 - Metronidazole: 2 g orally in a single dose OR
 - Tinidazole: 2 g orally in a single dose





Differential diagnosis of vaginitis

إحفظه زي إسمك

Bacterial vaginosis (<i>Gardnerella vaginalis</i>)	Trichomoniasis (<i>Trichomonas vaginalis</i>)	Candida vaginitis (<i>Candida albicans</i>)
 - Thin, off-white discharge with fishy odor - No inflammation	 - Thin, yellow-green, malodorous, frothy discharge - Vaginal inflammation	 - Thick, cottage cheese discharge - Vaginal inflammation
 - pH >4.5 - Clue cells - Positive whiff test (amine odor with KOH)	 - pH >4.5 - Motile trichomonads	 - Normal pH (3.8-4.5) - Pseudohyphae
Metronidazole or clindamycin	Metronidazole; treat sexual partner	Fluconazole

KOH = potassium hydroxide.

©UWorld

