



Pharmacology of eye

PREPARED BY: HEBA AHMED HASSAN

ASSISTANT PROFESSOR OF CLINICAL PHARMACOLOGY

FACULTY OF MEDICINE, MUTAH UNIVERSITY, JORDAN

Sympathetic Innervation:

- 1- to dilator pupillae muscle radial muscles of the iris, leading to pupil dilation (mydriasis)
- 2- to blood vessels within the eye, influencing ocular blood flow and intraocular pressure
- 3- to Müller's muscle leading to eyelid retraction
- 4- Beta-2 adrenoceptors in the ciliary body increase the secretion of aqueous humor but Alpha-2 adrenoceptors in the ciliary body suppress it.

quantity not quality ← through α-Receptor.
 5- Relaxation of ciliary muscle → Accommodation of far vision
 protrusion of eye → hyperthyroidism
 quality not quantity
 miosis due to M₃ Receptor.

Parasympathetic innervation:

- 1-To constrictor pupillae muscle, narrowing the pupil in response to bright light (light reflex).
- 2-To the ciliary muscle, causing it to contract, leading to lens accommodation.

Drainage of aqueous humor:

Aqueous humor flows from the posterior chamber → anterior chamber → exits via two routes:

1- Conventional Pathway (90% of outflow):

Fluid traverses the trabecular meshwork → Schlemm's canal → episcleral veins.

space of Fontana


2-Unconventional Pathway (10% of outflow):

Fluid drains through the ciliary muscle, suprachoroidal space, and sclera (uveoscleral route).

passive	active	سبب التقيح بين ال
	dilatation (mydriasis)	Constriction (miosis)
active	stimulation of dilator	stimulation of constrictor
passive	cut the constrictor (قطع ال مقصر)	cut the dilator (قطع ال مدبر)


Drugs

- 1- Drugs affecting pupil size
- 2-Treatment of Glaucoma
- 3-Drugs that ↑↑ IOP




contraindicated with patient of glaucoma.

through M_3 Receptor.
Circular sphincter muscles contract




Bright light



Normal light

through α -Receptor.
Radial dilator muscles contract



Dim light

① To Reduce Amount of Light enter the pupil



→ through Parasympathetic stimulation.
↳ Active miosis

→ Or sympathetic inhibition
↳ passive miosis

① To increase the Amount of Light Enter the pupil



→ through sympathetic stimulation.
↳ Active mydriasis

→ Or Parasympathetic inhibition
↳ Passive mydriasis.

A- Drugs affecting pupil size

1- Miotics drugs

Drug Class	Examples	Effect on Pupil	Mechanism
Opioids (systemic) <i>→ pin-point - Pupil.</i> <i>⊗ active miosis</i>	<u>Morphine</u> , Heroin, <u>Fentanyl</u> <i>stimulation of M-Receptor (inhibitory المثبط) ←</i>	Miosis <i>severe miosis شديد انقباض الحدقة Center</i>	Activates <u>μ-opioid receptors</u> , inhibiting sympathetic tone.
<u>Cholinergic Agonists</u> (local)	Pilocarpine, Carbachol	Miosis	Stimulates parasympathetic system (muscarinic receptors).
<u>Acetylcholinesterase Inhibitors</u> (local)	Physostigmine, Neostigmine,	Miosis	Increases acetylcholine levels, activating muscarinic receptors
<u>Guanthiden</u> <i>→ passive miosis</i>	<i>treat → prostatic hyperplasia</i>	Miosis	Reduces <u>Release of NE</u> in the eye:
<u>α1-Adrenergic Blockers</u> <i>Passive miosis</i>	Prazosin, Tamsulosin	Miosis	Blocks <u>sympathetic stimulation of the dilator muscle</u>
<u>Sedatives / Barbiturates</u> <i>inhibition of sympathetic ←</i>	Benzodiazepines (high doses)	Miosis <i>Mild</i>	<u>CNS depression reduces sympathetic tone.</u> <i>فسيكوسباتية passive Miosis ←</i>

Locally acting miotics

(parasympathomimetics): stimulate M3 receptors in

N.B:-

constrictor pupillae muscle

used in treatment of glaucoma.

- 1- CPM → miosis + wide angle of filtration & space of Fontana.
- 2- Ciliary muscle → accommodation to near vision + open canal of Schlemm.
- 3- Some stimulate Nm receptors in upper eye lid → upper eyelid twitches.

1- Direct parasympathomimetics:

- Choline esters: bethanichol (M only) & carbachol (M+N).
- Alkaloid: pilocarpine (M only).

2- Indirect parasympathomimetics:

- Reversible: physostigmine (eserine) & demecarium.
- Irreversible: organophosphorus → ecothiophate & isofluorophate: Long-lasting strong effect with extreme miosis, but produce irritation & cataract

Therapeutic uses:


← لم تذكر
الكثرة

1-Glaucoma.



2-Counteract mydriatics after fundus examination.

3-Alternatively with mydriatics to cut adhesion between iris & lens.

during inflammatory Reaction.



→ iritis



Opiate use or overdose is one of the most common causes of pinpoint pupils.

Guanthidine:

Paralysis of Dilator Pupillae Muscle → miosis + ↓↓ IOP

passively.

Relaxation of levator palpebrae superioris → ↓↓ exophthalmos of hyperthyroidism.
and Muller muscle.

Morphine stimulates ^Mopiate receptor in **3rd cranial nerve nucleus** → stimulates oculomotor nerve → **ciliary ganglia (Nn)** → eye → ACh → stimulates **M3** receptors of

CPM → marked **miosis** (pin point pupil).

Pin-point pupil of morphine can be antagonized by:

- 1) Systemic naloxone** → block opiate μ receptors in CNS.
- 2) Systemic ganglion blocker** → block Nn of ciliary ganglia.
- 3) Topical or systemic atropine** → block M3 receptors on CPM

Atropine. (muscarinic)

parasympatholytic - Anticholinergic effect
 passive mydriasis

2- Mydriatics drugs

contra-indicated in cases of glaucoma.

Drug Class	Examples	Effect on Pupil	Mechanism
<u>Sympathomimetics</u> (indirect).	Epinephrine, Cocaine, Amphetamines	Mydriasis	Stimulates adrenergic receptors, enhancing sympathetic activity
<u>Anticholinergics</u> للأظننال	Atropine, Tropicamide, Scopolamine	Passive Mydriasis	Blocks parasympathetic innervation to the constrictor pupille muscle.
<u>α1-Adrenergic Agonist</u>	Phenylephrine	Mydriasis	Stimulates dilator muscle via α1 receptors.
SSRIs & SNRIs	Fluoxetine, Venlafaxine	Mydriasis	Increased serotonin activity affects autonomic control
<u>Tricyclic Antidepressants</u> ↳ Atropine-like action (Anticholinergics)	Amitriptyline, Imipramine	Mydriasis	Strong anticholinergic effects block pupil constriction.
<u>Hallucinogens</u>	LSD, MDMA بكمون بيلات	Mydriasis	Serotonin and dopamine effects increase sympathetic tone
<u>Dopaminergic Drugs</u>	Levodopa, Bromocriptine	Mydriasis	Enhances dopamine signaling, indirectly increasing sympathetic effects

A- Sympathomemetic:

N.B: ciliary muscle is predominantly supplied by parasympathetic system, to lesser extent with sympathetic supply. If Muscarinic Receptor is Blocked; most of ciliary muscle will be lost → cycloplegia → Atropine

mechanism: Stimulate α_1 receptors leading to:

Contraction of **DPM** → **Active mydriasis** (intact light reflex) & no cycloplegia, BV → VC → decongestion & ↓ IOP.

→ paralysis of ciliary muscles

○ **Examples:** Direct: phenylephrine., Indirect: amphetamine, Mixed: ephedrine.

○ **Therapeutic uses:** fundus examination especially in elderly patients liable for glaucoma.

B- Cocaine:

□ **Surface anesthesia** → loss of sensory reflex (corneal & conjunctival reflex)

□ **Indirect sympathomimetic:** ↓ neuronal uptake (1) + MAO inhibitor → ↑

endogenous NA → stimulates α_1 receptors → **active mydriasis & decongestion.**
No cycloplegia.

Block Na⁺ channel
└─ adrenergic preceptors.

cocaine overdose lead to death; arrhythmia

C) Parasympatholytics:

Mechanism: Block M3 receptors in:

- 1) **CPM** → passive mydriasis → lost light reflex & narrow angle of filtration.
- 2) **Ciliary muscle** → cycloplegia (loss of accommodation) + closing canal of Schlemm.

○ Result is **lost light reflex + cycloplegia + ↑↑ IOP.** **Examples:**

- 1) **Natural belladonna alkaloids: atropine & hyoscine.**
- 2) **Synthetic:** homatropine, cyclopentolate, tropicamide & eucatropine.

○ **Therapeutic uses:**

- 1) **Atropine** is used in iritis and corneal ulcer (to prevent adhesions), and measurement of refraction in children.
↳ examine without the effect of ciliary muscle.
- 2) **Synthetic substitutes:** in fundus examination.

All are contraindicated in glaucoma

Treatment of Glaucoma

Normal Intra-Ocular Pressure (IOP) = 15-25 mmHg.

Glaucoma کئی طرح سے ہوتا ہے

Glaucoma may be :

α_2
 β_2


1) Closed angle (narrow-angle) glaucoma

crowded iris, closed canal by root

(emergency) surgical intervention is essential

2) Open-angle glaucoma

(Chronic glaucoma)



→ Acute congestive glaucoma

I- Closed angle (narrow-angle) glaucoma:

- ❑ Needs **surgical intervention** (iridectomy).
- ❑ Due to **occlusion of angle of filtration** by iris root coming in contact with periphery of the cornea (Acute congestive glaucoma).

Drugs used to decrease I.O.P before surgery are:

- 1) **Miotic eye drops:** a) **Pilocarpine** (of choice) with low concentration.
b) **Physostigmine** (not preferred due to congestion & extreme miosis).
- 2) **Carbonic anhydrase inhibitors:** acetazolamide (↓↓ aqueous secretion)
↪ systemic
- 3) 3- Osmotic agents (dehydrating agent): mannitol (20%) IV, MgSO₄ rectally & Glycerine (50%) orally: they produce rapid reduction of IOP.
- 4) 4- Brimonidine & apraclonidine (α₂ agonists). *↳ Reduce Aqueous Humor.*
- 5) 5- Recently β-Blockers can be used with pilocarpine
↳ Reduce Aqueous Humor

II- Open-angle glaucoma (Chronic glaucoma):

← مسئلہ کلیتاً حسن بال drainage
← مسئلہ کلیتاً بال Production ← overproduction

• Drugs used are:

1) Miotic eye drops

(Pilocarpine & Physostigmine).


1) Carbonic anhydrase inhibitors:

(inhibit aqueous formation):

a- Acetazolamide orally

b- Dorzolamide & Brinzolamide

(locally)



3- Sympathomimetic eye drops: (Adrenaline & Dipivefrin) → VC → decrease synthesis of aqueous humor.

selective to B.V of ciliary Body "Locally"
→ prodn of adrenaline

4- B-blockers: decrease cAMP → decrease aqueous humor e.g. timolol & betaxolol. Side effects: tolerance, systemic absorption.

SUDDEN DECREASE of DRUG effectiveness.

5) α_2 agonists: a- Apraclonidine: used only for short time due to tachyphylaxis. b- Brimonidine: decrease aqueous secretion & \uparrow uveoscleral outflow.

Side effects: allergic conjunctivitis, dry mouth & fatigue.

6) PGF 2α analogues e.g. Latanoprost, travaprost & bimatoprost:

They decrease IOP by $\uparrow\uparrow$ uveoscleral outflow.

The most potent ocular hypotensives.

بطلوا ال
eyebrow & eyelashes

Side effects: conjunctival hyperpigmentation & hyperemia, and headache.

7) Guanethidine

↓
melanin

causes VD

Drugs that ↑↑ IOP:

1. Parasympatholytics (atropine). *mydriatic drug*

2. Drugs with atropine-like effect:

a) Some H1 blockers (Diphenhydramine).

b) Some antiarrhythmics

(Disopyramide). *substitute of quinidine with more Atropine like reaction.*

1. Ganglion blockers *→ corticosteroids ↑ IOP & ↓ IOP
↑ IOP, edema, ↓ matrix of meshwork ↓* *mostly with topical use.*

2. Corticosteroids. *Steroid induced glaucoma due to ↑ resistance of aqueous humor secretion.*

3. Nitrates. *vasodilation of blood vessels → ↑ ciliary secretion.*

Thank you