


CARDIOVASCULAR SYSTEM

HEART CHAMBERS & CARDIAC VALVES

Dr. Aiman Qais Afar
Surgical Anatomist

College of Medicine / University of Mutah

2022-2023





Chambers of the Heart

The heart is divided by vertical septa into four chambers:

❖ The **right and left atria** and the **right and left ventricles**.

❖ The **right atrium** lies **anterior to the left atrium**

❖ The **right ventricle** lies **anterior to the left ventricle**.




Chambers of the Heart

The walls of the heart are composed of cardiac muscle The myocardium;

covered externally with serous pericardium The epicardium

And lined internally with a layer of endothelium called The endocardium




Chambers of the Heart

Right Atrium

The right atrium forms the right border of the heart and receives venous blood from the **SVC, IVC**, and **coronary sinus**


The ear-like right auricle is a **conical muscular pouch** that projects from this chamber like **an add-on room**, increasing the capacity of the atrium as it overlaps **the ascending aorta**.



Right Atrium


The interior of the right atrium has a:

□ Smooth, thin-posterior wall, on which the venae cavae (SVC and IVC) and coronary sinus open, bringing poorly oxygenated blood into the heart.



Right Atrium


- ❑ Rough, muscular anterior wall composed of pectinate muscles
- ❑ Right AV orifice through which the right atrium discharges the poorly oxygenated blood, it has received into the right ventricle.



Right Atrium

- ❖ The **smooth and rough parts of the atrial wall** are separated **externally by** a shallow vertical groove, **the sulcus terminalis** or (terminal groove)
- ❖ and **internally by** a vertical ridge, **the crista terminalis** or (terminal crest).

sulcus terminalis



Two ways into heart:
superior and inferior
vena cava

Superior
Vena Cava

Interatrial
septum


divides right and left atrium

fossa ovalis

Auricle

Crista
Terminalis


Pectinate Muscle



Right Atrium

✓ The **SVC** opens into the superior part of the right atrium at the level of **the right 3rd costal cartilage**.


✓ The **IVC** opens into the inferior part of the right atrium almost in line with the **SVC** at approximately the level of **the 5th costal cartilage**



Right Atrium

✓ The opening of the coronary sinus, receiving most of the cardiac veins, is between the **right AV orifice** and the **IVC orifice**

✓ The interatrial septum separating the atria has an oval, thumbprint-size depression, the **oval fossa** (L. fossa ovalis), which is a remnant of the oval foramen (L. foramen ovale) and its valve in the fetus.




Chambers of the Heart

Right Ventricle

- ✓ Forms the largest part of the anterior surface of the heart
- ✓ a small part of the diaphragmatic surface
- ✓ and almost the entire inferior border of the heart

✓ Superiorly it tapers into an arterial cone, **the conus arteriosus (infundibulum)** which leads into the **pulmonary trunk**.


✓ The interior of the **right ventricle** has irregular muscular elevations (**trabeculae carneae**).



Right Ventricle

✓ A thick muscular ridge, **the supraventricular crest**, separates the **ridged muscular wall** of the **inflow part** of the chamber from the **smooth wall** of the **conus arteriosus**, or **outflow part**.

✓ The inflow part of the ventricle receives blood from the right atrium through **the right AV (tricuspid) orifice** located posterior to the body of the sternum at the level of **the 4th and 5th intercostal spaces**.




Right Ventricle

The **tricuspid valve** guards the **right AV orifice**.

The bases of the valve cusps are attached to **the fibrous ring around the orifice**.

- ❖ Because the **fibrous ring maintains the caliber of the orifice**, the attached valve cusps contact each other in the same way with each heartbeat.




Right Ventricle

13

Dr. Aiman Qais Afar
8 November 2022

Tendinous cords (L. **chordae tendineae**) attach to the free edges and ventricular surfaces of the **anterior, posterior,** and **septal cusps**, much like the cords attaching to a parachute.

The **tendinous cords** arise from the apices of **papillary muscles**, which are **conical muscular projections** with bases attached to the ventricular wall.




Right Ventricle

Dr. Aiman Qais Afar
8 November 2022

Three papillary muscles in the right ventricle correspond to the cusps of the tricuspid valve:


❖ The anterior papillary muscle, the largest one, arises from the anterior wall of the right ventricle



❖ The posterior papillary muscle, smaller than the anterior muscle, may consist of several parts; it arises from the inferior wall of the right ventricle

Right Ventricle


❖ The **septal papillary muscle** arises from the **interventricular septum**, and its tendinous cords attach to the anterior and septal cusps of the tricuspid valve.



Right Ventricle

Dr. Aiman Qais Afar
8 November 2022


The interventricular septum (IVS), composed of muscular and membranous parts, is a strong, obliquely placed partition between the right and left ventricles.



Superiorly and posteriorly, a thin membrane, part of the fibrous skeleton of the heart, forms the much smaller membranous part of the IVS

Right Ventricle


Because of the much higher blood pressure in **the left ventricle**, **the muscular part of the IVS**, which forms the majority of **the septum**, has the thickness of the remainder of the wall of **the right ventricle** (two to three times as thick as the wall of the right ventricle)



Right Ventricle

Dr. Aiman Qais Afar
8 November 2022

The **septomarginal trabecula (moderator band)** is a curved muscular bundle that traverses the right ventricular chamber from the inferior part of the IVS to the base of the anterior papillary muscle.




This trabecula is important because it carries part of **the right branch of the AV bundle**, a part of the **conducting system** of the heart to the anterior papillary muscle

LEFT ATRIUM

Similar to the right atrium, the left atrium consists of a **main cavity** and a **left auricle**. **The left atrium** is situated behind the **right atrium** and forms the **greater part of the base** or the **posterior surface of the heart**

Behind it lies **the oblique sinus of the serous pericardium**, and the fibrous pericardium separates it from **the esophagus**




LEFT ATRIUM

The interior of the **left atrium** is smooth, but the **left auricle** possesses muscular ridges as in the right auricle.

Openings into the Left Atrium

The **four pulmonary veins**, two from each lung, open through the posterior wall and have no valves. The **left atrioventricular orifice** is guarded by the **mitral valve**




LEFT VENTRICLE

- ❖ Forms the **apex of the heart**, and **most of the diaphragmatic surface**.

The interior of the left ventricle has:

- ❖ Walls that are **two to three times as thick** as those of the right ventricle.

- ❖ Walls that are mostly covered with a mesh of **trabeculae carneae** that are **finer and more numerous** than those of the right ventricle



- ❖ A **conical cavity** that is longer than that of the right ventricle.


❖ **Anterior and posterior papillary muscles** that are **larger than** those in the right ventricle.

❖ A smooth-walled, non-muscular, superoanterior outflow part, **the aortic vestibule**, leading to the **aortic orifice and aortic valve**.

❖ A **double-leaflet mitral valve** that guards the left AV orifice

❖ An aortic orifice that lies in its right posterosuperior part .
The ascending aorta begins at the aortic orifice.


B. Section through the heart showing the posteroinferior portion



The mitral valve

guards the atrioventricular orifice. It consists of **two cusps**, one **anterior** and one **posterior**.



✓ The **anterior cusp is the larger** and intervenes between the atrioventricular and aortic orifices.



✓ The **mitral valve** is located posterior to the sternum at the level of the **4th costal cartilage**.


The aortic valve

- ✓ guards the aortic orifice
- ✓ One cusp is situated on the **anterior wall (right cusp)** and two are located on the **posterior wall (left and posterior cusps)**.
- ✓ Behind each cusp, the aortic wall bulges to form **an aortic sinus**.



The aortic valve

✓ The anterior aortic sinus gives origin to **the right coronary artery**, and the left posterior sinus gives origin to the **left coronary artery**




✓ It is located posterior to the **left side of the sternum** at the level of the **3rd intercostal space**.

ATRIAL SEPTAL DEFECTS

A congenital anomaly of the interatrial septum, usually incomplete closure of the oval foramen, is an atrial septal defect (ASD).


The small openings, by themselves, cause no hemodynamic abnormalities and are, therefore, of no clinical significance and should not be considered forms of ASDs.



VENTRICULAR SEPTAL DEFECTS

Membranous part is the common site of ventricular septal defects (VSDs), although defects also occur in the muscular part

- ❖ VSDs rank first on all lists of cardiac defects. Isolated VSDs account for approximately **25%** of all forms of congenital heart disease.
- ❖ A VSD causes a left to right shunt of blood through the defect. A large shunt **increases pulmonary blood flow**, which causes severe pulmonary disease (hypertension, or increased blood pressure) and may cause **cardiac failure**.



Dr. Aiman Qais Afar



8 November 2022