






Schistosoma and hydatid cyst

Presented by

Associate Professor Dina Abou Rayia



Schistosoma species







Classification of *Schistosoma species*



Geographical distribution and habitat



S. Japonicum

Superior mesenteric veins mainly that supply the small intestine but can invade inferior mesenteric veins that supply the large intestine too

S. mansoni

Inferior mesenteric veins that supply the large intestine



**Why does Jordan lack a high
number of cases of schistosomiasis
???????**


General characters



THE ADULTS

The female

The male






Schistosoma mansoni




Intestinal caeca reunite at the anterior 1/3 of the body

- ❖ **Size:** 140x60 μ
- ❖ **Shape:** Oval with lateral spine
- ❖ **Color:** Translucent
- ❖ **Content:** Mature miracidium



Male and female in copula






Miracidium



Mother sporocyst



Daughter sporocyst



Furcocercus cercaria

Miracidium, Sporocyst, Daughter sporocyst, Cercaria

Larval stages




S. mansoni LIFE CYCLE

systemic circulation

lungs

venous circulation

Maturation in the portal vein of the liver




Adults inhabit inferior mesenteric veins



Female lay eggs in mesenteric venules of the large intestine



SCHISTOSOMULUM



Cercariae penetrate human skin

D. H.

Fresh water



Eggs pass with stool




Miracidium hatches



Miracidium penetrates snail

Cercariae attach to human skin

Cercariae emerge from the snail host



Daughter sporocyst

Mother sporocyst

CERCARIA

I. H.

Habitat: Inferior mesenteric veins

Host




Intestinal schistosomiasis





Stages of disease



Intestinal Schistosomiasis (Bilharziasis)

Stages of disease

1- Stage of invasion

Manifestations

❖ Skin lesion due to cercarial penetration.

❖ Local dermatitis, irritation, itching and papular rash.



Intestinal Schistosomiasis (Bilharziasis)



Stages of disease

2- Stage of migration

- ❖ **Lung** : Irritation due to passage of schistosomulum causing minute haemorrhage, cough, sputum, dyspnea and eosinophilia, and pneumonitis.
- ❖ **Liver** : Enlarged tender liver and spleen.
- ❖ **Toxic symptoms**: Due to metabolic products of maturing parasites causing fever, anorexia, headache, malaise and muscle pain.





Intestinal Schistosomiasis (Bilharziasis)

Stages of disease

3- Stage of maturation (acute schistosomiasis-Katayama syndrome)

- The development of schistosomes into sexually mature, egg-producing adults with the beginning of egg-laying produces a form of acute schistosomiasis which is a systemic hypersensitivity reaction like serum sickness.
- It is manifested by fever, vomiting, diarrhea, enlarged lymph nodes and hepatosplenomegaly with marked eosinophilia.

4- Stage of egg deposition and tissue reaction



Eggs can reach the intestinal wall by:

- 1- Egg spine.
- 2- Proteolytic enzymes.
- 3- Elastic recoil of blood vessels.

The eggs have three destinations:

- 1- Pass through the wall to the lumen, or
- 2- Trapped in the wall which leads to granuloma, fibrosis, and strictures, or
- 3- Eggs moved with the venous circulation forming embolism. (Liver, lung, CNS, skin,)

Intestinal Schistosomiasis (Bilharziasis)



Stages of disease


4- Stage of egg deposition and tissue reaction

- ❖ Trapped eggs in the intestinal wall → formation of **polyps, ulcers and granuloma causing** abdominal pain, diarrhea and dysentery.
- ❖ Later on, the intestinal wall becomes **fibrosed** and thickened → **stricture of the wall.**
- ❖ Sinuses or fistula can occur.
- ❖ Rectal prolapse.
- ❖ The eggs secrete **proteolytic** enzymes that provoke typical eosinophilic inflammatory and granulomatous reactions (**bilharzial granuloma**), which are progressively replaced by fibrotic tissue which is the main cause of pathology and complications.



Embolitic lesions

Some eggs are swept back into the blood stream to different organs:-



Clinical picture summary




Stages	Clinical aspect	Manifestations
Early	1. Cercarial dermatitis	At the penetration sites of cercariae → itching & papular eruption
	2. Schistosomular migration	Migration of schistosomula → lungs : pneumonitis (fever, cough and haemoptysis) and → liver (tender hepatomegaly)
	3. Acute schistosomiasis (Katayama syndrome)	It occurs when worms mature in the liver, migrate to the small venules and begin to lay eggs. There is fever, abdominal pain, diarrhoea, wheezing, urticaria, marked eosinophilia, sometimes lymph node enlargement and hepatosplenomegaly.

Clinical picture



Stages	Clinical aspect	Manifestations
Late manifestations	1. Chronic Intestinal schistosomiasis	Oviposition in the mesenteric plexus → diarrhoea with blood and mucus (schistosomal dysentery)
	2. Chronic hepatosplenic schistosomiasis	Granuloma Formation in the liver → periportal fibrosis → Obstruction of the portal venous branches → portal hypertension → hepatomegaly & splenomegaly
	3. Advanced complications	Hypersplenism → Anaemia + thrombocytopenia Extensive periportal fibrosis → Hepatic failure Portal hypertension → Opening of porto-systemic collateral → oesophageal varices → fatal haematemesis Egg embolism → Lung & CNS Ascites due to hypoproteinaemia + portal hypertension




(A)



(B)



(C)





Laboratory diagnosis

Direct

- 1) Detection of **eggs in the stool** by direct smear or concentration .
- 2) Thick faecal smear .
- 3) **Rectal swab.**
- 4) **Rectal biopsy or liver biopsy** in chronic stage

Indirect

- 1) Intradermal test.
- 2) **Serological tests :**
IHAT, CFT, and ELISA.
- 3) **Recently:** Detection of circulating *Schistosoma* antigens by using of monoclonal antibodies
- 4) **Anaemia:-**
 - Iron deficiency anaemia due to blood loss.
 - Haemolytic anaemia due to hypersplenism.
- 6) **Eosinophilia**



Treatment


Medical

Praziquantel effective against adult worms

Artemisinin effective against schistosomulum




Surgical

For complications




Hydatid cyst disease

Life Cycle of *Echinococcus granulosus*





Unilocular hydatid cyst



Hydatid cyst





Hydatid cyst disease (Cystic Echinococcosis or Hydatidosis)

- ❖ It is a parasitic infection of both humans and other mammals such as sheep, and cattle with hydatid cysts, the larval stage of *Echinococcus granulosus*.
- ❖ Man is an intermediate and blind host for *Echinococcus granulosus*

Pathogenesis & Symptomatology



Local inflammatory reaction around the hydatid cyst, ending in formation of a fibrous capsule which may become calcified or even ossified.

The symptoms depend on the size & site of the cyst.

Large sized cysts \Rightarrow pressure atrophy of affected organs:-
Liver (70%) \Rightarrow enlargement and dysfunction (fever, pain and jaundice).
Lung (20%) \Rightarrow pain, cough and dyspnea.
Brain \Rightarrow epilepsy.
Eye \Rightarrow protrusion of the eye ball.
Bones \Rightarrow Pain & spontaneous fracture.
Kidney \Rightarrow membranous nephropathy.

Spontaneous rupture of cyst into peritoneal cavity or pleura may lead to severe allergic reaction (**anaphylactic shock**) or secondary cysts.



Diagnosis

Clinical

- History of contact with dogs.
- Slowly growing cystic tumour.
- Hydatid thrill.

Laboratory

Direct

- X-ray for calcified cyst.
- Ultrasonography, CT scan and MRI.
- Scolices in sputum or urine due to rupture of the cyst in bronchus or urinary tract.
- Puncture or aspiration of hydatid fluid
 - ⊖ may lead to anaphylactic shock due to leakage of the fluid.

Indirect

- Eosinophilia.
- Intradermal test (**Casoni test**).
- Serological tests.
- PCR



Treatment

1) **Surgical removal of the cyst:** The most efficient treatment but it may cause mortality (2%) and recurrence of the disease (2 - 25%).

2) **Percutaneous treatment (PAIR):** In three steps:

Puncture (P) and needle aspiration (A) of the cyst.

Injection (I) of a scolicidal solution usually hypertonic sodium chloride solution or ethanol and left for 5 - 30 minutes.

Cyst-re-aspiration (R) and final washing.

- ✓ This procedure is indicated in **inoperable cases** and who have **drug resistance** (no response to medical treatment).





3) **Medical treatment:**

Indications: In inoperable cases and before and after surgery.

- **Albendazole (Drug of choice).**
- **Mebendazole.**
- **The combination of ABZ and Praziquantel (PZQ) may provide synergistic effect and better efficacy.**





Identify ??????





Case study

- A 24-year-old man presented to the hospital complaining of a swelling in the right upper quadrant of his abdomen. Clinical examination revealed the presence of a mass on the right side of the abdomen that elicited a thrill on palpation. Blood examination revealed eosinophilia. Abdominal ultrasound showed a medium-sized cyst with heterogenous contents occupying the right liver lobe.