

1\_One of the following is not a complication of celiac disease

Select one:

- a, T-cell lymphoma
- b. Osteoporosis
- c. Aplastic anemia
- d. Ulcerative jejunitis
- e Increased risk of esophageal carcinoma

تعليق 

أعجبني 

### ▼ All comments

وليد عزايزة  
اعرفوا شغلة بس ناخذ biopsy ونشوفها بالهستو  
راح نلاقي  
villus atrophy (1  
crypt hyperplasia (2  
lymphocytosis (3



١١ دقيقة أعجبني رد المزيد

وليد عزايزة  
واكثر complication هو  
T cell lymphoma



NOT B cell

١٠ دقائق أعجبني رد المزيد

77 Regarding the clinical features of celiac disease. One is false

Select one:

- a. Can be diagnosed after the age of 60
- b. Can appear in infancy upon weaning from milk to solid foods
- c. Has a peak of incidence in the fifth decade
- d. Patients can be asymptomatic and present only with laboratory abnormalities
- e. Mouth ulcers and angular stomatitis are indicators of very severe disease

٢ تعليق

تعليق

أعجبني

وليد عزايذة

اهم اشى تعرفوا

Celiac disease can occur at ANY AGE

يعني من ساعة ما بنولد لغاية ما يموت يمكن يصير معاه celiac disease

١٢ دقيقة أعجبني رد المزيد

وليد عزايذة

وهو بصير معاه

Mouth ulcer & angular stomatitis

بس لا يعني انه severe disease

١٢ دقيقة أعجبني رد المزيد

30) A 27-year-old woman presents for review. She describes herself as having 'IBS' and for the past two years has suffered intermittent bouts of abdominal pain, bloating and loose stools. For the past two weeks however her symptoms have been much worse. She is now passing around 3-4 watery, grey, 'frothy' stools per day. Her abdominal bloating and cramps have also worsened and she is suffering from excessive flatulence. Judging by the fitting of her clothes she also feels that she has lost weight. Some blood tests are ordered Hb 10.9 g/dl Platelets 199 \* 10<sup>9</sup>/l WBC 7.2 \* 10<sup>9</sup>/l Ferritin 15 ng/ml Vitamin B12 225 ng/l Folate 2.1 nmol/l! What is the most likely diagnosis?

Select one:

- a. Crohn's disease
- b. Celiac disease
- c. Infective exacerbation of irritable bowel syndrome
- d. Ulcerative colitis
- e. Bacterial overgrowth syndrome

تعليق 

أعجبني 

## ▼ All comments

وليد عزايزة  
اعرفوا شغلة انه  
Patient with celiac disease  
بصير معاهم steatorrhea



٨ دقائق أعجبني رد المزيد

60\_Celiac patients are instructed to report to clinical follow up every 3-6 months. What is the best way to ensure strict diet if you suspect that your patient is not compliant on the requested dietary program?

Select one:

- a. Anti Endomysial antibodies
- b. Anti Tissue Transglutaminase antibodies**

c. Endoscopy with duodenal biopsy

d. Detailed history of what he eats

١ تعليق

تعليق

أعجبني

وليد عزايذة



Most specific antibody is

Anti -TTG

اختصارها هو

Anti- tissue transglutaminase antibody

١٠ دقائق أعجبني رد المزيد

اكتب تعليقًا...



Deficiency of which immunoglobulin is frequently encountered in celiac disease?

Select one:

- a. IgA**
- b. IgE
- c. IgM
- d. IgG
- e. Complement system

تعليق

أعجبني

اكتب تعليقًا...



60\_Celiac patients are instructed to report to clinical follow up every 3-6 months. What is the best way to ensure strict diet if you suspect that your patient is not compliant on the requested dietary program?

Select one:

- a. Anti Endomysial antibodies
- b. Anti Tissue Transglutaminase antibodies**

c. Endoscopy with duodenal biopsy

d. Detailed history of what he eats

تعليق 

أعجبني 

### ▼ All comments

وليد عزايزة



Most specific antibody is

Anti -TTG

اختصارها هو

Anti- tissue transglutaminase antibody

١٣ دقيقة أعجبني رد المزيد



100- most common HLA subtype seen in celiac disease is:

Select one:

- a. HLA DR3
- b. HLA DR4
- c. HLA DQ2
- d. HLA DQ8
- e. HLA B27

١ تعليق

تعليق 

أعجبني 

وليد عزايذة

Celiac disease is associated with both HLA-DQ2 & HLA-DQ8

بس اكثررر اشى يعنى 90% بكون HLA-DQ2

١٣ دقيقة أعجبني رد المزيد



Deficiency of which immunoglobulin is frequently encountered in celiac disease?

Select one:

- a. IgA
- b. IgE
- c. IgM
- d. IgG
- e. Complement system