




# Hydrogenation of Alkenes and Alkynes


Hydrocarbons that have carbon-carbon double bond are called alkenes; those with a carbon-carbon triple bond are alkynes

Their general formulas are  $C_nH_{2n}$  alkenes and  $C_nH_{2n-2}$  alkynes




Both alkenes and alkynes are unsaturated hydrocarbons







Compounds with more than one double or triple bonds exist. Multiple double bonds may lead to dienes, trienes, tetraenes and polyenes.  $\beta$ -carotene and lycopene are examples of polyenes



When two or more multiple bonds are present in a molecule, they can be classified depending on the relative positions of the bonds




Which of the following compounds have conjugated multiple bonds?




# Nomenclature


1. The ending -ene is used to designate carbon-carbon double bond. When more than one double bond is present, the ending is *-diene*, *triene*, *tetraene* and so on. The ending -yne is used for triple carbon-carbon bond.
2. Select the longest chain that includes both carbons of the double bond or triple bond.




Number the chain from the end nearest the multiple bond so that the carbon atoms in that bond have the lowest possible numbers.




*not*



If the multiple bond is equidistant from both ends of the chain, number the chain from the end nearest the first branch point.



*not*




2-methyl-2-butene

Indicate the position of the multiple bond using the lower numbered carbon atom of that bond.




1-butene, *not* 2-butene

If more than one multiple bond is present, number the chain from the end nearest the first multiple bond.



*not*




= END =





If the double bond and the triple bond are equidistant from the end of the chain, the **double** bond receives the lowest number.

*alkene takes priority over alkyne*











The root name is from the longest carbon chain containing the multiple bond




*So are  
1-butene & 2  
2-butene*






## The numbering rules applied




4-methyl-2-pentene ✓

(Not 2-methyl-3-pentene; the chain is numbered so that the double bond gets the lower number.)



2-ethyl-1-butene ↗

(Named this way, even though there is a five-carbon chain present, because that chain does not include both carbons of the double bond.)



1,3-butadiene


(Note the *a* inserted in the name, to help in pronunciation.)

With the cyclic hydrocarbons, we start numbering the ring with the carbons of the multiple bond.




cyclopentene

(No number is necessary, because there is only one possible structure.)




3-methylcyclopentene

(Start numbering at, and number through the double bond; 5-methylcyclopentene and 1-methyl-2-cyclopentene are incorrect names.)



1,3-cyclohexadiene




1,4-cyclohexadiene

Write the structural formula for



3-methyl-2-pentene




1,4-dichloro-2-pentene




3-hexyne




1,2-dimethylcyclobutene



2-bromo-1,3-pentadiene



## Some Common names




ethylene

(ethene) ✓











acetylene

(ethyne) ✓




propylene

(propene) ✓






# Some Facts about Double Bonds



$sp^2$   $sp^2$

trigonal planar

$120^\circ$




= END =



**TABLE 3.1** Comparison of C—C and C=C Bonds


Property	C—C	C=C
1. Number of atoms attached to a carbon	<u>4</u> (tetrahedral)	<u>3</u> (trigonal)
2. Rotation	relatively <u>free</u>	<u>restricted</u>
3. Geometry	<u>many</u> conformations are possible; staggered is preferred	<u>planar</u>
4. Bond angle	109.5°	120°
5. Bond length	<u>1.54 Å</u>	<u>1.34 Å</u>


# The Orbital Model of a Double Bond; the pi Bond




Atomic orbitals  
of carbon

The 2s and two 2p orbitals are combined to form three hybrid  $sp^2$  orbitals, leaving one electron still in a  $p$  orbital.









side view



side view with  
the  $sp^2$  orbitals  
represented by lines




top view




$p-p$   $\pi$  bond

$sp^2-s$   $\sigma$  bond



$sp^2-sp^2$   $\sigma$  bond






restriction since  
rotation will distort  
and break the p-p  $\pi$   
bond

# Cis-Trans Isomerism in Alkenes






# Are cis-trans isomers possible for 1-butene and 2-butene?





*cis-2-butene*

bp 3.7°C, mp -139°C




*trans-2-butene*

bp 0.3°C, mp -106°C



1-butene


is identical to



1-butene

Whenever you have identical groups on one of the C=C bonds  $\Rightarrow$  NO <sup>of cis/trans</sup> possible


Geometric isomers of alkenes can be interconverted if sufficient energy is supplied to break the pi bond and allow rotation about the remaining sigma bond.



## Reactions of Alkenes

- The chemistry of alkenes can be divided into two general types of reactions:

### (1) Electrophilic Addition Reactions



*Addition of Symmetric and Unsymmetric Reagents to symmetric Alkenes.*

1. Addition of Hydrogen: Catalytic Hydrogenation
2. Addition of Halogens: Halogenation

*Addition of Unsymmetric Reagents to Unsymmetric Alkenes; Markovnikov's Rule.*

1. Addition of Hydrogen Halides
2. Addition of Sulfuric Acid
3. Addition of Water: Hydration
4. Addition of HOX: Halohydrin Formation

# Reactions of Alkenes



## (2) Oxidation Reactions

1. Ozonolysis
2. Oxidation Using  $\text{KMnO}_4$

Electrophilic Addition Reactions

1. Addition of Hydrogen: Hydrogenation

Addition of a mole of hydrogen to carbon-carbon double bond of Alkenes in the presence of suitable catalysts to give an *Alkane*.





## Electrophilic Addition Reactions

56


### 2. Addition of Halogen: Halogenation

When an **alkene** is treated at room temperature with a solution of **bromine** or **chlorine** in carbon tetrachloride to give the corresponding **vicinal dihalide** (two halogens attached to adjacent carbons)




- **Iodine** is **too unreactive** and will not add to the double bond.
- **Fluorine** is **too reactive** and reacts explosively with an alkene.

## Addition and Substitution Reactions Compared



Addition of halogens  $X_2$  ( $\text{Br}_2, \text{Cl}_2, \text{I}_2, \text{F}_2$ )





2-butene

bp  $1-4^\circ\text{C}$



2,3-dichlorobutane


bp  $117-119^\circ\text{C}$





### Addition of Water (Hydration)






Unsaturated Cycloalkene  
 non polar

alcohol  
 polar


$\Rightarrow$  hydrogen bonding



Bromine added  
to →  
saturated




← Br<sub>2</sub> to  
unsaturated Hydrocarbons  
alkene  
alkyne





- END -

## Addition of Acids



Acids that add this way are the hydrogen halides (H-F, H-Cl, H-Br, H-I) and sulfuric acid (H-OSO<sub>3</sub>H)





Write the equation for each of the following reactions

a) 2-butene + HCl

b) 3-Hexene + HI




a) 4-methylcyclopentene + HBr

## Table 3.2 ▀ Classification of Reagents and Alkenes by Symmetry with Regard to Addition Reactions




**Symmetric**

**Unsymmetric**

Reagents




Alkenes




mirror plane



not a mirror plane


# Addition of Unsymmetric Reagents to Unsymmetric Alkenes; **Markovnikov's Rule**













# Mechanism of Electrophilic Addition to Alkenes





tertiary (3°)  
most stable



secondary (2°)


primary (1°)

methyl (unique)  
least stable





## Hydroboration of Alkenes








transition state  
for hydroboration














What alkene is needed to obtain the alcohol below via hydroboration-oxidation sequence, what product would this alkene give with acid-catalyzed hydration.





## Addition of Hydrogen






## Oxidation with permanganate; a Chemical Test




## Ozonolysis of Alkenes

- The first product, a **molozone**, is formed by cycloaddition of the oxygen at each end of the ozone molecule to the carbon-carbon double bond.
- This product then rearranges rapidly to an **ozonide** (explosive if isolated).
- They are usually treated directly with a reducing agent, commonly **zinc and aqueous acid**, to give **carbonyl compounds** as the isolated products.









# Reactions of Alkenes

## Oxidation Reactions

67

## 2. Oxidation Using $\text{KMnO}_4$

Alkenes react with alkaline potassium permanganate to form glycols (compounds with two adjacent hydroxyl groups).





Hexane does not react with purple  $\text{KMnO}_4$  (left); cyclohexene (right) reacts, producing a brown-black precipitate of  $\text{MnO}_2$ .

# The Structure of Alkynes

69


- **Alkynes** are hydrocarbons that contain a **carbon–carbon triple bond**.
- **Alkynes** are also known as **Acetylenes**.
- General formula is  **$C_nH_{2n-2}$**
- Hybridization;  **$sp$ -hybridized orbitals**
- The angle between them is  **$180^\circ$**  and the bond length  **$1.20 \text{ \AA}$**
- The geometry is **Linear**.





Atomic orbitals  
of carbon

The 2s and one 2p orbital are combined to form two hybrid *sp* orbitals, leaving one electron in each of two *p* orbitals.




The resulting carbon–carbon triple bond, with a hydrogen atom attached to each remaining  $sp$  bond. (The orbitals involved in the C–H bonds are omitted for clarity.)

# Reaction of Alkynes

## Electrophilic Addition Reactions

### 1. Addition of Hydrogen: Hydrogenation

- With an ordinary nickel or platinum catalyst, alkynes are hydrogenated all the way to alkanes.
- However, a special palladium catalyst (called Lindlar's catalyst) can control hydrogen addition so that only one mole of hydrogen adds. In this case, the product is a **cis alkene**.
- On the other hand, reduction using metals such as sodium or lithium in liquid ammonia results in formation of **trans alkenes**.




# Reaction of Alkynes

## Electrophilic Addition Reactions

75


### 2. Addition of Halogen: Halogenation

Bromine adds as follows; In the first step, the addition occurs mainly *trans*.



### 3. Addition of Hydrogen Halide


With unsymmetrical triple bonds and unsymmetrical reagents, Markovnikov's Rule is followed in each step, as shown in the following example:




# Acidity of Alkynes

71

- A hydrogen atom on a triply bonded carbon (**Terminal Alkyne**) is weakly acidic and can be removed by a very strong base ( as Sodium amide).



- Internal alkynes (**Non-Terminal Alkyne**) have no exceptionally acidic hydrogens.
  - Relative Acidity of the Hydrocarbon.  
Terminal alkynes, are more acidic than other hydrocarbons








# Preparation of Alkynes


73


## 2) Reaction of Sodium Acetylide with Primary Alkyl Halides

### ○ Acetylene



### ○ Monosubstituted Acetylenes






$sp^3$

25%  $s$ ,

75%  $p$



$sp^2$

$33\frac{1}{3}\%$   $s$ ,

$66\frac{2}{3}\%$   $p$



$sp$

50%  $s$ ,

50%  $p$

increasing acidity



## Electrophilic Addition Reactions

76

### 4. Addition of Water: Hydration

- **Addition of water to alkynes** requires not only an acid catalyst but mercuric ion as well.
- Although the reaction is similar to that of alkenes, the initial product - a vinyl alcohol or enol - rearranges to a carbonyl compound (keto form).
- The keto form of aldehydes and ketones are in equilibrium with the enol form.
- The keto form predominates at equilibrium for most simple aldehydes and ketones.
- The inter conversion is called **keto-enol tautomerization**.

