


Cholinergic Antagonist Drugs



Dr. Saed M. Aldalaen

Mutah Univ. 2017

Cholinergic Antagonist Drugs



- **Anti-muscarinic drug:** Atropine-like drugs, Hyoscine (Scopolamine)
- **Anti-nicotinic drugs**
- **Ganglion blockers:** Used in experimental pharmacology. E.g. Nicotine, Trimethapan.
- **Neuro-muscular blockers:** Used in surgery to produce complete muscle relaxation.

Anti-muscarinic anti-cholinergic drugs



- Natural agents:
 - **Atropine, Hyoscine**
 - Semi-synthetic
 - **Homatropine**
 - Synthetic
 - **Ipratropium, Pirenzepine, Propantheline**

Anti-muscarinic

- Atropine (Hyoscyamine)
- Alkaloids obtained from *Atropa Belladonna*,
- Considered as prototype for parasympatolytics



- Hyoscine (Scopolamine)
- Obtained from *Hyocyamus niger* plant (*Datura Stramonium*)



Note: Antihistamines, phenothiazides and some antidepressants have anti-muscarinic effects

Clinical pharmacology of anti-muscarinic drugs

Mechanism of action:

- **Reversible blockade** of **M** receptors
- **Exocrine glands** are most **sensitive**
- **Gastric secretion** is the **least** affected
- **Heart** is **intermediate**

Note: Atropine blocks all 3 subtypes receptors (M1,M2,M3)

Pharmacokinetics

- **Absorption:**
- **Natural** and most **tertiary amines: good**
- **Wide distribution** and **cross BBB**
- **Quaternary amines: poorly absorbed** and poor crossing **BBB** (Ipratropium)
- Atropine $t_{1/2}$: 2hrs
- **Partly metabolized** and partly **excreted unchanged**

Routes of administration



Oral



Parenteral



Topical (suppositories)



**Some by inhalation
(Ipratropium)**

Pharmacodynamics

- **Exocrine glands: at low doses reduced secretions**
- Salivary
- Bronchial
- Sweet glands

CNS



- Central **stimulant** effects (Atropine) *Cross 131313*
- Some may produce **sedation** (Hyoscine) *given with anaesthesia*
- Hyoscine blocks M receptors in vomiting centre and has **anti-emetic effect**
- **Toxic doses:** hallucination, convulsion, coma

Eye



- **Mydriasis** (dilatation of pupil)
- **Cycloplegia** (relaxation of the ciliary muscle) cause: blurred vision and impaired accommodation to near vision
- **Decreased lacrimation**
- **Increase IOP**

glaucoma is a contraindicator.

CVS



Depending in the doses

- **Central effect:**
- **Decrease heart rate**
- **Peripheral effect:**
- Blockade of vagus nerve and **increase heart rate**
- **ABP:**
- **No change**

Respiratory system



- **Bronchodilatation**
- **Reduced** bronchial **secretion**
- Ipratropium (quaternary amine derivate of Atropine) inhalation:
- Useful in asthma and chronic obstructive pulmonary disease (COPD), also in patient who are unable to take adrenergic agonists.

block M_1 / M_3

GIT



- **Decrease salivation**
- **Decrease acid secretion**
- **Decrease motility**
- **Delay gastric emptying**
- **Prolong intestinal transit time**
- **Anti-diarrhoeal and anti-spasmodic effects**

genitourinary tract
GUT M₃



- **Relaxation of bladder wall**
- Useful in **inflammatory spasm** and **pains** of the urinary tract
- **Risky** in patients with **BPH** (Benign Prostatic Hypertrophy)

Therapeutic uses

CNS disorders:

- **Parkinson's** disease → *"excessive stimulation of cholinergic receptors"*
- Drug-induced parkinsonism as Phenothiazine (induced acute dystonias)
- Benztropine, Benzhexol: useful
- **Motion sickness**: Hyoscine oral, injection, transdermal patches

Therapeutic uses

Ocular uses:

- In **eye examination** (Tropicamide) produce mydriasis and cycloplegia
- In **iritis** (Atropine eye drop) prevent synechia (adhesion of the iris to the lens)

Note:

- Atropine eye drops effects: 7 days
- Tropicamide eye drops effects: 4-12hrs

Therapeutic uses

- **Premedication:** Hyoscine and Atropine (use as adjunct in anaesthetic procedure)
- **Bronchial asthma:** Ipratropium inh. (produce bronchodilatation)

Cardiovascular:

- **Bradycardia and heart block following AMI:**
Atropine

Therapeutic uses

GI disorders:

- **Anti-diarrhoeal**

- Lomotil= atropine + diphenoxylate

- **Anti-spasmodics** (in intestinal colic, IBS)

- Atropine, hyoscine, clidinium, prifinium

"Buscopan" "Libraxam" "Kibal"

Urinary disorders:

- **Urinary urgency** with UTI
- **Renal colic**

Therapeutic uses

- **Cholinergic poisoning as:**
 - Irreversible CEI insecticide poisoning
 - Chemical warfare intoxication
- To **counteract muscarinic effects**
- (nicotinic effects can not be reversed)
- Atropine IV

Adverse effects of anti-muscarinic agents



- **Dry mouth**
- **Blurred vision**
- **Tachycardia**
- **Constipation**
- **Hot flushed dry skin & hyperthermia may occur with high doses**

Contraindications



- **Glaucoma**
 - Increase IOP
- **BPH**
 - Bladder wall relaxation & sphincter contraction

Atropine poisoning



- **Hot flushed dry skin & hyperthermia,**
- **Agitation, delirium, hallucination,**
- **Convulsions & coma**
- **Treatment is symptomatic**

Individual drugs



- **Atropine**
- **Hyoscine**
 - Buscopan
- **Clidinium**
 - Libraxam
- **Prifinium**
 - Riabal