



# 3- MUSCLE PHYSIOLOGY- I



**Prof. Sherif W. Mansour**  
Physiology dpt., Mutah School of medicine  
2024

# The Muscle

Muscles are divided into 3 types:

	<b>Skeletal</b>	<b>Cardiac</b>	<b>Smooth</b>
-Striations	Striated	Striated	Non-striated
-control	Voluntary	Involuntary	Involuntary
-innervation	Somatic	Autonomic N.S.	Autonomic N.S.
-Function	Movement	Pumping of blood	According to site
-site	Attached to bone	Heart	Gut, bl.vs, others

## Skeletal muscle

- Functions of the skeletal muscle:

1-It generates **force** which is helpful in daily life.The muscle fibres are arranged in variable axis, longitudinal, oblique or radial to exert force in different planes.

2-It is very important in **respiration**.

3-It is very important in **producing heat** and keeping body temperature constant.

4-It is very important in **venous return** of the blood to heart against gravity (muscle pump).

5- It is very important in keeping **equilibrium** of the body.

6- It has a **protective** function (withdrawal reflex).


7- It helps **posture (Equilibrium)**.

8- Contraction of anterior abdominal wall and pelvic diaphragm muscle keeps **intra-abdominal pressure** positive and keep organs in position and help respiration.


9-It is important in mechanism of **hearing**.

10-Movement of the **extra-ocular muscles** increase range of vision.


**(b) Diagram of part of a muscle fiber showing the myofibrils. One myofibril extends from the cut end of the fiber.**




**(c) Small part of one myofibril enlarged to show the myofilaments responsible for the banding pattern. Each sarcomere extends from one Z disc to the next.**



**(d) Enlargement of one sarcomere (sectioned lengthwise). Notice the myosin heads on the thick filaments.**



**(e) Cross-sectional view of a sarcomere cut through in different locations.**



-The arrangement of these contractile proteins in the filament shows cross striations with alternative dark (A) and light (I) bands.

**-The dark band:**

- .Called **A** band, 100 A° in diameter.
- .Contain myosin filaments and ends of actin filaments.
- .Has lighter central **H-zone** (has no actin filaments).
- .**M-line** in middle of H-zone due to central bulge on thick myosin.

**-The light band:**

- . Called **I** (Isotropic to light) band - 50 A° diameter.
  - . Contain thin actin filaments + troponin.
  - . Contain central **Z-line** (site of attachment of thin filaments).
- The area between two **Z-lines** is called sarcomere which is the contractile unit of the muscle.
- Longitudinal striations are due to longitudinal arrangement of the muscle fibres and transverse striations are due to alternating light and dark bands.

# Sarco-tubular system

-It is a system of tubules present in the sarcoplasm of the muscle fibre and made of:

## 1) T- tubules (transverse system):

-It is a deep infolding of the sarcolemma near endings of the sarcomere (z-line) as an extension of the extracellular space, this infolding is in perpendicular direction of the myofibril so it is called transverse tubule.

-T-tubule membrane contains a voltage sensitive **dihydropyridine (DHP)** receptor that open the ryanodine  $Ca^{++}$  release channel on the sarcoplasmic reticulum


- Through this tubule the action potential from the cell membrane is rapidly spread to fibrils and fluids with ions move into and out the cell.

## 2) The sarcoplasmic reticulum:

- It is formed of irregular longitudinal tubules that run parallel and surround the muscle fibre to be a source of  $Ca^{++}$  ions, via Ryanodine  $Ca^{+2}$  channels.

- At site of contact to T-tubules it dilated to form terminal cisterns.

-Each 2 terminal cisterns with adjacent T-tubules is called triad.



# Muscle proteins

## [A] Contractile proteins:

**1- Myosin:** -Myosin is complex protein with M.W. 480,000.


-Composed of **6** polypeptide chains (2 heavy chains and 4 light chains).

-The 2 heavy chains wrap spirally around each other as double helix forming long tail (light meromyosin) and arm (heavy meromyosin) while the terminal part combine with the 4 light chains forming 2 globular heads ,one head contains actin-binding sites and the other contain sites of ATP hydrolysis.

-Cross bridges arise from the head with arm of 2 flexible points called **hinges** (one between arm and tail and the other between the arm and heads) to bind to the actin.

**2- Actin:** - It is small **globular** protein with M.W. 42,000.

- The globules attached to each other to form filamentous structure arranged in two chains as long double helix.



## [B] Regulatory protein:

### 1- Tropomyosin:

-It is long filament of two polypeptide chains twisting on each other and located between the 2 chains of actin covering its active sites which combine to myosin and keeps the actin structure.

### 2- Troponin:


-Small globules located at intervals along tropomyosin.

-Of 3 subunits with MW 18,000-25,000.

1-Troponin T: binds troponin to tropomyosin.

2-Troponin I: inhibit binding of actin & myosin.


3-Troponin C: bind  $Ca^{+2}$  ions → contraction.



# Mechanism of muscle contraction (Excitation - contraction coupling)

It is the process by which depolarization of the muscle fiber initiate contraction.

- 1-When a nerve impulse reach the MEP, it leads to Ach release from the nerve terminals.
- 2-Ach combines with the **cholinergic receptors** on the muscle membrane  $\rightarrow \uparrow \text{Na}^+$  permeability  $\rightarrow$  depolarization of the membrane (End plate potential).
- 3-When the **EPP** reaches the firing level  $\rightarrow$  action potential that propagates along the muscle membrane and transmitted to all fibers via the T tubules.
- 4-This action potential triggers the release of  $\text{Ca}^{++}$  ions from the terminal cisternae of sarcoplasmic R.
- 5-The concentration of  $\text{Ca}^{+2}$  increases and initiates contraction by binding to troponin-c leading to:
  - a- Weakness of the binding of troponin I to actin.
  - b- Movement of tropomyosin laterally into the groove between the thin filaments  $\rightarrow$  uncovering the binding sites of actin for the myosin heads.
- 6-The interaction between actin & myosin heads leads to sliding of actin filaments between myosin filaments  $\rightarrow$  muscle contraction.
- 6-The energy required for this mechanism is provided by breakdown of ATP to ADP by ATPase activity of myosin heads in the presence of  $\text{Ca}^{++}$  ions.



## Results of contraction

- The **sarcomere** becomes **short**.
- The width of **I** band is **decreased**.
- The width of **A** band is remained **constant**.
- H-zone** becomes **narrow**.

## Mechanism of muscle relaxation

- The  $\text{Ca}^{+2}$  is actively reuptake back to the SR by active  $\text{Ca}^{+2}$  pump to be stored in the cisterns.
- Decrease the intracellular  $\text{Ca}^{+2}$  ions to  $10^{-7}$  mol/L, the troponin-tropomyosin complex return to its original position separating myosin head from actin that is covered and inhibited by tropomyosin resulting in muscle relaxation.
- Breakdown of ATP is necessary to  $\text{Ca}^{+2}$  pump.

So, ATP hydrolysis is required for contraction and relaxation and  $\downarrow$  ATP  $\rightarrow$  no relaxation.

**Electrical changes during M. contraction:** Similar to the nerve with the following differences

	<b>Muscle</b>	<b>Nerve</b>
- RMP	- 90 mV.	- 70 mV. to - 90
- Firing level	- 70 mV.	- 55 mV or - 65
- Overshoot	+ 40 mV	+ 35 mV
-Amplitude of A.P.	130 mV	105 mV or 125
- Duration of spike P.	2-4 msec	2 msec
- Rate of conduction	5 meter/sec	Variable
- Chronaxi	Long (less excitable)	Short (more excitable)

# Rigors : (contracture)

-It is a state of stiffness (sustained contraction) in the muscle. -Of four types:

## 1-Rigor mortis:

-It is a state of stiffness in the muscle **five hours** after death due to **depletion of ATP**, which is required for muscle relaxation.

-Then autolysis of the muscle proteins by lysosomes and bacterial putrefaction within **15-25** hours.

-It is important **medico legally** for determination of death time.

**2-Ca<sup>2+</sup> rigor:** increase in extracellular Ca<sup>2+</sup> may stop the heart in the systole.

**3-Heat rigor:** in case of increase body temperature as high fever lead to destruction of enzymes responsible for relaxation and Ca<sup>2+</sup>

**4-Physiological rigor: (Contracture)** means contraction not preceded by stimulation and not followed by relaxation because in extreme fatigue or untrained muscle → ↓ ATP → sustained contraction.

## N.B.: Difference between:

**Contraction:** Shortening of muscle , **preceded by** stimulation and **followed** by relaxation.

**Contracture:** Shortening of muscle, **not preceded** by stimulation, **not followed** by spontaneous relaxation.

**-Mechanical changes:** there are **2 types** of contraction:

**-In isometric contraction:** (CE) shortens but (SE) stretched and elongated so the total muscle length is **constant**.

**-In isotonic contraction:** CE shortens but (SE) **not stretched** → **shorten ms.**

	<b>Isotonic</b>	<b>Isometric</b>
- The muscle length	Decrease	Constant
- Muscle tension	Constant	Increased
- Energy	Great and used for work & waste heat	Less and is waste heat (no work)
- Sliding	Great sliding	less
- Duration	Longer	Short
- O <sub>2</sub> required	Greater	Less
- Heat production	Less	Great waste heat
-Mechanical efficiency and work done	25-30% e.g. contraction of biceps to lift objects	Zero (no work done) e.g. when the weight is too heavy to be moved.

## -Muscle tension:

- **Passive (rest) tension:** is that exerted by unstimulated muscle.
- **Active tension:** is the increase in tension during contraction.
- **Total tension** = passive t. + active t. in isometric cont.


-Muscle length: • **Resting length:** is the length of unstimulated muscle

- **Equilibrium length:** is the length of relaxed and detached muscle from bony attachment
- **Optimal length:** is the length at which we get the best contraction.

## -Relation between muscle length and tension:

**Starling law** states that “Within limits, the greater the initial muscle length (preload), the more will be the tension developed in isometric contraction” as increase in muscle length leads to increase number of cross bridges → ↑ tension.

But overstretching → ↓ number of **cross bridges** → ↓ tension. This law is applied also in the cardiac and smooth muscles as it is myogenic law (depend on the muscle not on the nerve supply).




## -Relation between the load and velocity of contraction:

-The velocity of contraction is **inversely** proportional to the load carried as in the cardiac muscle.

-Maximal velocity is obtained when the load is zero.

-Zero velocity is obtained when the load is more than power of contraction.



Thank You