

وسهلا



أهلا

يُمنع أخذ السلايدات بدون
إذن المحرر واي اجراء
يخالف ذلك يقع تحت طائلة
المسؤولية القانونية
جميع المعلومات للاستخدام
التعليمي فقط

الأستاذ الدكتور يوسف حسين

رئيس قسم التشريح والأنسجة والأجنة

كلية الطب - جامعة مؤتة - الأردن

دكتورة من جامعة كولونيا المانيا

Prof. Dr. Youssef Hussein Anatomy - YouTube

الواتس 00201224904207

SPERMATOGENESIS

البيوتوب د. يوسف حسين

- ** Definition:** It is the process of formation of the sperms (mature male gametes) from the primordial germ cells (spermatogonia)
- **Time:** starting at puberty (15-17 years) and continues till old age.

epididymis



seminiferous tubules

- Site of spermatogenesis
- in seminiferous tubules in testis and sperms stored in epididymis.
- **Testis** primary Male sex gland, Located in the Scrotum.
- Produce Sperms.
- Testis have interstitial cells (**Leydig cells**) that produce male sex hormone (**testosterone** or androgen hormone).
- Seminiferous tubules is structural unit of testes, site of developmental phases of sperms, containing **Sertoli cells**.

اليوتوب د. يوسف حسين

❖ Functions of Sertoli cells

1. Secrete protein for **nutrition** of developing sperms So called **mother cells or Nurse cells**
2. **Phagocytosis** for residual cytoplasm from spermatogenesis.
3. Maintain the **environment** of sperms via forming a **blood-testis barrier**.
4. **Secrete anti-Müllerian hormones** during early stage of fetal life which represses formation of derivatives of müllerian duct
5. **Secrete inhibin and activin** which controls pituitary gland to regulate FSH
6. **Secrete androgen binding protein** leading to increase testosterone hormone to stimulate spermatogenesis

Stage of proliferation

Stage of growth


اليوثيوب د. يوسف حسين

1st meiotic division

Stage of maturation

2nd meiotic division

Stage of transformation



At puberty

** Stages of spermatogenesis

** It includes 4 stages:

اليوثيوب د. يوسف حسين

(1) Stage of proliferation

- Each primordial germ cell (**spermatogonium**, 46 chromosome) is divided by many rounds of **mitotic division** successively to form a large stock of **daughter spermatogonium** (each contains 46 chromosomes).

(2) Stage of growth

- A number of daughter spermatogonia acquired more cytoplasm and increased in size forming **primary spermatocyte** (46 chromosomes).
- **Other numbers of daughter spermatogonia** remain a source of new **Spermatogonium cells**.

(3) Stage of maturation

- After many mitotic divisions, The primary spermatocytes, divide by **meiotic division** as follows:


I) 1st meiotic division (reduction division): each primary spermatocyte divide by meiotic division into two **secondary spermatocytes** {23 chromosome (one contains 22+X and one 22+Y)}.

II) 2nd meiotic division (equational division): each of the developed secondary spermatocytes divide into two cells called **spermatids** (each contains 23 chromosomes 22+X or 22+Y).

N.B; Each **daughter spermatogonium** gives 4 spermatids (2= 22+X and 2= 22+Y).

اليوتيوب د. يوسف حسين


<https://www.youtube.com/@ProfDrYoussefHusseinAnatomy/playlists>



Spermatid

الـيـوتـيـوب د. يوسف حسين

Stage of transformation



Sperm

(4) Stage of transformation (Spermiogenesis):

- It occurs in male only.
- Transformation of the spermatid into a mature sperm.
- The **length** of the sperm is about **50- 60 micrometer**
 1. The **nucleus** forms the **head** of sperm.
 2. The **Golgi** apparatus forms **acrosomal cap** contains **hydrolytic enzymes** which covers the anterior 1/2 of the head.
 3. **Centrioles**,
 - a- **Anterior** centriole: in the neck, immediately behind the head.
 - b- **Posterior** centriole: forms a **ring** shaped structure at end of the body.
 - **Axial filament** arises from anterior centriole and passes through the ring shaped posterior centriole to the tail.
 4. **Mitochondria** forms a mitochondrial sheath around axial filaments between centrioles.
 5. **Cytoplasmic membrane** forms a cytoplasmic sheath around the body and tail.

الـيـوتـيـوب دـ. يـوسـف حـسـين

- **Characters of normal sperm**

- 1. Number of sperm** about 20-200 million per ml and survive on fructose.
- 2. Parts of mature sperm:** head, neck, body and tail.
- 3. Motile of sperm at ejaculation:** more than 80%.
- 4. Rate of movement of sperms** in the female genital tract about 1- 3 mm per minute.
- 5. Survival of sperm** in the female genital tract about 3-4 days.
- 6. The average volume of semen** at ejaculation is 3-5 ml
- 7. Appearance** is whitish to gray
- 8. PH** is 7.2 – 8.0 (If low is acidic while high is alkaline)
- 9. The seminal fluid is secreted by** testis, seminal vesicle, prostate, and bulbourethral glands (Cooper's gland)

اليوثيوب د. يوسف حسين

- **Abnormal spermatogenesis**

- i) **Azoospermia:** absence of sperms in the seminal fluid
- ii) **Oligospermia:** decreased number of sperms in the seminal fluid
- iii) **Asthenospermia** reduced sperm motility
- iv) **Necrospermia: sperms found dead.**
- v) **Genetic abnormalities:** Sperm having abnormal chromosomal content
- vi) **Morphological Abnormalities:** If more than 20% affect fertility
 - Giant. - Dwarf. - Joined in head or in tail.



اليوثوب د. يوسف حسين

Oogenesis

البيوتوب د. يوسف حسين

**** Definition:** It is a process by which mature ovum is formed from primitive germ cell (oogonium)

• OOGENESIS

dr_youssefhussein@yahoo.com

**** Site:** in the cortex of ovary in female.

**** Time:**

- It **started** during foetal (intrauterine) life
- **Continues** after puberty
- **Completed** after fertilization
- **Arrested** at the age of menopause.

اليوثيوب د. يوسف حسين

- Many of the **primary oocytes degenerate before birth**
- **At birth, the ovary contains about two million primary oocyte,** most of them degenerate and, by puberty, when ovulation begins only about **300,000- 400,000 primary oocytes** are left in the ovary.

Before birth

Stage of Proliferation

daughter

Daughter oogonium

Stage of growth

Primary oocyte

primary oocyte

اليوتوب د. يوسف حسين

1st meiotic division

At puberty (ovulation)

Stage of maturation

Secondary oocyte

2ry oocyte and 1st polar body

2nd meiotic division


At fertilization

Ootid

First polar body

Ovum and 2nd polar body

Second polar bodies



**** Stages of Oogenesis**

A- During foetal life

dr_youssefhussein@yahoo.com

(1) Stage of proliferation:

- Each primordial germ cell; {**oogonium**} (46 chromosome) is divided by mitosis into 2 **daughter** oogonia (each contains 46 chromosomes).

(2) Stage of growth:

- Each of the daughter oogonia increases in size forming the **primary oocyte** containing 46 chromosomes (at birth).

B- After puberty

اليوثيوب د. يوسف حسين

(3) Stage of maturation:

- I) **1st meiotic division (reduction division):** each primary oocyte divide by meiotic division into one **secondary oocytes** (contains 22 +X chromosomes and **1st polar body** (contains 22 +X chromosomes and minimal amount of the cytoplasm)).

C- After fertilization

II) 2nd meiotic division (equational division):

- The secondary oocyte divides into **mature ovum** (contains 23 chromosome, 22 + X and most of the cytoplasm) and **2nd polar body** (contains 22 +X chromosomes and minimal amount of the cytoplasm) in the Fallopian tube.
- The 1st polar body divided by **secondary meiotic division** into two **2nd polar bodies**.

اليوثيوب د. يوسف حسين

N.B;

- The role of the polar body is to reduce the number of chromosomes. Later on decay and disintegrated

<https://www.youtube.com/@ProfDrYoussefHusseinAnatomy/playlists>

• Differences between spermatogenesis and oogenesis


| | Spermatogenesis | Oogenesis |
|-------------------|--|---|
| Definition | Formation of the sperms from the primordial germ cells. | Formation of the ova from the primordial germ cell. |
| Site | In the seminiferous tubules of testis | In the cortex of the ovary |
| Time | - It started at puberty and continues till very old age (all over life). <div style="border: 2px solid red; padding: 5px; display: inline-block;"> اليوتيوب د. يوسف حسين </div> | - It started during intrauterine life, and continues after puberty to be completed after fertilization. - It arrested at the age of menopause. |
| Hormones | - Follicle stimulating hormone, Luteinizing hormone, and testosterone | - FSH and LH |
| Stages | Proliferation, growth, maturation and transformation. | Proliferation, growth, maturation (No transformation stage) |
| Results | Each daughter spermatogonium gives 4 sperms. | Each daughter oogonium gives 1 ova and 3 polar bodies. |



https://www.youtube.com/channel/UCVSNqbibj9UWYaJdd_cn0PQ

يُمنع أخذ السلايدات بدون
إذن المحرر واي اجراء
يخالف ذلك يقع تحت
طائلة المسؤولية القانونية

اليوتيوب د. يوسف حسين



<https://www.youtube.com/@ProfDrYoussefHusseinAnatomy/playlists>