

# **PERIPHERAL NERVOUS SYSTEM**

**CN IX , X , XI ,XII**

**Dr. Aiman Qais Afar**  
**Surgical Anatomist**


**College of Medicine / University of Mutah**  
**2023-2024**

**Tuesday 5 March 2024**

# Glossopharyngeal Nerve CN IX


**Nuclei:** Four nuclei in the medulla send or receive fibers via CN IX:

- ✓ **two motor** (nucleus ambiguus and inferior salivary nucleus) and
- ✓ **two sensory** (sensory nuclei of the trigeminal nerve [CN V] and nuclei of the solitary tract).
- ✓ **Three of these nuclei are shared with CN X**



# Glossopharyngeal Nerve CN IX


- ❖ The glossopharyngeal nerve is a **motor** and **sensory nerve**
- ❖ It emerges from the anterior surface of the medulla oblongata **between the olive and the inferior cerebellar peduncle.**
- ❖ It passes laterally in **the posterior cranial fossa** and leaves the skull by passing **through the jugular foramen.**



# Glossopharyngeal Nerve CN IX

❖ The **superior** and **inferior sensory ganglia** are located on the nerve as it passes through the foramen.

❖ The glossopharyngeal nerve then descends through the upper part of the neck **to the back of the tongue**



# Functions:




❖ Somatic (Branchial) Motor

❖ Somatic (General) Sensory

❖ Visceral (Parasympathetic) Motor

❖ Special Sensory (Taste)


❑ Visceral Sensory



# Glossopharyngeal Nerve CN IX

❑ Somatic (Branchial) Motor


❑ Motor fibers pass to one muscle, the **stylopharyngeus**, derived from the 3rd pharyngeal arch.



# Glossopharyngeal Nerve CN IX

❑Visceral (Parasympathetic) Motor

❑Following a circuitous route initially involving **the tympanic nerve**, presynaptic parasympathetic fibers are provided to **the otic ganglion** for innervation of the parotid gland.




# Glossopharyngeal Nerve CN IX

## □ Somatic (General) Sensory

✓ The tympanic nerve.

Via the tympanic plexus, CN IX supplies the:

- mucosa of the tympanic cavity,
- pharyngotympanic tube, and
- the internal surface of the tympanic membrane.






# Glossopharyngeal Nerve CN IX

Dr. Aiman Al Maathidy  
Tuesday 5 March 2024

9

## □ Somatic (General) Sensory

- ✓ The pharyngeal,
- ✓ tonsillar, and
- ✓ lingual nerves to the:
  - mucosa of the oropharynx and
  - isthmus of the fauces (L., throat), including palatine tonsil, soft palate, and
  - posterior third of the tongue.



# Glossopharyngeal Nerve CN IX

Dr. Aiman Al Maathidy



1  
0

In addition to **general sensation** (touch, pain, temperature), there is a **tactile (actual or threatened) stimuli** determined to be unusual or unpleasant here may evoke **the gag reflex** or even vomiting.

Pharyngeal "gag" reflex



Palatal reflex





Tuesday 5 March 2024

# Glossopharyngeal Nerve CN IX

Dr. Aiman Al Maathidy  
Tuesday 5 March 2024

## ❑ Visceral Sensory


- ✓ The **carotid sinus nerve** to the carotid sinus, a **baro-(presso-)** receptor sensitive to changes in blood pressure,
- ✓ and **the carotid body**, a **chemoreceptor** sensitive to blood gas (oxygen and carbon dioxide levels).



# Glossopharyngeal Nerve CN IX


## □ Special Sensory (Taste)

Taste fibers are conveyed from **the posterior third of the tongue** to the sensory ganglia, the superior and inferior ganglia of **CN IX**



# Vagus Nerve CN X


- ❖ The vagus nerve is composed of **motor and sensory fibers**
- ❖ It emerges from the anterior surface of the medulla oblongata **between the olive and the inferior cerebellar peduncle.**
- ❖ The nerve passes laterally through **the posterior cranial fossa** and leaves the skull **through the jugular foramen.**



# Vagus Nerve CN X

The vagus nerve has both **superior** and **inferior sensory ganglia**.


□ Below the inferior ganglion, **the cranial root of the accessory nerve joins the vagus nerve and is distributed mainly in its pharyngeal and recurrent laryngeal branches.**



# Vagus Nerve CN X


❑ The vagus nerve descends through the neck alongside **the carotid arteries** and **internal jugular vein** within **the carotid sheath**

❑ It passes through the mediastinum of the thorax passing behind the root of the lung, and enters the abdomen through the **esophageal opening** in the diaphragm



# Important Branches of the Vagus Nerve in the Neck

- ■ **Meningeal and auricular branches**
- ■ **Pharyngeal branch** contains nerve fibers from the **cranial part of the accessory nerve.**
- ■ **Superior laryngeal nerve** divides into the internal and the external laryngeal nerves.



- ✓ **The internal laryngeal nerve** is **sensory** to the piriform fossa and the larynx down as far as **the vocal cords.**
- ✓ **The external laryngeal nerve** is **motor** and it supplies **the cricothyroid muscle.**



# Important Branches of the Vagus Nerve in the Neck

## ■ Recurrent laryngeal nerve


On the right side, the nerve hooks around the first part of **the subclavian artery**

On the left side, the nerve hooks around the **arch of the aorta**

The nerve is closely related to **the inferior thyroid artery**, and it supplies all the muscles of the larynx, except the cricothyroid muscle, the mucous membrane of the larynx below the vocal cords, and the mucous membrane of the upper part of the trachea.

■ ■ Cardiac branches (two or three) arise in the neck, descend into the thorax, and end in the cardiac plexus

Tuesday 5 March 2024  
Dr. Aiman Al Maathidy




# Vagus Nerve CN X

Dr. Aiman Al Maathidy  
Tuesday 5 March 2024

The vagus nerve thus innervates:

- ✓ The heart and great vessels within the thorax
- ✓ The larynx, trachea, bronchi, and lungs
- ✓ Much of the alimentary tract from the pharynx to the splenic flexure of the colon.
- ✓ It also supplies glands associated with the alimentary tract, such as the liver and pancreas.



✓ The vagus nerve has the most extensive distribution of all the cranial nerves and supplies the fore mentioned structures with afferent and efferent fibers.

# Vagus Nerve CN X

## Summary (Vagus nerve function)

- ✓ Somatic sensory
- ✓ Visceral sensory from the thoracic and abdominal organs.
- ✓ Taste and somatic (general) sensation
- ✓ Somatic motor
- ✓ Proprioceptive to the muscles that innervate
- ✓ Visceral (parasympathetic) motor to thoracic and abdominal viscera.

# Accessory Nerve CN XI


Dr. Aiman Al Maathidy  
Tuesday 5 March 2024

- ❑ The accessory nerve is a motor nerve.
- ❑ It consists of a cranial root (part) and a spinal root (part)

## Cranial Root

Emerges from the anterior surface of the medulla oblongata **between the olive and the inferior cerebellar peduncle**


The nerve runs laterally in **the posterior cranial fossa** and joins the spinal root



# Accessory Nerve CN XI

## Spinal Root


- ✓ Arises from nerve cells in the anterior gray column (horn) of **the upper five segments of the cervical part of the spinal cord**
- ✓ The nerve ascends alongside **the spinal cord** and enters the skull through **the foramen magnum**. It then turns laterally to **join the cranial root**



# Accessory Nerve CN XI

❖ The two roots unite and leave the skull through **the jugular foramen**.

The roots then separate:



❖ The **cranial root** joins **the vagus nerves** and is distributed in its branches to the muscles of the soft palate and pharynx (**via the pharyngeal plexus**) and to the muscles of the larynx (**except the cricothyroid muscle**).


# Accessory Nerve CN XI

❖ The spinal root runs downward and laterally and enters the deep surface of the sternocleidomastoid muscle, which it supplies, and then crosses the posterior triangle of the neck to supply the trapezius muscle

The accessory nerve thus brings about:


✓ Movements of the soft palate, pharynx, and larynx

✓ Controls the movements of the sternocleidomastoid and trapezius muscles, two large muscles in the neck



# Hypoglossal Nerve CN XII

- ❖ The hypoglossal nerve is a **motor nerve**.
- ❖ It emerges on the anterior surface of the medulla oblongata between the pyramid and the olive, crosses **the posterior cranial fossa**, and leaves the skull through **the hypoglossal canal**.
- ❖ The nerve then passes downward and forward in the neck and **crosses the internal and external carotid arteries** to reach **the tongue**
- ❖ In the upper part of its course, it is **joined by C1 fibers from the cervical plexus**






# Important Branches of the Hypoglossal Nerve

- Meningeal branch
- Descending branch (C1 fibers) passes downward and joins the descending cervical nerve (C2 and 3) to form the ansa cervicalis.

Branches from this loop supply the omohyoid, the sternohyoid, and the sternothyroid muscles.


- Nerve to the thyrohyoid muscle (C1)



# Important Branches of the Hypoglossal Nerve


- ■ **Muscular branches** to all the muscles of the tongue except **the palatoglossus** (pharyngeal plexus)
- ■ **Nerve to the geniohyoid muscle (C1).**

**The hypoglossal nerve** thus innervates the muscles of the tongue (**except the palatoglossus**) and therefore **controls the shape and movements of the tongue.**




**VAGUS NERVE INJURY** Injury to **pharyngeal branches of CN X** results in dysphagia (difficulty in swallowing).

- ❑ Injury of the **recurrent laryngeal nerve** causes hoarseness and dysphonia (difficulty in speaking) because of paralysis of the vocal folds (cords).
- ❑ Paralysis of **both recurrent laryngeal nerves** causes aphonia (loss of voice) and inspiratory stridor (a harsh, high pitched respiratory sound). All these may result from **cancer of the larynx and thyroid gland** and/or from **injury during surgery on the thyroid gland, neck**.



27  
Dr. Aiman Al Maathidy  
Tuesday 5 March 2024



# SPINAL ACCESSORY NERVE INJURY

Because of its nearly subcutaneous passage through the posterior cervical region, **CN XI is susceptible to injury during surgical procedures** such as lymph node biopsy, cannulation of the internal jugular vein, and carotid endarterectomy result in **marked ipsilateral weakness of shoulder (Drooping of the shoulder)**

Test for sternocleidomastoid function



Red arrow:  
Direction of movement  
White arrow:  
Direction of resistance

Test for trapezius function





FIGURE B9.6. Muscle test for sternocleidomastoid and trapezius.



## Injury to Hypoglossal Nerve

Injury to CN XII **paralyzes the ipsilateral half of the tongue**. After some time, the tongue atrophies, making it appear shrunken and wrinkled. When the tongue is protruded, its apex **deviates toward the paralyzed side** because of the unopposed action of **the genioglossus muscle** on the normal side of the tongue



**FIGURE B9.7.** Hypoglossal (CN XII) nerve lesion.

**Dr. Aiman Qais Afar**  
**Tuesday 5 March 2024**

