



Neurophysiology of smell and taste (gustation)

Dr. Arwa Rawashdeh




Physiology of taste





Taste Bud

Mechanism of stimulation




Gustatory Nerves

Pharynx, Epiglottis,
Esophagus

1/3 of anterior
tongue innervated by
glossopharyngeal nerve


2/3 of anterior tongue
innervated by tympanic
nerve



Vagus
Nerve (X)

Glossopharyngeal
Nerve (IX)

Facial
Nerve (VII)





- Geniculate ganglion

Herpes zoster, also known as shingles, is caused by the reactivation of the varicella-zoster virus (VZV), the same virus that causes varicella (chickenpox)



- Bell's palsy is an unexplained episode of facial muscle weakness or paralysis. This condition results from damage to the facial nerve (the 7th cranial nerve)


Physiology of olfaction





- A cilia of specific cilia express a specific protein receptors which can respond to different odorants
- One odor can bind to many different types of olfactory receptors protein
- G olfactory protein bind to GDP . But gets rid of GDP and binds with GTP and become very active
- Bind with AC that makes ATP converts into c AMP and bind to sodium channels and flow in sodium and calcium (adaptation response) and leaving chloride (component of mucus layer)





Granule cells connected glomerular with mitral and when they get excited, they releasing GABA inhibitory neurotransmitter .Granule cells neurons are interneurons that are thought to be involved with fine-tuning the processing of olfactory information by doing things like helping to sharpen the contrast between different odorants.

Lateral olfactory area gives branches to :

1. Deep part in the temporal lobe ; the incus and supply the piriform cortex

2. Limbic system

Entorhinal cortex (EC) , Hippocampus (memories)

Amygdale (emotion)

Medial olfactory area gives branches to :

Subcallosal gyrus

Orbital frontal cortex

Some small fibers can cross over

Other fibers are ipsilateral

So smell can be bilateral

All the olfactory nerves and taste nerves are intermingled

Taste is 80% smell

Anosmia

Nasal infection

Paranasal sinus infection

Olfactory groove meningiomas

Trauma rhinorrhea

sign of neurodegenerative disease