

Genomic DNA Extraction



Dr. Nesrin Mwafi

Biochemistry & Molecular Biology Department

Faculty of Medicine, Mutah University



Part I

Introduction

Extraction of DNA



- **DNA Extraction:** is a process of isolation and purification of DNA from a sample using a combination of physical and chemical methods
- Routine procedure widely used in: (downstream(pathway , application)):
 1. Molecular biology labs (scientific research labs) for example to study a gene involved in a cancer .



In purification
We used
ependorf
tube

Extraction of DNA



2. Genetic testing : is used to diagnose or rule out suspected genetic or inherited diseases. Also to identify disease carriers or to predict those at high risk for specific conditions



- Adult
- Newborn
- Fetus



Extraction of DNA




3. Forensic analyses : to gather evidences from the **crime scene that can be used in the court**



DNA Fingerprinting



- DNA fingerprinting (DNA profiling or typing):
- is a technique to identify individuals by features of their DNA .
- 99.9% of Human DNA sequences are identical in every person. Only 0.1% of our DNA
- sequences that make us unique .
- (ALL human organ, which in turn perform the same functions)
- DNA profiles: are small set of DNA variations that are very likely to be different in all unrelated individuals (except identical twins). مهم جداً
- DNA fingerprinting is used in criminal investigations and paternity or parentage testing .




DNA Fingerprinting



Victim

Crime Scene



Suspect 1

Suspect 2

Suspect 3



Mother

Child



'Dad' 1

'Dad' 2

'Dad' 3




Types of DNA



- We need cells to isolate DNA. In eukaryotes (multicellular) like human (*Homo sapiens*), **genomic DNA or chromosomal DNA** is found inside the nucleus of every cell (exceptions like the anucleated RBCs). Also, DNA is found within mitochondria (**mitochondrial DNA**), the energy producing organelles.

- Mitochondrial DNA is 16Kbp small circular dsDNA (2-10 copies) codes for 37 genes (inherited from mother).
- Genomic DNA contains: genetic data of organism with size of 300 million bp and codes for 25,000 genes arranged on 23 chromosomes (haploid cell, gametes/sex cells).


Chromosomal DNA



Types of DNA




- Somatic cell (diploid) consists of 23 pairs of chromosomes (46 chromosomes) code for different 25,000 genes (duplicate, one copy from each parent) with total size of 600 million bp.
- Genomic DNA is inherited from both parents (half from the mother and half from the father).
- In diploid cell, if DNA is laid out end to end, the total length will be approximately 2 meters. (located in nucleus which size $6\mu\text{m}$) .
- We have enough DNA to make 70 round trips from earth to sun .
- Human genome project (1990-2003) .
- 98.5% of human genome is junk coding DNA (1.5% only is coding DNA) . مهم جداً



DNA Packaging




- DNA is arranged into three different levels of organization:
 1. **Nucleosome**: DNA is wrapped around histone proteins (octamer)
 2. **Chromatin fibers**: nucleosomes coil and form loops
 3. **Chromosomes**: chromatin fiber is further condensed



Bacterial DNA



- In prokaryotes (unicellular) like bacterial cells, there are a single circular chromosome (bacterial DNA) found in the cytoplasm (with size of 5 million bp).
- Extrachromosomal DNA called plasmid also found in the cytoplasm: جداً مهم
 1. Small circular dsDNA (e.g. A.B resistance genes)
 2. Replicate independently of chromosomal DNA .
 3. Hundred of copies of single plasmid .
 4. Synthetically modified plasmids are used as vectors .



DNA Extraction Kits



- Obtaining high quality (***purity***) and quantity (***concentration/amount***) of ***intact DNA (not degraded)*** is often the first and most critical step in many fundamental molecular biology applications, such as DNA cloning, sequencing, PCR and electrophoresis.
- Different kinds of kits are available from different companies: **Qiagen**, **Invitrogen**, **Promega** and **Bio Basic Inc.**

DNA Extraction Kits



DNA Isolation from Various Samples




- DNA can be extracted from variety of samples:

1. Human Samples



Forensic Sample



Tissues



Cells



Buccal swab
From squamous epithelium cell




**Embryonic or Fetal biopsy for
Prenatal genetic testing**

Human Samples



- Genomic material can be extracted from different human samples like cells or tissues (extraction from the liver tissue is easier than spleen or brain tissues depending on the biochemical nature of that tissue).
- Buccal swab (cheek cells): is the most convenient and easiest way (non-invasive) because the squamous epithelial cells of buccal mucosa divide every 24 hours.
- Fetal sample (screening/diagnostic): cells of the placenta (in late first trimester) or from amniotic fluid (in second trimester) to do chromosomal analysis.



DNA Isolation from Various Samples




- In Vitro Fertilization (IVF) technique:

Preimplantation Genetic Diagnosis (PGD) test



Three days embryo biopsy
for PGD test



DNA Isolation from Various Samples



- **Non-invasive prenatal testing (NIPT)** is a test to determine the risk that the fetus will be born with certain abnormalities like Down syndrome (trisomy 21) .
- A blood sample from pregnant woman is **taken at the beginning of week 10.**
- Free DNA molecules from placenta and mother are circulating in mother's blood
- Placental DNA is identical to fetal DNA



DNA Isolation from Various Samples




- DNA can be extracted from variety of samples:

1. Human Samples



Forensic Sample



Tissues



Blood



Cells



Buccal swab




**Embryonic or Fetal biopsy for
Prenatal genetic testing**

Human Samples




- **In vitro fertilization (IVF):** is a technique of doing fertilization of eggs in the lab.
- One cell sample can be taken from the zygote to perform genetic testing **called PGD.**
- **Blood sample is more common although RBCs are anucleated .**
- **After centrifugation of the sample, we take the buffy coat layer containing the WBCs** مهم جداً

Blood Sample



centrifugation



Plasma (55% of total blood)
Proteins, electrolytes, clotting factors, hormones , etc...

Buffy coat (<1% of total blood)
contains WBC's

RBC's (45% of total blood)



Human Samples



- **In vitro fertilization (IVF):** is a technique of **doing fertilization of eggs in the lab.**
- A one cell sample can be taken from the zygote for genetic testing
- Blood sample is more common although RBCs are anucleated
- After centrifugation of the sample, we take the buffy coat layer containing the WBCs
- Nowadays, there are kits available for extraction of DNA from whole blood sample

DNA Isolation from Various Samples




Not important

غير مطالب



2. Animal cells/tissues (e.g. zebrafish fins)

- Genotyping for fishes



2. Plant material (e.g. banana and strawberry)

- Plant cell wall
- High levels of polysaccharides and polyphenols present in plant tissue lysate so negatively affects the quality of extracted DNA and may inhibit downstream reactions



DNA Isolation from Various Samples




4. Viral and Bacterial cells


Not important

غير مطالب

- Nasal swab like COVID-19
- Oral swab (throat)
- Blood sample (serum)
- Stool sample



Viral cells



Bacterial cells

DNA Isolation from Various Samples




مهم جداً

5. Plasmid DNA (containing the gene of interest)

- After transformation of bacterial cells (competent cells), the amplified plasmid is extracted using different types of kits

- Miniprep (50-100 μg) kits.
- Midiprep (100-350 μg) kits.
- Maxiprep (500-850 μg) kits.

- The purified plasmid is stored as stock at $-20\text{ }^{\circ}\text{C}$.



Extraction happened in : cell , tissue , Embryo, Blood , Animal , plant plasmid DNA.