### Archive Lecture 1

Collected By:

Medical card

Date of



1. Which of the following pairs is mismatched ?

- Flucloxacillin - atypical pneumonia (Explanation: Atypical pneumonia is caused by organisms that either lack a cell wall or have cell walls that beta-lactams cannot target effectively, so beta-lactams aren't an effective treatment.)

2.Which of the following drugs is not commonly used to treat community-acquired pneumonia?

A) AmoxicillinB) AzithromycinC) LevofloxacinD) Tetracycline

**Answer: D** 

3.All of the following are drugs used in acute bronchitis except:
A) Clarythromycin
B) Amphotericin B
C) Erythromycin
D)Trimethoprim sulfamethoxazole

**Answer: B** 

4.Which of the following conditions can co-trimoxazole be used to treat? A) Otitis media

- B) Sinusitis<sup>e of</sup>
- C) Pharyngitis
- D) Acute bronchitis

**Answer: B** 

Archive Lecture 2

Collected By:

Medical card

Date of



- 1. All the following are adverse effects of first-generation histamine antagonists except: A) Blurred vision
- B) Dry mouth
- C) Dizziness
- D) Diarrhea

**Answer: D** 

Archive Lecture 3

Collected By:

Medical card

Date of



- 1.All of the following are true regarding the treatment of acute severe asthma except: A) Humidified oxygen B) Nebulized B2 adrenergic agonist C) IV fluid D) Potassium and sodium bicarbonate E) Oral glucocorticoid Answer: D 2. Which of the following statements about the treatment of asthma is incorrect? A) Salmeterol can be used as a rescue inhaler during acute asthma attacks. B) Inhaled corticosteroids are effective for long-term control of asthma symptoms. C) Short-acting beta-agonists (e.g., albuterol) are preferred for immediate relief during an acute attack. D) Systemic corticosteroids are often used for severe asthma exacerbations. **Answer:** A 3. Patient with daily shortness of breath, loss of 40% of pulmonary function test, diagnosed with acute asthma attack with moderate severity. Best corticosteroid for this emergency: A) IV Methylprednisolone **B)** Prednisolone **C)** Budesonide
- D) Fluticasone
- E) Beclomethasone

**Answer:** A

70) A 6-years-old girl experiencing frequent allergic reactions, Which medication is recommended to inhibit the release of mediators that cause this?

A) TheophyllineB) Sodium CromoglycateC) LoratadineD) Prednisone

Ans:B

### Archive Lecture 4 Eicosanoids

Collected By:

Medical card

Date of

Malak Khaled



1. Which of the following is the best explains about COX1, COX2 : A)coxl is primary involved in inflammation ,cox2 responsible for normal physiological functions B)coxl is constitutional, cox2 inducible C)both cox1 & cox2 found only in brain D) cox2 is for gastric protection, cox1 is a thermal regulator **Answer: B** 2. One of the following is used for induced abortion or labor: A)epoprostenol B)pgI2 C)indomethacin D) pgE2 Answer: D 50)mechanism of action of LT receptor antagonist: A)chemical antagonism **B)**pharmacological antagonism **C)non competitive D**)partial E)physiological Ans: not sure

> -إذا كان يومُك أغلبه مذاكرة ، فأنت ممّن اصْطَفاهم اللَّهُ وأشغلهم بالعلمِ لا بغيره، وأنت ممّن سهّل اللَّهُ عليهِم الطريق إلىٰ الجنة، فأخلصْ في عملِك، بين يديك علمٌ ستنفع به الأمة.-

وتذّكّر قولهِ تعالى :: ﴿يَوْمَ تَشْهَدُ عَلَيْهِمْ أَلْسِنَتُهُمْ وَأَيْدِيهِمْ وَأَرْجُلُهُم بِمَا كَانُوا يَعْمَلُونَ﴾.

### Archive Lecture 5 TB

Collected By:

Medical card

Date of

Malaak Al Zaidaneeyen



1. \*All are adverae effect of isoniazid except;

Gout (high levels of uric acid)

2. Mechanism of action of isoniazid?

-interferes with mycolic acid synthesis

<u>3. Multi drug in TB therapy is to:</u> reduce resistance

<u>4.Patient with TB, after treatment it notice dark coloration in urine (red / orange), and new neuropathy symptoms. Which would be likely true:</u>

A ) the red coloration caused by isoniazid , but the deficincy in vitB6 caused by rifampin B ) the red coloration caused by isoniazid , but the deficincy in vitB6 caused by rifampin C)the red coloration caused by rifampin, but the deficiency in vitB6 leading to neuropathy by isoniazid

#### Ans:c

All of the following is true regarding rifampcin except :

Interfere with mycolic acid synthetic

Gout is a side effect of one of the following drugs:

#### pirazinamide



Lecture 5

Archive Lecture 6

Collected By:

Medical card

Date of



- 1.Which of the following is not typically used as an antitussive drug ?A) DextromethorphanB) Methadone
- C) Diphenhydramine
- D) Pholcodine

Answer: B

Answer: D

Answer: d

2.Which of the following statements is not true about dextromethorphan ?
A)Does not cause respiratory depression
B)Available in over-the-counter (OTC) preparations
C)Has antitussive effects
D)Has analgesic effects
E)Is not used for the treatment of cough due to asthma

All of the following is antitussive except:

a. Morphine b. Dextromethorphan c. Diphenhidramin D. Bromhexine

ard

Not related to Methadone:

A)long duration B)severe cough in lung cancer C)replacements of morphine addiction D)not used in children

Ans:c