



Biochemistry of Taste and Smell

Dr. Nesrin Mwafi


Biochemistry & Molecular Biology Department
Faculty of Medicine, Mutah University

Chemical Senses



- Taste (**Gustation**) and Smell (**Olfaction**) are chemical senses as they are stimulated by chemical compounds which bind the specialized receptors that they contain.
- Upon binding, the generated impulse is sent to the brain and then registered as a certain taste or smell
- The term “**flavour**” is used as an alternative to “**taste**” to avoid confusion with flavour referring to the overall perception that results from both taste and smell.

Detection of Flavour



Integration of odour and taste sensations
in the brain enables detection of flavour

Flavour = Taste + odour

Chemical Senses




- Taste (**Gustation**) and Smell (**Olfaction**) are chemical senses as they are stimulated by chemical compounds which bind the specialized receptors that they contain.
- Upon binding, the generated impulse is sent to the brain and then registered as a certain taste or smell
- The term “**flavour**” is used as an alternative to “**taste**” to avoid confusion with flavour referring to the overall perception that results from both taste and smell.
- Flavour of food protects us from poisons (mostly taste bitter) and spoiled food (taste sour).

TASTE



Gustatory System




Lingual Papillae



- There are 4 types of papillae scattered on the surface of human tongue:
 1. Circumvallate papillae: 8-12 dome or V-shaped papillae present at the back of the tongue
 2. Foliate papillae: leaf-like (4-5) short vertical folds located on the sides at the back of the tongue
 3. Fungiform papillae: mushroom-shaped projections mostly present on the tip and sides of the tongue
 4. Filiform papillae: threadlike structures scattered on the frontal 2/3 of the tongue. They are the only papillae that don't contain taste buds (mechanical papillae)
- Taste buds are also scattered throughout the mouth and throat (e.g. soft palate and epiglottis). These assist the primary taste buds located in papillae.

Taste Buds & Taste Receptors




Taste Buds & Taste Receptors



- Taste buds are flask-shaped structures surrounded by epithelial cells. They contain the taste receptor cells (**chemoreceptor cells**)
- Each taste bud is made up of 50-150 elongated cells arranged around a central taste pore:
 1. Taste receptor cells (TRCs): spindle-shaped neuroepithelial cells with microvilli projecting from the apical end whereas the other end is innervated by afferent nerve fibers. Each TRC has a particular receptor on its apical surface to bind specific tastant.
 2. Supporting cells: columnar cells that support TRCs
 3. Basal cells: cells that migrate from the adjacent lingual epithelium into the bud and differentiate into TRCs (regeneration time every 10 days)

Taste Signal Transmission



Taste Signal Transmission



Three cranial nerves are involved in the transport of taste message to the brain:

- Facial nerve (cranial nerve VII) carries taste sensations from the anterior two thirds of the tongue and soft palate
- Glossopharyngeal nerve (cranial nerve IX) carries taste sensations from the posterior one third of the tongue
- Vagus nerve (cranial nerve X) carries taste sensations from the back of the oral cavity (e.g. pharynx and epiglottis)

The Five Basic Tastes




There are 5 primary taste sensations:

1. **Sweet taste:** usually indicates energy rich nutrients (e.g. glucose)
2. **Salty taste:** allows modulating diet for electrolyte balance (e.g. NaCL)
3. **Sour taste:** typically the taste of acids (e.g. lemon)
4. **Bitter taste:** allows sensing of diverse natural toxins. Bitter tastants include alkaloids like quinine and caffeine
5. **Umami taste:** the taste of amino acid glutamate found in breast milk (10X) & many foods (e.g. meat, tomatoes, mushrooms, onions)



Tongue map

The Principle of Taste Stimulus Transduction



Molecular Mechanism of Taste Perception




- Taste sensation is realized in the mouth when the chemicals in food are dissolved by saliva and the free floating molecules enter a taste bud through the pore at its center
- Flavor molecules (tastants) fit into receptors on the microvilli at the top of the cell (TRC), causing either electrical changes (depolarization) that increase the intracellular level of Ca^{2+} ions (influx through voltage-gated Ca^{2+} channel) or by Ca^{2+} releasing from intracellular Ca^{2+} stores thus releasing neurotransmitters onto the nerve ending at the bottom of the cell
- The taste message is then carried to the brain by the appropriate cranial nerve: VII, IX or X

1. Salty Taste Perception




- With salty substances (e.g. table salt or NaCl), the Na^+ ions influx through amiloride -sensitive Na^+ channels expressed on the surface of cells in the tongue
- This causes the depolarization of the cell \rightarrow the influx of Ca^{2+} (voltage-gated Ca^{2+} channels) triggers the release of neurotransmitters at the synapse to the attached sensory neuron and fires an action potential




2. Sour Taste Perception

- Sour taste can result from either the passage of H^+ ions through **amiloride-sensitive Na^+ channels** or from the blockade of **K^+ channels**, which are normally open at resting potential → membrane depolarization
- These channels are permeable to protons when salivary Na^+ concentration is low. At high Na^+ concentration, protons block Na^+ flux through the channels and inhibit the response to NaCl.




3. Bitter Taste Perception

- Although at least one bitter stimulus “quinine” may depolarize taste cells by blocking apical K^+ channels, most bitter stimuli are thought to bind to G protein-coupled receptors (GPCRs) of the family **T2R receptors**.
- Binding of bitter tastant to T2R receptor stimulates the membrane-bound G-protein “gustducin” which activates phospholipase C (PLC) to increase production of inositol 1,4,5-trisphosphate (IP_3) as a second messenger that causes the release of Ca^{2+} from intracellular stores.



4. Sweet Taste Perception

- Sweet receptor is heteromeric GPCR made of two subunits: T1R2 and T1R3 proteins. This receptor can respond to both natural and non-carbohydrate artificial sweeteners (e.g. saccharin and aspartame)
- Substances that generate the sense of sweet flavour (e.g. fructose sugar) act on receptors that are coupled with G- proteins. Activation of G proteins results in activation of adenylate cyclase (AC) which increases the levels of cAMP (second messenger). cAMP activates a protein kinase (PKA) that depolarizes the receptor cells by closing K^+ channels at the basolateral membrane.
- Artificial sweeteners were found to act via the alternative signaling pathway : through the generation of the second messenger IP3 which induces the release of Ca^{2+} from intracellular stores




5. Umami Taste Perception



- Glutamate amino acid has a taste called *Umami* (from the Japanese word for “delicious”), that is distinct from the other four basic tastes.

- Umami taste is mediated by:

1. Metabotropic glutamate receptor 4 “ mGluR4”
a G-protein coupled receptor. mGluR4 is also expressed in certain regions of the brain.




The activation of these receptors causes Ca^{2+} to be released from intracellular stores such as the endoplasmic reticulum (ER).

2. Heteromeric GPCR consists of T1R1 and T1R3 proteins. This receptor is broadly tuned to the 20 standard L-amino acids found in proteins (but not to their D-amino acid enantiomers). Activation of these receptors also increases the intracellular level of Ca^{2+}

SMELL



Olfactory System



Olfactory System



- The olfactory system is located at the dorsal roof of the nasal cavity. It consists of 3 main structures:
 1. Olfactory bulb: it is an extension of the brain. It contains a bundle of nerves that carry the olfactory messages to the brain
 2. Cribriform plate of ethmoid bone: a bone that separates the olfactory bulb from the olfactory epithelium. It has holes through which the olfactory receptor neurons send projections (axons) to the olfactory bulb
 3. Olfactory epithelium: it consists of 3 different types of cells. Also, it contains the mucus secretory gland called “Bowman’s gland”.

Olfactory Epithelium




- The olfactory epithelium layer contains 3 types of cells:
 1. Olfactory receptor cells (olfactory sensory neurons): these are bipolar neurons with their axons extended at the basal surface towards the olfactory bulb (where they make synapses with mitral and tufted neural cells). At the apical surface, these cells have projections called cilia which extend into the underlying mucus layer containing the dissolved odorants. Cilia have the odorant receptors (proteins).
 2. Basal cells: responsible for regeneration of receptor cells every 6-8 weeks.
 3. Supporting cells

Olfactory Bulb



- The olfactory bulb contains structures called glomeruli (a spherical structures where the axonal ends of receptor cells cluster). Each glomerulus receives input primarily from olfactory receptor neurons that express the same olfactory receptors.
- Mitral cells and tufted cells are neurons located in olfactory bulb. They form synapses with receptor cells inside glomeruli. They send olfactory inputs into brain for integration.

Cellular Mechanism of Olfactory Transduction



Cellular Mechanism of Olfactory Transduction



- Each olfactory receptor cell is thought to contain a single type of olfactory receptor responding to a single type of odorant
1. The cilia of the olfactory sensory neurons are embedded in a layer of mucus in which odor molecules are dissolved.
 2. Binding of odorant ligands with G-protein-coupled receptors activates olfactory specific G-protein (G_{olf}) by removal of GDP and substitution with GTP .
 3. The active G-protein dissociates into β , γ and α -GTP subunits with the latter activating the enzyme adenylate cyclase.
 4. The adenylate cyclase increases the formation of cAMP from ATP.
 5. The increased amount of cAMP (second messenger) opens cAMP-gated cation channels causing the influx of Na^+ and Ca^{+2} ions

Cellular Mechanism of Olfactory Transduction



6. This causes membrane depolarization
7. The depolarization is further amplified by Ca^{+2} activated Cl^- channels (outflow of Cl^-)
8. The increased intracellular calcium ion concentration initiates calcium mediated exocytosis of neurotransmitter vesicles
9. The neurotransmitter molecules are released inside the synaptic cleft and diffuse towards the postsynaptic membrane (mitral or tufted cells of the olfactory bulb)