

1. **Which of the following statements is false?**
  - A- Drugs with molecular weight more than 1000 cross the placenta
  - B- Highly ionized drugs cross the placenta slowly
  - C- Drugs with high lipid solubility cross the placenta poorly **xx**
  - D- Warfarin should be avoided during pregnancy
  
2. **All the following statements are true EXCEPT:**
  - A- Drugs can harm the fetus anytime during pregnancy
  - B- Drugs may produce congenital abnormalities during first trimester
  - C- Drugs may affect growth and development of fetus during second trimester
  - D- Drugs may produce congenital abnormalities anytime during pregnancy **xx**
  
3. **All the following drugs are contraindicated in pregnancy EXCEPT:**
  - A- Thalidomide
  - B- Warfarin
  - C- Tetracycline
  - D- Heparin **xx**
  
4. **All the following drugs should be avoided during lactation EXCEPT:**
  - A- Chloramphenicol
  - B- Tetracycline
  - C- Radioactive iodine
  - D- Penicillin **xx**
  
5. **Which of the following statements is false?**
  - A- Enzyme systems that inactivate drugs are presents at birth
  - B- Clearance of drugs is low in the neonate
  - C- Drugs eliminated by kidney should be given in high dose **xx**
  - D- Glomerular filtration of drugs is low in the neonate
  
6. **All the following drugs are safe to give during pregnancy EXCEPT:**
  - A- Paracetamol
  - B- Insulin
  - C- Cephalosporins
  - D- Phenytoin **XX**
  
7. **All the following drugs should be avoided during lactation EXCEPT:**
  - A- Aspirin
  - B- Tetracycline
  - C- Radioactive iodine
  - D- Amoxicilin **xx**