

MORPHOLOGIC PATTERNS OF ACUTE INFLAMMATION

Eman Krieshan, M.D.

29-10-2025

- Special morphologic patterns are often seen in addition to the general features which are characteristic of most acute inflammatory reactions, depending on :
 - ❖ The severity of the reaction.
 - ❖ Its specific cause.
 - ❖ Particular tissue.
 - ❖ Site involved.
- ❖ They can provide valuable clues about the underlying cause.




1. SEROUS INFLAMMATION

- Marked by the exudation of cell poor fluid into spaces created by injury to surface epithelial or into body cavities such as peritoneal, pleural, or pericardial cavities.
- The fluid in serous inflammation is not infected by destructive organisms and does not contain large numbers of leukocytes
- Accumulation of fluid in these cavities is called an effusion.




❖ Peritoneal effusion



❖ SKIN BLISTER

- Resulting from a burn or viral infection.
- Represents accumulation of serous fluid within or immediately beneath the damaged epidermis of the skin



2. FIBRINOUS INFLAMMATION

- A fibrinous exudate develops when the vascular leaks are large or there is a local procoagulant stimulus.
- A fibrinous exudate is characteristic of inflammation in the lining of body cavities, such as the meninges, pericardium and pleura.



MECHANISM OF FORMATION

- Large increase in vascular permeability.




- higher-molecular weight proteins such as fibrinogen pass out of the blood.



- fibrin is formed and deposited in the extracellular space



❖ GROSSLY




The **pericardial surface** is **dry** with a **coarse granular appearance** caused by **fibrinous exudate**

Normally, the visceral **pericardium** is **translucent**




HISTOLOGY



Normal pericardium composed of thin fibrous wall
Covered by single layer of mesothelial cells





the pericardial surface here shows strands of pink fibrin extending outward. There is underlying inflammation.

fibrin appears as an eosinophilic meshwork of threads



3. PURULENT (SUPPURATIVE) INFLAMMATION, ABSCESS


- Purulent inflammation is characterized by the production of pus, an exudate consisting of neutrophils, the liquefied debris of necrotic cells, and edema fluid.
- The most frequent cause is infection with pyogenic (pus-producing) bacteria, such as staphylococci.



A COMMON EXAMPLE OF AN ACUTE SUPPURATIVE INFLAMMATION IS ACUTE APPENDICITIS



Acute appendicitis



Acute inflammation with predominance of neutrophils; involves some or all layers of the appendiceal wall.




- Abscesses:

- Localized collections of pus caused by suppuration buried in a tissue, an organ, or a confined space.
- They are produced by seeding of pyogenic bacteria into a tissue . In time the abscess may become walled off and ultimately replaced by connective tissue



Abscesses have multiple areas:

- * central region with necrotic leukocytes and tissue cells.
- * zone of preserved neutrophils around this necrotic focus.
- *vascular dilation, parenchymal and fibroblastic proliferation.




4. ULCERS




- An ulcer is a local defect, or excavation, of the surface of an organ or tissue that is produced by the sloughing (shedding) of inflamed necrotic tissue.
- Ulceration can occur only when tissue necrosis and resultant inflammation exist on or near a surface



- It is most commonly encountered in:
- (1) the mucosa of the mouth, stomach, intestines, or genitourinary tract.
- (2) the skin and subcutaneous tissue of the lower extremities in older persons




HISTOLOGY



sloughing (shedding) of inflamed necrotic tissue



Microscopic features of Ulcers



Acute stage:


Intense polymorphonuclear infiltration and vascular dilation in the margins of the defect.

With chronicity:

the margins and base of the ulcer develop fibroblast proliferation, scarring, and the accumulation of lymphocytes, macrophages, and plasma cells.



Outcomes of acute inflammation



OUTCOMES OF ACUTE INFLAMMATION


- Acute inflammatory reactions typically have one of three outcomes:
 - 1. Complete resolution:
 - Occur when the injury is limited or short-lived or when there has been little tissue destruction and the damaged parenchymal cells can regenerate.
 - Resolution involves removal of cellular debris and microbes by macrophages, and resorption of edema fluid by lymphatics.



- 2. Healing by connective tissue replacement (scarring, or fibrosis).
- occurs after substantial tissue destruction, when the inflammatory injury involves tissues that are incapable of regeneration, or when there is abundant fibrin exudation.
- connective tissue grows into the area of damage or exudate, converting it into a mass of fibrous tissue.
- 3. Progression of the response to chronic inflammation.
- occurs when the acute inflammatory response cannot be resolved, as a result of either :
 - the persistence of the injurious agent
 - interference with the normal process of healing



CHRONIC INFLAMMATION



- Chronic inflammation is a response of prolonged duration (weeks or months) in which:
 - inflammation.
 - tissue injury.
 - attempts of repair.
- coexist,
in varying combinations.**
- It may follow acute inflammation, as described earlier, or may begin insidiously,




CAUSES OF CHRONIC INFLAMMATION

- Persistent infections, e.g??.
- Hypersensitivity diseases.
- Autoimmune disease.
- Allergic diseases.
- Prolonged exposure to potentially toxic agents, e.g Silica.



Histopathology of chronic inflammation



- healing by connective tissue replacement of damaged tissue,



CELLS AND MEDIATORS OF CHRONIC INFLAMMATION

- Macrophages
- Lymphocytes




1. MACROPHAGES

- The dominant cells in most chronic inflammatory reactions .
- Become the dominant cell population in inflammatory reactions within 48 hours of onset.
- There are two major pathways of macrophage activation, (depends on the nature of the activating signals):
 - Classical:
 - designed to destroy the offending agents.
 - Alternative :
 - initiates tissue repair.




Macrophages are professional phagocytes.

Circulating
macrophage



Activation pathways




- They secrete growth factors that promote
- Angiogenesis.
 - activate fibroblasts.
 - stimulate collagen synthesis.

THE PRODUCTS OF ACTIVATED MACROPHAGES

- **Eliminate injurious agents such as microbes.**
- **Initiate the process of repair.**
- **Responsible for much of the tissue injury in chronic inflammation**





Lymphocyte

2. LYMPHOCYTES


- Microbes and other environmental antigen activate T and B lymphocytes, which amplify and propagate chronic inflammation.
- Some of the strongest chronic inflammatory reactions, such as granulomatous inflammation, are dependent on lymphocyte responses.



LYMPHOCYTES AND MACROPHAGES INTERACT IN A BIDIRECTIONAL WAY.



recruit and activate macrophages


display antigens to T cells



GRANULOMATOUS INFLAMMATION

- Granulomatous inflammation is a form of chronic inflammation characterized by collections of activated macrophages, often with T lymphocytes.
- Granuloma formation is a cellular attempt to contain an offending agent that is difficult to eradicate











Where??



TYPES OF GRANULOMAS;

- 1. Immune granulomas:
 - caused by persistent T cell–mediated immune response.
 - when the inciting agent cannot be readily eliminated.
- **2. Foreign body granulomas:**
 - seen in response to inert foreign bodies, in the absence of T cell– mediated immune responses.
 - May form around materials such as talc (associated with intravenous drug abuse), sutures, or other fibers





The foreign material can usually be identified in the center of the granuloma, particularly if viewed with polarized light, in which it may appear refractile.



SYSTEMIC EFFECTS OF INFLAMMATION

- Inflammation is associated with cytokine-induced systemic reactions that are collectively called the acute-phase response.
- The cytokines TNF, IL-1, and IL-6 are important mediators of the acute phase reaction.



THE ACUTE-PHASE RESPONSE CONSISTS OF SEVERAL CLINICAL AND PATHOLOGIC CHANGES:

- 1.Fever:
- Substances that induce fever are called pyrogens.
- caused by prostaglandins especially **PGE2** that are produced in the vascular and perivascular cells of the hypothalamus.

2.Acute-phase proteins

3.Leukocytosis

