

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

Question # 1

A 71-year-old man is admitted to the hospital one hour after he was found unconscious. His pulse is 80/min and systolic blood pressure is 98 mm Hg; diastolic blood pressure cannot be measured. He is intubated and mechanically ventilated with supplemental oxygen at a tidal volume of 450 mL and a respiratory rate of 10/min. Arterial blood gas analysis shows:

P _{CO2}	43 mm Hg
O ₂ saturation	94%
O ₂ content	169 mL/L

Pulmonary artery catheterization shows a pulmonary artery pressure of 15 mm Hg and a pulmonary capillary wedge pressure of 7 mm Hg. Bedside indirect calorimetry shows a rate of O₂ tissue consumption of 325 mL/min. Given this information, which of the following additional values is sufficient to calculate the cardiac output in this patient?

	Answer	Image
A	End-tidal carbon dioxide pressure	
B	Total peripheral resistance	
C	Partial pressure of inspired oxygen	
D	Left ventricular end-diastolic volume	
E	Pulmonary artery oxygen content	

Hint

Cardiac output can be calculated using the Fick principle. One additional value is required for the formula if the rate of oxygen consumption and the arterial oxygen content are known.

Correct Answer

A - End-tidal carbon dioxide pressure

Explanation Why

End-tidal carbon dioxide pressure (PetCO₂) is used with the [partial pressure](#) of arterial oxygen (PaO₂) and [tidal volume](#) to calculate [physiologic dead space](#). [Dead space](#) volume represents the amount of inspired air that does not participate in [gas exchange](#); however, neither PetCO₂ nor [dead space](#) volume can be used to determine [cardiac output](#).

B - Total peripheral resistance

Explanation Why

While [total peripheral resistance](#) (TPR) can be used to calculate [cardiac output](#) (CO) using the equation $CO = \text{mean arterial pressure} / TPR$, the necessary values are not given. To use this equation, [mean arterial pressure](#) (MAP) must first be determined. However, [diastolic](#) pressure (DBP) cannot be measured in this patient by sphygmomanometry; thus, the [MAP](#) is not calculable (recall that $MAP = \frac{2}{3} DBP + \frac{1}{3} SBP$). DBP may not be measurable via sphygmomanometry in patients with severe [hypotension](#) (as seen here), stiff [arteries](#), [arteriovenous malformation](#), or [aortic dissection](#). Therefore, another equation using the provided values should be utilized.

C - Partial pressure of inspired oxygen

Explanation Why

[Partial pressure](#) of inspired oxygen (PIO₂) is used in the [alveolar gas equation](#) to determine the partial pressure of oxygen in the alveoli, which is then used to calculate the [A-a gradient](#). The [A-a gradient](#) demonstrates the difference between the partial pressure of oxygen in the alveoli and the [arteries](#), which is useful to measure oxygenation or, conversely, determine the cause of [hypoxemia](#). However, neither PIO₂ nor the [A-a gradient](#) can be used to determine [cardiac output](#).

D - Left ventricular end-diastolic volume

Explanation Why

[Left ventricular end-diastolic volume \(EDV\)](#) can be used to find [stroke volume \(SV\)](#), and thus calculate [cardiac output \(CO\)](#) using the equation $CO = \text{heart rate} \times SV$. However, either [end-systolic volume \(SV = end-diastolic volume - end-systolic volume\)](#) or [ejection fraction \(SV = EF x EDV\)](#) is also required to determine [SV](#). Because these necessary values are not given in this patient, another equation should be utilized.

E - Pulmonary artery oxygen content

Explanation Why

[Pulmonary artery](#) oxygen content, i.e., the oxygen content of venous blood (CvO_2), is the missing value needed to calculate [cardiac output \(CO\)](#) in this patient. The CvO_2 can be measured via [catheterization of the right heart](#) or the [pulmonary artery](#). According to the [Fick principle](#), the CO is equal to the [ratio](#) of the rate of oxygen consumption (VO_2) to the arteriovenous difference, or $CO = (VO_2 / (\text{arterial } O_2 \text{ content} - \text{venous } O_2 \text{ content}))$. Given that VO_2 is 325 mL/min and arterial O_2 content is 169 mL/L, only venous O_2 content is needed to calculate this patient's CO.

Question # 2

A 30-year-old woman comes to the emergency department because of a 1-day history of sharp chest pain radiating to the left shoulder. The pain worsens when she coughs, breathes deeply, or lies down. She has had a mild fever and a cough for the past 2 days. She takes ibuprofen for hip and knee pain, which started 4 months ago. Examination shows a symmetric, erythematous rash on her nose and cheeks, and erythematous patches with scaling and plugged follicles over her hands. Cardiovascular evaluation of this patient is most likely to show which of the following findings?

	Answer	Image
A	Increased distension of jugular veins during inspiration	
B	18 mm Hg decrease in systolic blood pressure during inspiration	
C	High-pitched pericardial sound occurring slightly earlier than a third heart sound	
D	Triphasic scratchy heart sound of varying intensity	
E	Jugular venous pulsation visible 9 cm above the sternal angle	
F	Holosystolic blowing murmur heard best at the apex	

Hint

Joint pain, malar rash, and a discoid rash (erythematous patches with scaling and follicular plugging) should immediately raise suspicion of systemic lupus erythematoses (SLE). The acute onset of low-grade fever and pleuritic chest pain that is aggravated by lying down (and thus improved by sitting up) is suggestive of acute pericarditis, the most common cardiovascular manifestation of SLE.

Correct Answer

A - Increased distension of jugular veins during inspiration

Explanation Why

The [Kussmaul sign](#) is classically seen in [right ventricular infarction](#), [restrictive cardiomyopathy](#), [constrictive pericarditis](#), and tricuspid stenosis. [Constrictive pericarditis](#) may occur with [SLE](#) but would be characterized by signs of decreased [cardiac output](#) such as fatigue and [tachycardia](#). It would also take months to years to develop. The patient's acute presentation with [pleuritic chest pain](#), [cough](#), and [fever](#) is more consistent with [acute pericarditis](#), in which case the [Kussmaul sign](#) would not be expected.

B - 18 mm Hg decrease in systolic blood pressure during inspiration

Explanation Why

A decrease of 18 mm Hg in [systolic](#) blood pressure during inspiration suggests [pulsus paradoxus](#), which may be seen in [cardiac tamponade](#), [constrictive pericarditis](#), and in noncardiac diseases such as [asthma](#), [COPD](#), and [tension pneumothorax](#). Although [SLE](#)-associated [pericarditis](#) can be associated with a [pericardial effusion](#), it rarely progresses to tamponade. Furthermore, other findings of [cardiac tamponade](#), including [dyspnea](#), [hypotension](#), [tachycardia](#), and [jugular venous distention](#), would be expected. [Constrictive pericarditis](#) is also rarely associated with [SLE](#) and would cause signs of decreased [cardiac output](#) (fatigue) and fluid overload ([jugular venous distention](#), [Kussmaul sign](#)).

C -

High-pitched pericardial sound occurring slightly earlier than a third heart sound

Explanation Why

[Pericardial knock](#) is a high-pitched, early-[diastolic](#) sound in patients with [constrictive pericarditis](#).

[Constrictive pericarditis](#) may occur with [SLE](#) but would be characterized by signs of decreased [cardiac output](#) such as fatigue and [tachycardia](#). It would also take months to years to develop. The patient's acute presentation with [pleuritic chest pain](#), [cough](#), and [fever](#) is more consistent with [acute pericarditis](#), in which case the [Kussmaul sign](#) would not be expected.

D - Triphasic scratchy heart sound of varying intensity

Explanation But

[ECG](#) changes include widespread [ST-segment elevation](#) and/or PR depression.

Explanation Why

A triphasic scratchy or squeaky [heart sound](#) on auscultation suggests a [pericardial friction rub](#), which is highly specific for [pericarditis](#). Acute [pericarditis](#) typically manifests with [fever](#) and [pleuritic chest pain](#) that is usually sharp in nature, improves on sitting and leaning forward, and often radiates to the shoulder. It is the most common cardiac manifestation of [SLE](#), occurring in ~ 25% of patients.

E - Jugular venous pulsation visible 9 cm above the sternal angle

Explanation Why

Elevated jugular venous pulsation can be seen in [cardiac tamponade](#) or [constrictive pericarditis](#), both of which are uncommon in [SLE](#). Furthermore, other findings of [cardiac tamponade](#), including [dyspnea](#), [hypotension](#), [tachycardia](#), and [jugular venous distention](#), would be expected. [Constrictive pericarditis](#) would cause signs of decreased [cardiac output](#) (fatigue) and fluid overload ([jugular venous distention](#), [Kussmaul sign](#)).

F - Holosystolic blowing murmur heard best at the apex

Explanation Why

[Holosystolic](#) blowing [murmur](#) best heard at the apex suggests [mitral regurgitation](#). [Libman-Sacks endocarditis](#), which is caused by noninfectious [verruccous vegetations](#) and is seen in [SLE](#), favors the

[mitral valve](#) and can result in [mitral regurgitation](#). Most patients, however, are asymptomatic. This patient's [fever](#), [cough](#), and [pleuritic chest pain](#) are more consistent with [acute pericarditis](#), which is also the most common cardiac manifestation of [SLE](#).

Question # 3

A 59-year-old man is brought to the emergency department one hour after developing shortness of breath and “squeezing” chest pain that began while he was mowing the lawn. He has asthma, hypertension, and erectile dysfunction. Current medications include salmeterol, amlodipine, lisinopril, and vardenafil. His pulse is 110/min and blood pressure is 122/70 mm Hg. Physical examination shows diaphoresis. An ECG shows sinus tachycardia. Sublingual nitroglycerin is administered. Five minutes later, his pulse is 137/min and his blood pressure is 78/40 mm Hg. Which of the following is the most likely mechanism of this patient's hypotension?

	Answer	Image
A	Bradykinin accumulation	
B	Calcium channel antagonism	
C	Alpha-1 receptor antagonism	
D	Cyclic GMP elevation	The diagram illustrates the mechanism of nitroglycerin-induced vasodilation. It shows an endothelial cell producing NO from L-arginine via eNOS. NO diffuses into a smooth muscle cell, where it activates guanylyl cyclase, converting GTP to cGMP. cGMP then inhibits PDE5, leading to an increase in cGMP levels. This elevated cGMP inhibits the MLCK-myosin pathway, resulting in smooth muscle relaxation and vasodilation. The diagram also shows that nitrates and PDE5 inhibitors can enhance this pathway. A legend indicates: Red box = Effects of cGMP, Yellow box = Breakdown of cGMP, Green box = Physiological NO production.
E	Decreased nitric oxide production	
F	Histamine release	

	Answer	Image
G	Beta-2 receptor agonism	

Hint

Vardenafil is a PDE-5 inhibitor and should not be used in combination with nitrates.

Correct Answer

A - Bradykinin accumulation

Explanation Why

ACE inhibitors (e.g., [lisinopril](#)) can lead to the accumulation of [bradykinin](#), which can cause [hypotension](#) (due to vascular dilation). However, an acute, life-threatening drop in blood pressure is unlikely. Furthermore, if an accumulation of [bradykinin](#) had occurred in this patient, symptoms such as bronchoconstriction, [angioedema](#), and dry [cough](#) would also be expected.

B - Calcium channel antagonism

Explanation Why

[Calcium channel antagonists](#) (e.g., [amlodipine](#)) can cause vascular [smooth muscle](#) relaxation, which may lead to a drop in blood pressure. However, this class of drugs rarely causes acute, severe [hypotension](#), even when taken in conjunction with [nitroglycerin](#).


C - Alpha-1 receptor antagonism

Explanation Why

Alpha-1 antagonists (e.g., [prazosin](#)) cause peripheral [vasodilation](#) by inhibiting the [alpha-1 receptor](#) in [smooth muscle cells](#). This can also cause [hypotension](#), which is why this class of drugs is contraindicated in patients who take [nitrates](#). However, this patient is not currently taking any alpha-1 antagonists.

D - Cyclic GMP elevation

Image



Explanation Why

Nitrates (e.g., sublingual [nitroglycerin](#)) and **PDE-5 inhibitors** (e.g., [vardenafil](#)) both work by increasing the intracellular concentration of **cGMP**. **Nitrates** increase intracellular NO, which is responsible for activating [guanylate cyclase](#), leading to increased synthesis of **cGMP**. **PDE-5 inhibitors**, as their name implies, inhibit [phosphodiesterase \(PDE\)](#), which leads to a decreased breakdown of **cGMP**. Accumulation of **cGMP** leads to activation of [myosin](#)-light-chain [phosphatase \(MLCP\)](#), which dephosphorylates the [light chains](#) of [myosin](#). This results in [smooth muscle cell](#) relaxation in blood vessels with subsequent [vasodilation](#). Life-threatening [hypotension](#) can occur if these two drugs are taken in combination.

E - Decreased nitric oxide production

Explanation Why

[Nitroglycerin](#), a [nitric oxide](#) (NO) donor, causes [vasodilation](#), which leads to [smooth muscle](#) relaxation via a process that starts with the activation of [guanylyl cyclase](#). So NO production would be increased rather than decreased in this patient.

F - Histamine release

Explanation Why

A release of [histamine](#), which acts as a peripheral [vasodilator](#), is the mechanism underlying [hypotension](#) in [anaphylactic shock](#). [Anaphylactic shock](#) could account for this patient's [hypotension](#) and [tachycardia](#), but he lacks other typical features of [anaphylactic reactions](#), such as [urticaria](#), [angioedema](#), wheezing, nausea, and vomiting. Furthermore, [allergic reactions](#) to [nitroglycerin](#) are rare.



G - Beta-2 receptor agonism

Explanation Why

[Beta-2 receptor agonists](#) (e.g., [salmeterol](#)) cause relaxation of the [smooth muscle](#) in bronchial tubes and blood vessels. They are primarily used as bronchodilators in the management of [asthma](#) and other [obstructive pulmonary diseases](#) (e.g., [COPD](#)). Although [smooth muscle cell](#) relaxation leads to [vasodilation](#), severe [hypotension](#) is rare, even when [beta-2 receptor agonists](#) are taken in combination with [nitrates](#).

Question # 4

A 27-year-old woman comes to the physician because of a 1-year history of progressive shortness of breath. She is now unable to jog for more than 10 minutes without stopping to catch her breath. Cardiac examination shows a harsh systolic, crescendo-decrescendo murmur best heard at the lower left sternal border. The murmur increases in intensity when she moves from a squatting to a standing position and decreases when she clenches her fists. The lungs are clear to auscultation. Which of the following is the most likely cause of her condition?

	Answer	Image
A	Dystrophin defect	
B	Fibrillin-1 defect	 Marfan syndrome Epidemiology Prevalence: 1/5,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15, autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissections) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups) Features and complications shown in the diagram: - Ectopia lentis, severe myopia - Heart valve defects - Pectus deformity (pectus carinatum or pectus excavatum) - Aortic disease (e.g., aortic dissection, aortic aneurysm) - Arachnodactyly, joint hypermobility - Hyperextensible skin - Tall stature - Berry aneurysms - High-arched palate - Spiral deformities (scoliosis, hyperkyphosis) - Long extremities - Pes planus, hindfoot valgus
C	Myosin heavy chain defect	 The diagram illustrates the difference between a normal heart and two types of hypertrophic cardiomyopathy (HCM): the nonobstructive type and the obstructive type. In the obstructive type, the interventricular septum is hypertrophied, leading to obstruction of the left ventricular outflow tract (LVOT). This obstruction is caused by the systolic anterior motion (SAM) of the mitral valve, which is pulled into the LVOT during systole. Labels include: Normal, Hypertrophic cardiomyopathy, Nonobstructive type, Obstructive type, Left ventricular outflow tract (LVOT) obstruction due to: Interventricular septum hypertrophy, and Systolic anterior motion (SAM) of the mitral valve.
D	GAA trinucleotide repeats	

	Answer	Image
E	Myocyte ion channel defect	
F	CTG trinucleotide repeats	
G	Deletion in chromosome 7q	

Hint

A systolic crescendo-decrescendo murmur at the left sternal border indicates left ventricular outflow tract (LVOT) obstruction. This patient's LVOT obstruction is dynamic, i.e., it increases with maneuvers that decrease venous return (standing from squatting) and decreases with maneuvers that increase afterload (clenching fists). This finding suggests hypertrophic cardiomyopathy.

Correct Answer


A - Dystrophin defect

Explanation Why

Dystrophin defects are responsible for **Duchenne muscular dystrophy (DMD)**, which is associated with **dilated cardiomyopathy**. Patients with **dilated cardiomyopathy** can present with exertional dyspnea and a **systolic mitral regurgitation (MR) murmur**. However, an **MR murmur** would be heard loudest at the 5th left **intercostal space** in the **midclavicular line (mitral region)**, and increase in intensity with maneuvers that increase **afterload** (e.g., hand clenching). Moreover, **DMD** would cause rapidly progressive muscle weakness with an onset during childhood, and female individuals are very rarely affected by this X-linked recessive condition.

B - Fibrillin-1 defect

Image




Explanation Why

A [fibrillin-1](#) defect is the cause of [Marfan syndrome](#), which can result in [mitral valve prolapse](#) (MVP), [aortic aneurysms](#), and/or [aortic dissections](#). Severe MVP can cause exertional dyspnea and a [systolic murmur](#) that increases in intensity with maneuvers that decrease [preload](#) (e.g., standing up suddenly) and decreases in intensity with maneuvers that increase [afterload](#) (e.g., hand clenching). However, the [murmur](#) would be heard loudest at the 5th left [intercostal space](#) in the [midclavicular line](#) ([mitral region](#)) and would typically be preceded by a midsystolic click. Also, this patient does not have any [marfanoid features](#) (e.g., [arachnodactyly](#), tall stature, long extremities, hypermobile joints).

C - Myosin heavy chain defect

Image



Explanation But

All symptomatic patients should receive a [beta blocker](#) and be considered for an [implantable cardioverter defibrillator](#) (ICD) if there is a risk of sudden death. Positive [inotropic](#) and [afterload](#)-reducing or [preload](#)-reducing drugs (e.g., [digitalis](#), [glyceryl trinitrate](#), [calcium channel blockers](#) of the dihydropyridine class, ACE inhibitors) are contraindicated in cases of obstructive

[hypertrophic cardiomyopathy](#). If symptoms persist, myectomy or [transcoronary ablation of septal hypertrophy](#) are further treatment options.

Explanation Why

Mutations in the cardiac beta-[myosin heavy chain gene](#) cause [hypertrophic cardiomyopathy](#), which is the most common hereditary [cardiomyopathy](#). The mode of inheritance is typically [autosomal dominant](#). [Echocardiography](#) (or examination of an autopsy specimen) would most likely demonstrate concentric [left ventricular hypertrophy](#) or [hypertrophy](#) of the interventricular septum. Microscopic analysis will show [hypertrophic myocytes](#) with disoriented [myofibrils](#) (myofibrillar disarray).

D - GAA trinucleotide repeats

Explanation Why

GAA [trinucleotide repeats](#) within the [frataxin gene](#) on [chromosome 9](#) result in [Friedreich ataxia](#), which is associated with [hypertrophic cardiomyopathy](#). [Friedreich ataxia](#) would also manifest with impaired muscle coordination, deformities due to muscle [spasticity](#) (e.g., kyphoscoliosis, [pes cavus](#)), [hearing loss](#), visual impairment, and/or [diabetes mellitus](#). The absence of these features makes a GAA [trinucleotide repeat disorder](#) unlikely.

E - Myocyte ion channel defect

Explanation Why

[Myocyte](#) ion channel defects are responsible for several inherited long QT syndromes (e.g., [Romano-Ward syndrome](#), Jervell and Lange-Nielson syndrome), which manifest as acute episodes of [palpitations](#) and/or [syncope](#) (due to polymorphic [ventricular tachyarrhythmias](#)). The auscultatory findings in this patient suggest LVOT obstruction, which is not caused by [myocyte](#) ion channel defects.

F - CTG trinucleotide repeats

Explanation Why

CTG [trinucleotide repeats](#) within the [DMPK gene](#) result in [myotonic dystrophy](#). The typical cardiac manifestations in this condition are conduction blocks and/or [tachyarrhythmias](#). Also, [myotonic dystrophy](#) is associated with other features such as progressive muscle weakness, [myotonia](#), frontal balding, and/or visual impairment (due to [cataracts](#)), none of which are seen here.

G - Deletion in chromosome 7q

Explanation Why

A deletion in [chromosome 7q](#) occurs in [Williams syndrome](#), which is associated with [left ventricular outflow obstruction](#) due to [supravalvular aortic stenosis](#). Clinically, this can present with exertional dyspnea and a [systolic murmur](#) that decreases in intensity with maneuvers that increase [afterload](#) (e.g., fist clenching). However, a [murmur](#) due to [aortic stenosis](#) is best heard at the 2nd right [intercostal space](#) ([aortic area](#)), and decreases in intensity with maneuvers that decrease [preload](#) (e.g., standing up suddenly). Moreover, this patient does not have other findings indicative of [Williams syndrome](#), such as [elfin facies](#) and mental disabilities.

Question # 5

A 70-year-old man comes to the physician for evaluation of worsening nocturia, fatigue, and shortness of breath on exertion. While he used to be able to walk for 15 minutes at a time, he now has to pause every 5 minutes. Recently, he has started using two pillows to avoid waking up short of breath at night. He has a history of hypertension treated with daily amlodipine and prazosin; he has difficulty adhering to his medication regimen. His pulse is 75/min, and blood pressure is 150/90 mm Hg. Physical examination shows a laterally displaced apical heartbeat and 2+ bilateral pitting edema of the lower legs. Auscultation shows an S4 gallop and fine bibasilar rales. Further evaluation is most likely to show which of the following pathophysiologic changes in this patient?

	Answer	Image
A	Constriction of efferent renal arterioles	
B	Increase in urinary bicarbonate excretion	
C	Decrease in total peripheral vascular resistance	
D	Decrease in left ventricular elastance	
E	Reduction of alveolar surface tension	
F	Retention of potassium	

Hint

This patient has classic signs and symptoms of congestive heart failure (CHF), including dyspnea on exertion, paroxysmal nocturnal dyspnea, nocturia, bibasilar rales from pulmonary edema (suggestive of left-sided heart failure), and pitting edema (suggestive of right-sided heart failure). CHF is characterized by reduced cardiac output, which activates numerous compensatory neurohumoral mechanisms.

Correct Answer

A - Constriction of efferent renal arterioles

Explanation But

Although [RAAS](#) activation is a compensatory mechanism for maintaining [cardiac output](#) in patients with [CHF](#), prolonged renal hypoperfusion (particularly in conjunction with precipitating factors, such as [NSAID](#) use and [dehydration](#)) can lead to renal dysfunction (increased [BUN](#) and [creatinine](#)).

Explanation Why

The reduced [cardiac output](#) seen in [CHF](#) results in organ hypoperfusion, which also affects the [kidneys](#). Reduced [renal blood flow](#) leads to compensatory activation of [RAAS](#) in order to maintain the [GFR](#). [Angiotensin II](#) causes vasoconstriction of the afferent and [efferent arterioles](#) within the [kidney](#), decreasing the net [renal blood flow](#). Vasoconstriction is, however, more pronounced in the [efferent arteriole](#), which results in an increase in intraglomerular pressure to maintain [GFR](#). In combination with the other effects of [RAAS](#) activation ([aldosterone](#)-induced renal $\text{Na}^+/\text{H}_2\text{O}$ resorption and systemic vasoconstriction), this maintains [cardiac output](#).

B - Increase in urinary bicarbonate excretion

Explanation Why

An increase in [bicarbonate](#) excretion can be a compensatory mechanism in patients with [respiratory alkalosis](#) or a cause of [metabolic acidosis](#), e.g., in type 2 acute tubular [acidosis](#). This patient presents with [CHF](#), which would not result in increased [bicarbonate](#) excretion. Instead, elevated [aldosterone](#) levels due to upregulation of [RAAS](#) lead to an increase in renal sodium resorption in exchange for both hydrogen and potassium. This would cause a net loss of hydrogen in the [urine](#). For this reason, patients with [CHF](#) are [prone](#) to developing [metabolic alkalosis](#).

C - Decrease in total peripheral vascular resistance

Explanation Why

This patient presents with [CHF](#), which is characterized by reduced [cardiac output](#) and activation of the [sympathetic](#) system and [RAAS](#). This compensatory mechanism results in systemic vasoconstriction, which increases rather than decreases [systemic vascular resistance](#).

D - Decrease in left ventricular elastance

Explanation Why

Elastance ($E = \text{pressure}/\text{volume}$) is the reciprocal value of compliance ($C = 1/E = \text{volume}/\text{pressure}$), the more commonly used concept to evaluate the relationship between transported volume and applied pressure. An increase in [left ventricular](#) compliance is seen in patients with [dilated cardiomyopathy](#) ([eccentric hypertrophy](#)), which results in [systolic dysfunction](#). This patient, however, presents with [CHF](#) that is most likely due to chronic [hypertension](#), as suggested by his history of poor compliance with his medication regimen. Chronic [hypertension](#) results predominantly in concentric ventricular [hypertrophy](#), which leads to decreased [left ventricular](#) compliance (as indicated by the [S₄ gallop](#) seen here) and [diastolic dysfunction](#). This translates to an increased [left ventricular](#) elastance.

E - Reduction of alveolar surface tension

Explanation Why

[Surfactant](#) ensures low alveolar surface tension to prevent alveolar collapse. This patient has signs of [pulmonary edema](#), which is characterized by fluid accumulation within the pulmonary [interstitium](#) and alveoli. This, in turn, causes increased (not decreased) alveolar surface tension, which results in alveolar collapse that causes [dyspnea](#), decreased exercise capacity, and [paroxysmal nocturnal dyspnea](#), as seen in this patient.

F - Retention of potassium

Explanation Why

An increase in potassium retention occurs secondary to [hypoaldosteronism](#). However, this patient presents with [CHF](#), characterized by reduced [cardiac output](#) and organ hypoperfusion, which also affects the [kidneys](#). Reduced [renal blood flow](#) results in [RAAS](#) activation. This patient is therefore more likely to have elevated [aldosterone](#) levels, resulting in increased renal Na^+ and H_2O resorption as well as potassium excretion.

Question # 6

A 72-year-old woman comes to the physician because of a 1-month history of frequent headaches, double vision, fatigue, and myalgias. Physical examination shows pallor, tenderness to palpation over the temples, and inward deviation of the left eye. Laboratory studies show an erythrocyte sedimentation rate of 65 mm/h. Temporal artery biopsy shows transmural granulomatous inflammation and fragmentation of the internal elastic lamina. Treatment with high-dose intravenous corticosteroids is begun. The patient's condition puts her at greatest risk of developing which of the following complications?

	Answer	Image
A	Pulmonary hypertension	
B	Lower extremity gangrene	
C	Nasal septum perforation	
D	Palpable purpura	
E	Posterior uveitis	
F	Thoracic aortic aneurysm	Polymyalgia rheumatica (PMR) Giant cell arteritis (GCA; temporal arteritis) Etiology Autoimmune, granulomatous inflammation of medium and large arteries (GCA esp. affects the carotid arteries). A common pathophysiology is presumed. Epidemiology ♀ > ♂; incidence rises with age Antibody serology No autoantibodies Laboratory analysis ↑↑ ESR, ↑ CRP Complications GCA: 20–30% risk of blindness without therapy Note Distal joints are not involved! GCA: 50% associated with PMR General symptoms 8 symptoms - Depressive mood - Weight loss PMR - Symmetrical pain (esp. at night) and morning stiffness - Synovitis and carpal tunnel syndrome - Nonerosive arthritis of medium-sized joints GCA - Pain in the temporal region (hence "temporal arteritis"); thickened superficial temporal artery - Ocular involvement: vision disorders, e.g., amaurosis fugax, AION - Jaw claudication - Aortic involvement

Hint

A headache, temporal tenderness, double vision, fatigue, and an increased ESR in this woman over 50 years of age are highly suggestive of giant cell arteritis (GCA). Concurrent myalgia possibly indicates polymyalgia rheumatica, which is commonly associated with GCA. Focal granulomatous inflammation of the temporal artery layers with fragmentation of the internal elastic lamina confirms the diagnosis of GCA in this patient.

Correct Answer

A - Pulmonary hypertension

Explanation Why

[Pulmonary hypertension](#) may develop in [diffuse systemic sclerosis](#), a condition that usually presents with [Raynaud phenomenon](#), cutaneous findings such as [sclerodactyly](#), and inner organ involvement (e.g., dysmotility of the [gastrointestinal tract](#), cardiac disease, renal disease). Because [GCA](#) involves medium to large-sized blood vessels, the pulmonary vessels are not typically involved.

B - Lower extremity gangrene

Explanation Why

Lower extremity [gangrene](#) is often seen as a late complication of [thromboangiitis obliterans](#), also known as [Buerger disease](#). As the disease progresses, patients develop [critical limb ischemia](#) and [pain](#) at rest, absent pulses, and digital [gangrene](#). This type of [dry gangrene](#) (involving the small and medium-sized blood vessels) is not seen in [GCA](#), which typically only affects larger blood vessels.

C - Nasal septum perforation

Explanation Why

[Nasal septum perforation](#) may be seen in [granulomatosis with polyangiitis](#), which typically presents with constitutional symptoms and localized manifestations, such as [chronic sinusitis](#), [rhinitis](#), [otitis media](#), ocular conditions, and/or [skin](#) lesions. [GCA](#) affects medium to large-sized vessels and therefore does not usually lead to [nasal septum perforation](#).

D - Palpable purpura

Explanation Why

Palpable [purpura](#) is commonly seen in small vessel [vasculitides](#) including [granulomatosis with polyangiitis](#) ([Wegener disease](#)), [eosinophilic granulomatosis with polyangiitis](#) ([Churg-Strauss syndrome](#)), Henoch-Schönlein [purpura](#), and [polyarteritis nodosa](#). Because [GCA](#) involves medium to large-sized blood vessels, it does not result in cutaneous [purpura](#).


E - Posterior uveitis

Explanation Why

Inflammatory conditions associated with the development of [posterior uveitis](#) include [Behcet disease](#), [sarcoidosis](#), spondyloarthritis, and [IBD](#). Although [eye](#) involvement can occur in patients with [GCA](#) (e.g., [anterior ischemic optic neuropathy](#) or [central retinal artery occlusion](#)) and blindness is a possible complication of [GCA](#), [posterior uveitis](#) is not associated with this condition.

F - Thoracic aortic aneurysm

Image



Explanation But

There is currently no general recommendation to screen patients with [GCA](#) for large vessel disease, but some medical centers perform screening with serial chest [x-rays](#) or CT/MR angiography for up to 10 years after the diagnosis of [GCA](#).


Explanation Why

[Giant cell arteritis](#) ([GCA](#)) causes chronic [inflammation](#) of large and medium-sized [arteries](#), in particular the carotid [arteries](#), its major branches, and the aorta. Approximately 10–20% of patients with [giant cell arteritis](#) ([GCA](#)) will have involvement of the [thoracic aorta](#), which significantly increases the risk of thoracic [artery aneurysms](#) and dissections. The [median](#) time between the diagnosis of [GCA](#) and [thoracic aortic aneurysm](#) is around 10 years.

Question # 7

A 3-year-old girl is brought to the cardiologist because of sweating and respiratory distress while eating. She is at the 30th percentile for height and 15th percentile for weight. Echocardiography shows a defect in the membranous portion of the interventricular septum and a moderately decreased left ventricular ejection fraction. Physical examination is most likely to show which of the following findings?

	Answer	Image
A	Systolic murmur that increases with hand clenching	The image displays two waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The background is divided into alternating Diastole (light blue) and Systole (light orange) phases. The ECG shows a normal QRS complex. The PCG shows a normal S1 and S2, with a systolic murmur occurring between them. The murmur is labeled as increasing with hand clenching, which is characteristic of aortic stenosis.
B	Wide, fixed splitting of the second heart sound	
C	Systolic murmur that increases with forced exhalation against a closed glottis	
D	Diastolic murmur preceded by opening snap	

	Answer	Image
E	Continuous murmur that is loudest at the second heart sound	 The image displays two synchronized cardiac waveforms. The top waveform is an Electrocardiogram (ECG) in blue, showing a regular rhythm with labeled phases: Diastole, Systole, Diastole, Systole, and Diastole. The bottom waveform is a Phonocardiogram (PCG) in orange, showing heart sounds corresponding to the ECG. The first heart sound (S1) is marked at the start of the first systole, and the second heart sound (S2) is marked at the start of the second systole. A continuous murmur is visible as a high-frequency, continuous oscillation that spans across the diastolic and systolic periods, with its amplitude peaking during the second heart sound (S2).


Hint

Examination findings for a symptomatic ventricular septal defect (VSD) in a patient this age will reflect shunting of blood from the left to right ventricle.

Correct Answer

A - Systolic murmur that increases with hand clenching

Image



Explanation Why

Hand clenching increases [systemic vascular resistance](#) and [afterload](#), which increases the left-to-right shunting of blood and results in increased volume of the [systolic murmur](#) heard in [VSD](#). Small [VSDs](#) may cause a loud [systolic murmur](#) with palpable thrill due to the large pressure differential between the left and right ventricles, while large [VSDs](#) are typically less obvious on auscultation. If left untreated, [VSDs](#) can lead to [pulmonary hypertension](#) and [heart failure](#).

B - Wide, fixed splitting of the second heart sound

Explanation Why

A wide, fixed (i.e., does not change with respiration) [splitting of the second heart sound](#) is characteristic of [atrial septal defect](#) (ASD). It occurs due to [right ventricular](#) overload and subsequent delay in [pulmonary valve](#) closure. Fixed [splitting of the second heart sound](#) is not a common feature of [VSD](#).

C -

Systolic murmur that increases with forced exhalation against a closed glottis

Explanation Why

A [systolic murmur](#) that increases with forced exhalation against a closed [glottis](#) ([Valsalva maneuver](#)) is characteristic of [hypertrophic obstructive cardiomyopathy](#) (HOCM). The [Valsalva maneuver](#) reduces both [preload](#) and [left ventricular afterload](#), which would exacerbate the subaortic stenosis in [HOCM](#) but reduce the [left-to-right shunt](#) of blood in [VSD](#). The [Valsalva maneuver](#) would therefore decrease this patient's [murmur](#) intensity.


D - Diastolic murmur preceded by opening snap

Explanation Why

A [diastolic murmur](#) preceded by an [opening snap](#) at the apex is characteristic of [mitral stenosis](#) (MS). The [opening snap](#) occurs when leaflet motion suddenly stops during [diastole](#) after the stenosed valve has opened fully. While a short, mid-[diastolic murmur](#) can sometimes be heard in [VSD](#) due to increased blood flow across the [mitral valve](#), an [opening snap](#) would not be expected.

E - Continuous murmur that is loudest at the second heart sound

Image



Explanation Why

A [continuous murmur](#) that becomes louder at the [second heart sound](#) is characteristic of [patent ductus arteriosus \(PDA\)](#). The [murmur](#) is best appreciated in the left infraclavicular region and results from the constant flow of blood from the high pressure aorta to the low pressure [pulmonary artery](#). [VSD](#) is characterized by a different finding on auscultation.

Question # 8

A 68-year-old woman with hypertension and congestive heart failure is brought to the emergency department because of a 2-day history of shortness of breath and confusion. Her pulse is 112/min, respirations are 22/min, and blood pressure is 73/45 mm Hg. Examination shows cool extremities, jugular venous distention, and diffuse crackles in the lungs. Treatment with intravenous dobutamine is begun. Which of the following is the primary mechanism of action of this drug?

	Answer	Image
A	α 2-adrenergic agonism	
B	β 2-adrenergic antagonism	
C	β 1-adrenergic agonism	
D	V1 antagonism	
E	α 1-adrenergic antagonism	
F	D1 agonism	

Hint

Dobutamine is used in patients with cardiogenic shock to increase cardiac output via an increase in heart rate as well as contractility.

Correct Answer

A - α 2-adrenergic agonism

Explanation Why

α -2 adrenergic agonists (e.g., [clonidine](#)) would cause a decrease in [sympathetic](#) tone and further exacerbate this patient's [cardiogenic shock](#). [Dobutamine](#) may have a mild effect on [\$\alpha\$ -1 receptors](#) but does not have a clinically meaningful effect on [\$\alpha\$ -2 receptors](#).

B - β 2-adrenergic antagonism

Explanation Why

[\$\beta\$ -2 receptor](#) antagonism (e.g., due to [propranolol](#)) would cause bronchial constriction, vasoconstriction, and decreased [cardiac output](#), none of which are desired effects in this patient with [cardiogenic shock](#). [Dobutamine](#), the medication this patient received, has a weak agonistic effect on [\$\beta\$ -2 receptors](#).

C - β 1-adrenergic agonism

Explanation Why

[Dobutamine](#) is used to increase cardiac contractility and [cardiac output](#) in patients with [cardiogenic shock](#) primarily via its agonistic effects on [\$\beta\$ -1 adrenergic receptors](#). At higher doses, it also has weak [agonism](#) at both β -2 and [\$\alpha\$ -1 adrenergic receptors](#). Major adverse effects include [hypertension](#) and increased risk of [arrhythmia](#).

D - V1 antagonism

Explanation Why

[V1 receptor](#) antagonists (e.g., VAPTANS) cause vascular [smooth muscle](#) relaxation, producing an antihypertensive effect (which would worsen this patient's [hypotension](#)). The [dobutamine](#) this patient received does not possess V1 antagonistic properties.

E - α 1-adrenergic antagonism

Explanation Why

[\$\alpha\$ -1 adrenergic receptor](#) antagonists (e.g., [prazosin](#), [phenoxybenzamine](#)) cause [vasodilation](#), which would further exacerbate the [hypotension](#) seen in this patient. The [dobutamine](#) administered here has no α 1 antagonistic properties; at high doses, it exerts a weak agonistic effect on [\$\alpha\$ -1 receptors](#).


F - D1 agonism

Explanation Why

[D1 receptor agonists](#) (e.g., [fenoldopam](#) and [dopamine](#)) cause vascular [smooth muscle](#) relaxation and would worsen this patient's [hypotension](#). [Dobutamine](#), which this patient received, does not possess D1 agonistic properties.

Question # 9

A 69-year-old man is brought to the emergency department because of severe epigastric pain and vomiting that started 30 minutes ago while gardening. His pulse is 55/min, respirations are 30/min, and blood pressure is 90/50 mm Hg. Physical examination shows diaphoresis and jugular venous distention. Crackles are heard in both lower lung fields. An ECG shows P waves independent of QRS complexes and ST segment elevation in leads II, III, and aVF. Coronary angiography is most likely to show narrowing of which of the following vessels?

	Answer	Image
A	Left coronary artery	
B	Proximal right coronary artery	 Anterior surface of the heart (ventral view) Diaphragmatic surface of the heart (dorsocaudal view) Regions supplied by: - RCA - AMA - LAD - LCX - PDA
C	Left circumflex artery	
D	Left anterior descending artery	
E	Posterior interventricular artery	

Hint

The symptoms of epigastric pain and vomiting, combined with ST segment elevation in leads II, III, and aVF and complete heart block, are indicative of an inferior wall myocardial infarction.

Correct Answer


A - Left coronary artery

Explanation Why

The [left coronary artery \(LCA\)](#) branches to the [left anterior descending artery \(LAD\)](#) and the [left circumflex artery \(LCX\)](#), so occlusion of the [LCA](#) can lead to a massive [myocardial infarction](#) involving the a combination of the [anterior](#) (ECG changes in I, aVL, V1–6), anteroseptal (I, aVL, V1–4), anterolateral (I, aVL, V5–6/7), apical (I, aVL, V3–5), and posterolateral (II, III, aVF, V5–7, and sometimes I, and aVL) walls. In a patient with left-dominant coronary circulation, [LCA](#) occlusion leading to [distal](#) occlusion of the [posterior interventricular artery \(PIV\)](#) could also cause a [posterior](#) wall [myocardial infarction](#).

B - Proximal right coronary artery

Image



Explanation Why

Occlusion of the [proximal right coronary artery \(RCA\)](#) can cause an inferior wall [myocardial infarction](#), which in turn can present atypically with epigastric [pain](#). High-degree [AV block](#) can occur in this setting because the [SA node](#) and [AV node](#) are usually supplied by the [RCA](#) (the [SA node](#) directly, and the [AV node](#) via the [AV nodal](#) branch). Right-sided leads (V4R, V5R, and V6R) should also be obtained to evaluate for [right ventricular \(RV\) infarction](#) (especially given the patient's [jugular venous distention](#)), as the [RCA](#) supplies the RV via its right marginal branch.

C - Left circumflex artery

Explanation Why

The [left circumflex artery \(LCX\)](#) supplies the [lateral](#) and [posterior](#) walls of the [left ventricle](#). Occlusion of the [LCX](#) causes [ECG](#) changes in the [lateral](#) (V5, V6, I, aVL) leads. In patients with left-dominant coronary circulation, occlusion of the [LCX](#) causes [distal infarction](#) of the [posterior descending artery](#), leading to an inferior wall [myocardial infarction](#) and [ECG](#) changes in the [posterior/posterolateral](#) (V7–V9) and inferior (II, III, aVF) leads. However, the [LCX](#) is not the most likely source of this patient's inferior wall [myocardial infarction](#) because only approx. 10% of the population is left-dominant.

D - Left anterior descending artery

Explanation Why

Occlusion of the [left anterior descending artery \(LAD\)](#) would cause a [myocardial infarction](#) of the [anterior](#) ([ECG](#) changes in I, aVL, V1–6), anteroseptal (I, aVL, V1–4), anterolateral (I, aVL, V5–6/7), and/or apical wall (I, aVL, V3–5). An anteroseptal [myocardial infarction](#) can sometimes cause serious [arrhythmias](#), such as high-degree [AV block](#), since the LAD supplies the left and right [bundle branches](#) and partially supplies the left [anterior](#) and [posterior](#) fascicles. However, conduction abnormalities are more common following an inferior [myocardial infarction](#), as is epigastric [pain](#).

E - Posterior interventricular artery

Explanation Why

Occlusion of the [posterior interventricular artery \(PIV\)](#) would cause a [posterior](#) wall [myocardial infarction](#), which would manifest as ST-segment depression in leads V1 and V2, sometimes with prominent R waves. [Posterior](#) leads (V7–9) would show ST-segment elevation. Epigastric [pain](#) is not common in [posterior](#) wall [myocardial infarctions](#), nor are conduction abnormalities, since the [PIV](#) does not supply the [conduction system](#).

Question # 10

A 25-year-old man comes to the emergency department because of a 1-week-history of progressively worsening dyspnea and intermittent chest pain that increases on inspiration. He had an upper respiratory tract infection 2 weeks ago. His pulse is 115/min and blood pressure is 100/65 mm Hg. Examination shows inspiratory crackles bilaterally. His serum troponin I concentration is 0.21 ng/mL. An x-ray of the chest shows an enlarged cardiac silhouette and prominent vascular markings in both lung fields; costophrenic angles are blunted. ECG shows nonspecific T wave changes. Which of the following additional findings is most likely in this patient's condition?

	Answer	Image
A	Sarcomere duplication	
B	Elevated brain natriuretic peptide	
C	Opening snap with low-pitched diastolic rumble	
D	Electrical alternans	
E	Myocardial perfusion defect	

Hint

Subacute onset of dyspnea, chest pain, and inspiratory crackles, as well as an enlarged cardiac silhouette and prominent vascular markings, are consistent with congestive heart failure (CHF).

Correct Answer

A - Sarcomere duplication

Explanation Why

[Sarcomere](#) duplication is found in concentric [left ventricular hypertrophy \(LVH\)](#), which results from the LV chronically pumping against high [afterload](#) (e.g., [hypertension](#), [aortic stenosis](#)). Untreated [LVH](#) can lead to an enlarged [left ventricle](#) and eventually [congestive heart failure](#). However, this patient has no history or evidence of [hypertension](#) or [aortic stenosis](#). In addition, [ECG](#) findings typically show signs of [left ventricular](#) enlargement (e.g., [left axis deviation](#), widened QRS, increased R wave amplitude), none of which are seen here. Moreover, [LVH](#) would not explain this patient's elevated [cardiac biomarkers](#).

B - Elevated brain natriuretic peptide

Explanation But

Other complications of [myocarditis](#) include concurrent [pericarditis](#) ([perimyocarditis](#)) that may lead to [cardiac tamponade](#), [arrhythmias](#), [AV block](#), and [sudden cardiac death](#).

Explanation Why

[Brain natriuretic peptide \(BNP\)](#) is a very sensitive marker for [congestive heart failure](#), as [BNP](#) is secreted by a stretched or overworked cardiac ventricle. In this patient, [heart failure](#) is most likely secondary to viral [myocarditis](#), which damages [cardiomyocytes](#) by virus-mediated lysis and immunological [molecular mimicry](#) and [cross-reactivity](#). Commonly implicated viruses include [coxsackie B](#), [adenovirus](#), [parvovirus B19](#), [HIV](#), and [HHV-6](#). This patient's recent history of upper respiratory tract infection, elevated [troponin](#) levels, and nonspecific [T-wave](#) changes are features of [myocarditis](#). [Myocarditis](#) has a broad range of clinical presentations, from subclinical (and often undiagnosed) to [acute heart failure](#), as seen in this patient.

C - Opening snap with low-pitched diastolic rumble

Explanation Why

An [opening snap](#) with low-pitched [diastolic](#) rumble is the classic finding in [mitral stenosis](#) (MS), which, if prolonged, can cause [left atrial enlargement](#) ([LAE](#)) and [left heart failure](#). While this patient has evidence of [left heart failure](#) ([pulmonary edema](#), [dyspnea](#)), neither MS nor [LAE](#) would explain his enlarged [cardiac silhouette](#). In addition, [P mitrale](#) on [ECG](#) would be an expected characteristic finding of [LAE](#). Moreover, elevated [troponin](#) indicates [myocardial injury](#), which is not caused by [mitral stenosis](#).

D - Electrical alternans

Explanation Why

[Electrical alternans](#) on [ECG](#) are seen in [pericardial effusion](#) due to a swinging beating [heart](#) surrounded by [pericardial](#) fluid, which can also be seen on [echocardiography](#). This patient's recent viral illness increases his risk of viral [pericarditis](#), which can cause [pleuritic chest pain](#), [dyspnea](#), and [pericardial effusion](#). Although a pericardial effusion would also cause an enlarged [cardiac silhouette](#) ([water bottle heart](#)), it would not cause [pulmonary edema](#). Moreover, [pericarditis](#) classically causes diffuse [ST elevations](#).



E - Myocardial perfusion defect

Explanation Why

A [myocardial](#) perfusion defect is seen in patients with [acute coronary syndrome](#) ([ACS](#)), which can also present with [chest pain](#), [dyspnea](#), and elevated [troponin](#) levels. However, this patient lacks other typical features of [ACS](#) including nausea, vomiting, [diaphoresis](#), and [palpitations](#). In addition, the [pain](#) associated with [ACS](#) does not typically worsen upon inspiration. Moreover, this patient is only 25-years-old and unlikely to have clinically significant [coronary artery disease](#).

Question # 11

A 74-year-old man comes to the physician for a 6-month history of progressively worsening fatigue and shortness of breath on exertion. He immigrated to the United States 35 years ago from India. His pulse is 89/min and blood pressure is 145/60 mm Hg. Crackles are heard at the lung bases. Cardiac examination shows a grade 3/6 early diastolic murmur loudest at the left sternal border in the third intercostal space. Further evaluation of this patient is most likely to show which of the following?

	Answer	Image
A	Pulsus parvus et tardus	 The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The background is divided into alternating light blue (Diastole) and light orange (Systole) vertical bands. The ECG shows a normal sinus rhythm. The PCG shows aortic regurgitation (A2) and aortic stenosis (A1). The A2 is significantly louder than the A1, indicating aortic regurgitation. The A1 is significantly softer than the A2, indicating aortic stenosis. This combination of findings is characteristic of aortic regurgitation with aortic stenosis, which can lead to aortic aneurysm. The PCG also shows aortic stenosis (A1) and aortic regurgitation (A2). The A2 is significantly louder than the A1, indicating aortic regurgitation. The A1 is significantly softer than the A2, indicating aortic stenosis. This combination of findings is characteristic of aortic regurgitation with aortic stenosis, which can lead to aortic aneurysm.
B	Pulsus paradoxus	
C	Fixed splitting of S2	
D	Water hammer pulse	 The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The background is divided into alternating light blue (Diastole) and light orange (Systole) vertical bands. The ECG shows a normal sinus rhythm. The PCG shows aortic regurgitation (A2) and aortic stenosis (A1). The A2 is significantly louder than the A1, indicating aortic regurgitation. The A1 is significantly softer than the A2, indicating aortic stenosis. This combination of findings is characteristic of aortic regurgitation with aortic stenosis, which can lead to aortic aneurysm. The PCG also shows aortic stenosis (A1) and aortic regurgitation (A2). The A2 is significantly louder than the A1, indicating aortic regurgitation. The A1 is significantly softer than the A2, indicating aortic stenosis. This combination of findings is characteristic of aortic regurgitation with aortic stenosis, which can lead to aortic aneurysm.

	Answer	Image
E	Paradoxical splitting of S ₂	1st heart sound (S1) 2nd heart sound (S2) Expiration P2 A2 Inspiration P2 A2

Hint

This patient presents with symptoms of left heart failure, a high pulse pressure, and an early diastolic murmur at Erb's point (left sternal border, third intercostal space), which suggests chronic aortic regurgitation (AR). Based on his history of emigration from India, this patient likely has rheumatic heart disease, the most common cause of AR in developing countries.

Correct Answer

A - Pulsus parvus et tardus

Image



Explanation Why

[Pulsus parvus et tardus](#) is seen in patients with [aortic stenosis](#), which can present with [angina](#), [syncope](#), and [heart failure](#). [Aortic stenosis](#) is characterized by a harsh crescendo-decrescendo late [systolic ejection murmur](#), unlike the [diastolic murmur](#) heard in this patient. [Pulsus parvus et tardus](#) is not a typical feature of [aortic regurgitation](#).

B - Pulsus paradoxus

Explanation Why

[Pulsus paradoxus](#) is characterized by decrease in [systolic](#) blood pressure during inspiration and is

classically seen in patients with [cardiac tamponade](#) and/or [constrictive pericarditis](#). [Pulsus paradoxus](#) is not a feature of [aortic regurgitation](#).


C - Fixed splitting of S₂

Explanation Why

Fixed [splitting of S₂](#) most commonly occurs as a result of pathologic processes that increase [right ventricular](#) volume (thereby delaying pulmonic valve closure), such as an [atrial septal defect](#). Fixed [splitting of S₂](#) is not characteristic of [aortic regurgitation](#).

D - Water hammer pulse

Image




Explanation Why

A [water hammer pulse](#) is a classic finding on [cardiovascular examination](#) in patients with significant

[aortic regurgitation](#), and results from the rapid ejection of a large [stroke volume](#) into a low-pressure arterial system. Other findings in patients with [aortic regurgitation](#) may include [pulse-synchronous head nodding](#) ([de Musset sign](#)) and [Austin Flint murmur](#).

E - Paradoxical splitting of S₂

Image



Explanation Why

Paradoxical [splitting of S₂](#) refers to the pathological closure of the [pulmonary valve](#) before the [aortic valve](#), resulting in an S₂ split with expiration (instead of the physiologic [splitting of S₂](#) heard normally with inspiration). [Paradoxical splitting](#) is caused by conditions that delay [left ventricular outflow](#), such as [aortic stenosis](#) or [left bundle branch block](#). It is not typically heard in [aortic regurgitation](#).

Question # 12

A 33-year-old man is evaluated by paramedics after being found unconscious outside of his home. He has no palpable pulses. Physical examination shows erythematous marks in a fern-leaf pattern on his lower extremities. An ECG shows ventricular fibrillation. Which of the following is the most likely cause of this patient's findings?

	Answer	Image
A	Lightning strike	
B	Cholesterol emboli	
C	Hypothermia	
D	Opioid overdose	

	Answer	Image
E	Infective endocarditis	

Hint

The fern-leaf pattern on the patient's lower extremities is caused by local hyperemia.

Correct Answer

A - Lightning strike

Image



Explanation Why

The fern-leaf pattern seen on this patient's lower extremities are [Lichtenberg figures](#), which are pathognomonic of a [lightning strike](#). In about 30% of cases, a [lightning strike](#) is fatal; the most common cause of death is [cardiac arrest](#) due to a fatal [arrhythmia](#), such as [ventricular fibrillation](#) or [asystole](#). [Superficial burns](#) and metallization of the [skin](#) may also occur after a [lightning strike](#) injury.

B - Cholesterol emboli

Explanation Why

[Cholesterol emboli](#), as well as other vascular diseases that cause slower blood flow due to

obstruction of blood vessels (e.g., [vasculitis](#), [atherosclerosis](#)), can lead to [livedo reticularis](#). However, [cholesterol emboli](#) are unlikely to lead to [ventricular fibrillation](#) and have a different appearance than the fern-leaf pattern seen here.

C - Hypothermia

Image



Explanation Why

[Severe hypothermia](#) (< 82°F) can lead to pulselessness or [ventricular fibrillation](#). However, the usual [skin](#) finding with [hypothermia](#) is [livor mortis](#), rather than the fern-leaf pattern seen here.

D - Opioid overdose

Explanation Why

[Opioid overdose](#) should always be considered when a person is found unconscious without a known

[history of present illness](#) or [past medical history](#). Although [opioid overdose](#) can result in cardiac abnormalities, the associated [skin](#) findings do not appear in a fern-leaf pattern.

E - Infective endocarditis

Image









Explanation Why

[Infective endocarditis](#) can cause [arrhythmias](#) if a perivalvular abscess forms. [Skin](#) findings associated with [infective endocarditis](#) include [Janeway lesions](#), [splinter hemorrhages](#), and [Osler nodes](#). However, these [skin](#) findings do not have the fern-leaf pattern seen in this patient. Furthermore, severe constitutional symptoms (e.g., high [fever](#), malaise) and progression over at least days would be expected with [infective endocarditis](#).

Question # 13

A 16-year-old boy is brought to the physician by his parents because of a 6-month history of progressive fatigue and worsening shortness of breath on exertion. The parents report that the boy “has always been tired out a bit more easily than other kids.” The family recently immigrated to the United States from rural South Korea. Pulse oximetry on room air shows an oxygen saturation of 96% on bilateral index fingers. Extremity pulses are synchronous bilaterally; radial pulses are bounding. There is mild bluish discoloration and bulbous enlargement of the distal toes bilaterally. Echocardiography is most likely to show which of the following?

	Answer	Image
A	Single overriding great vessel arising from the heart	
B	Anomalous shunting of blood through a defect in atrial septum	1  2  3 
C	Persistent blood flow between the pulmonary artery and descending aorta	1  2  3 

	Answer	Image
D	Positioning of the ascending aorta directly over a ventricular septal defect	
E	Abnormal narrowing of the aorta at the aortic isthmus	

Hint

The patient presents with symptoms of heart failure, lower extremity cyanosis, and clubbing with preserved upper extremity oxygenation – a finding known as differential cyanosis. Further evaluation of the patient will likely reveal a loud, continuous, machine-like murmur best heard in the left infraclavicular area.

Correct Answer

A - Single overriding great vessel arising from the heart


Explanation Why

A single overriding great vessel is called a [persistent truncus arteriosus](#). This single arterial trunk arising from the [heart](#) provides mixed blood to the [pulmonary arteries](#) and the [systemic circulation](#). Pulses may be accentuated and bounding; however, [oxygen saturation](#) would be decreased in all extremities. Furthermore, if it is not diagnosed prenatally, it will typically present perinatally with symptoms of global [cyanosis](#) and [heart failure](#), while this [adolescent](#) patient only has [differential cyanosis](#).


B - Anomalous shunting of blood through a defect in atrial septum

Image


1



2



3




Explanation Why

[Atrial septal defects](#) (ASD) are often asymptomatic. In adults, larger defects may lead to a distinct [left-to-right shunt](#), which can result in [right heart failure](#), [supraventricular arrhythmias](#) and [pulmonary hypertension](#), and ultimately lead to an [Eisenmenger reaction](#) and global [cyanosis](#). [Differential cyanosis](#) would not be seen.

C - Persistent blood flow between the pulmonary artery and descending aorta

Image



Explanation But

[Infants](#) with [PDA](#) usually receive an infusion of [indomethacin](#) to induce closure of the duct, while in older children symptomatic PDAs are typically closed surgically or via percutaneous catheter occlusion.


Explanation Why

Failure of the [ductus arteriosus](#) to completely close postnatally allows blood to flow from the [descending aorta](#) back to the [pulmonary artery](#) (left-to-right shunt). In a large [patent ductus arteriosus](#)

(PDA), this leads to volume overload of the [right ventricle](#) and the [lung](#) circuit. This increased blood flow causes [pulmonary hypertension](#) and irreversible remodeling of small pulmonary vessels, which may eventually lead to an [Eisenmenger reaction](#), manifesting as [cyanosis](#) and [clubbing](#). The symptoms of [differential cyanosis](#) in this patient are specific for [PDA](#). Because the ductus is located on the [descending aorta](#) after the origin of the [left subclavian artery](#), reversal of the shunt results in a mixture of deoxygenated and oxygenated blood reaching only the lower body.

D - Positioning of the ascending aorta directly over a ventricular septal defect

Image




Explanation Why

Positioning of the [ascending aorta](#) directly over a [ventricular septal defect](#) is seen in [tetralogy of Fallot](#). The other features of this cardiac abnormality are [pulmonary stenosis](#) and [right ventricular hypertrophy](#). It is typically symptomatic in [infancy](#) and presents with [tet spells](#) and mild [cyanosis](#) in all extremities.

E - Abnormal narrowing of the aorta at the aortic isthmus

Image



Explanation Why

Abnormal narrowing of the aorta at the [aortic isthmus](#) is consistent with [coarctation of the aorta](#), which may manifest with [bounding pulses](#) and [cyanosis](#) of the lower extremities in [neonates](#). Older children will have cool lower extremities with faint or unpalpable pulses, but no [cyanosis](#). Moreover, [physical examination](#) would demonstrate a brachio-femoral [pulse](#) delay, while this patient has synchronous pulses. Finally, additional symptoms, such as [headaches](#), [epistaxis](#), and [claudication](#) of the lower extremities, would be expected in a patient presenting with [coarctation of the aorta](#).

Question # 14

A 55-year-old man is brought to the emergency department after collapsing at the grocery store and losing consciousness for 1 minute following a sudden onset of palpitations. He has no chest pain or dizziness. He has a history of hypertension, type 2 diabetes mellitus, and is currently being treated as an outpatient for pneumonia. His current medications include lisinopril, metformin, and azithromycin. He has no sign of injury, and he appears well. During the examination, he loses consciousness again. Telemetry shows polymorphic ventricular tachycardia with cyclic alteration of the QRS axis, which spontaneously resolves after 30 seconds. Which of the following is the most likely underlying cause of this patient's syncope?

	Answer	Image
A	Hypomagnesemia	
B	Brugada syndrome	
C	Prolonged QT interval	
D	Accessory conduction pathway	
E	Hyperkalemia	

Hint

Polymorphic ventricular tachycardia with cyclic alteration of the QRS axis (sinusoidal waveform) is referred to as torsade de pointes and can progress to life-threatening ventricular fibrillation. In this patient, the torsade de pointes is probably due to a side effect of the macrolide antibiotic, azithromycin.

Correct Answer

A - Hypomagnesemia

Explanation Why

[Hypomagnesemia](#) is a predisposing factor for [torsade de pointes](#). However, this patient has no [risk factors](#) for [hypomagnesemia](#) (e.g., [diarrhea](#), [alcoholism](#), use of [diuretics](#)). He also lacks other typical features of [hypomagnesemia](#) such as neuromuscular manifestations (e.g., [tetany](#), [seizures](#)).

B - Brugada syndrome

Explanation Why

[Brugada syndrome](#) can present with [syncope](#) and [ventricular tachyarrhythmia](#), as seen in this patient. However, the characteristic [ECG](#) findings include incomplete [right bundle branch block](#) and ST-segment elevations in the [anterior precordial leads](#), which are not seen on this patient's [ECG](#).

C - Prolonged QT interval

Explanation Why

A [prolonged QT interval](#) may be congenital or acquired, and typically remains asymptomatic until patients present with [syncope](#), or life-threatening [arrhythmias](#) such as [torsade de pointes](#). Acquired [prolongation of the QT interval](#) can be caused by certain medications such as [antibiotics](#) ([macrolides](#), [fluoroquinolones](#)), [antiarrhythmics](#) ([dofetilide](#)), or [tricyclic antidepressants](#). This patient's history of [azithromycin](#) therapy suggests that drug-induced [QT prolongation](#) is the most likely cause of this patient's [torsade de pointes](#) and resulting [syncope](#).

D - Accessory conduction pathway

Explanation Why

A fast accessory conduction pathway can be seen in [Wolff-Parkinson-White \(WPW\)](#) syndrome, which can present with [palpitation](#) and [syncope](#), as seen in this patient. However, typical [ECG](#) findings include a shortened [PR interval](#) (< 120 ms), slurring and slow rise of the initial upstroke of the [QRS complex](#) (delta wave), and a widened [QRS complex](#), which are absent from this patient's [ECG](#). [WPW syndrome](#) causes [atrioventricular reentrant tachycardia](#) but not [ventricular tachycardias](#) such as [torsade de pointes](#).

E - Hyperkalemia

Explanation Why

While [hyperkalemia](#) can result in a widening of the [QRS complex](#) and a sinusoidal shape, it does not cause [torsade de pointes](#). [Hyperkalemia](#) classically results in peaked [T waves](#), followed by the development of disappearing [P waves](#), a prolonged QRS interval, and eventually, a sine wave appearance that can lead to [ventricular fibrillation](#) and death. Additionally, this patient lacks any etiology or additional symptoms of [hyperkalemia](#), including malaise, [palpitations](#), and muscle weakness.

Question # 15

A 70-year-old man comes to the emergency department after briefly losing consciousness and collapsing when rising from a chair. He did not sustain any injuries from his collapse. He has had a two-week history of dizziness upon standing. He has smoked one pack of cigarettes daily for 55 years. He drinks three beers and two glasses of whiskey daily. He currently takes dutasteride and tamsulosin for benign prostatic hyperplasia. His blood pressure is 120/80 mm Hg supine and 100/70 mm Hg one minute after standing with no change in pulse rate. Physical examination shows conjunctival pallor. On cardiac auscultation, there is a plopping sound followed by a low-pitched, rumbling mid-diastolic murmur heard best at the apex. Which of the following is the most likely cause of this patient's presentation?

	Answer	Image
A	Stokes-Adams attack	
B	Constricted aortic valve orifice	
C	Cardiac tumor	
D	Adverse medication effect	
E	Left ventricular dilation	

Hint

This patient presents with syncope, orthostatic hypotension (drop of ≥ 20 mm Hg in systolic pressure from supine to standing), and a diastolic murmur over the apex, all of which are suspicious for an obstruction of the blood flow into the left ventricle.

Correct Answer

A - Stokes-Adams attack

Explanation Why

[Stokes-Adams attacks](#) are characterized by a sudden loss of consciousness, usually without warning and lasting for a few seconds as in this patient. However, these episodes of fainting are due to an abnormal [heart](#) rhythm (especially [complete atrioventricular block](#)) and not due to a mechanical obstruction of blood flow. This patient's normal rhythm strip suggests a different etiology of his [syncope](#).

B - Constricted aortic valve orifice

Explanation Why

[Aortic valve stenosis](#) is the most common [valvular heart disease](#) and may also present with [dizziness](#) and [syncope](#). It should always be considered as a differential diagnosis in an elderly patient with a [heart murmur](#), such as this one. The auscultatory finding of [aortic valve stenosis](#) is a harsh crescendo-decrescendo late [systolic ejection murmur](#) heard best over the 2nd right [intercostal space](#), which is inconsistent with this patient's rumbling [diastolic murmur](#) heard best at the apex.

C - Cardiac tumor

Explanation Why

[Cardiac myxomas](#) are the most common primary cardiac [tumor](#), commonly located in the [left atrium](#). They may be pedunculated, in which case they often present with position-dependent symptoms of obstructed blood flow through the [atrioventricular valves](#), or they may embolize. Additionally, patients may present with constitutional findings (e.g., weight loss, [fever](#), or [anemia](#)). The typical auscultatory finding (also present in this patient) is an early [diastolic](#) “plop”, caused by a movement of the mobile [tumor](#) mass, followed by a mid-[diastolic](#) rumbling [murmur](#) best heard at the apex that results from the obstructed blood flow through the [mitral valve](#), and mimics the [murmur](#) of [mitral](#)

[stenosis](#).

D - Adverse medication effect

Explanation Why

Various drugs can cause (orthostatic) [hypotension](#) and may lead to [syncope](#), including alpha-1 blockers. This patient is taking an alpha-1 blocker and presents with [orthostatic hypotension](#). However, among the alpha-1 blockers, [tamsulosin](#) is the agent with the lowest potential of inducing [hypotension](#) because of its increased selectivity for [alpha-1 receptors](#) of the [bladder neck](#) and the [prostatic urethra](#). The peculiar [heart murmur](#) of a [diastolic](#) “plop” followed by a rumble suggests a different diagnosis.


E - Left ventricular dilation

Explanation Why

[Dilated cardiomyopathy](#) is most commonly idiopathic, but can also be alcohol-induced. It usually presents with signs of heart failure. While this patient consumes critical levels of alcohol, and displays symptoms that may suggest impaired pump function of the [heart](#) ([syncope](#), [hypotension](#)), [dilated cardiomyopathy](#) does not manifest with a position-dependent [diastolic heart murmur](#).

Question # 16

A 35-year-old man is brought to the emergency department 20 minutes after being involved in a motor vehicle collision in which he was a restrained passenger. The patient is confused. His pulse is 140/min and blood pressure is 85/60 mm Hg. Examination shows a hand-sized hematoma on the anterior chest wall. An ECG shows sinus tachycardia. Which of the following structures is most likely injured in this patient?

	Answer	Image
A	Papillary muscle	
B	Aortic isthmus	
C	Aortic valve	
D	Inferior vena cava	
E	Left main coronary artery	

Hint

This patient's presentation of shock (hypotension, tachycardia, altered mental status) after severe blunt chest trauma in the setting of a motor vehicle accident is highly suggestive of aortic injury.

Correct Answer


A - Papillary muscle

Explanation Why

[Papillary muscle rupture](#) is a notorious complication after a [myocardial infarction](#) and presents with acute [hypotension](#), new-onset [mitral regurgitation](#), and subsequent [pulmonary edema](#). Rapid deceleration injury in a motor vehicle accident is not associated with [papillary muscle rupture](#).

B - Aortic isthmus

Image



Explanation But

Perform a CT angiography on those patients who are hemodynamically stable or a [TEE](#) if hemodynamically unstable to assess the severity of injury.

Explanation Why

The [aortic isthmus](#), a physiological narrowing at the final section of the [aortic arch](#), is located just [distal](#) to the origin of the [left subclavian artery](#) and is tethered by the [ligamentum arteriosum](#). It is the most common site of injury due to blunt aortic trauma because it is extremely vulnerable to rapid deceleration injury, which can occur in motor vehicle accidents when the chest smashes against the steering wheel or is abruptly restrained by the seatbelt.

C - Aortic valve

Explanation Why

In rare cases, rapid deceleration injury in the setting of a motor vehicle accident, as seen in this patient, can lead to a large aortic tear that extends to the [aortic valve](#). Nevertheless, this is not a common site of injury. Moreover, a [diastolic murmur](#) would be heard due to new-onset [aortic regurgitation](#).

D - Inferior vena cava


Explanation Why

Injury to the [inferior vena cava](#) may occur in patients after blunt decelerating trauma, but it is extremely rare. Aside from that, the [hematoma](#) seen on the patient's chest would not be typical as injury to the [inferior vena cava](#) would likely result in [retroperitoneal hematoma](#).

E - Left main coronary artery

Explanation Why

Although the [coronary arteries](#) may also be affected together with aortic injury, the absence of ST-segment abnormalities on this patient's [ECG](#) makes involvement of the [left main coronary artery](#) unlikely.



Question # 17

A 55-year-old woman comes to the emergency room 30 minutes after the sudden onset of chest pain radiating to the left shoulder. Prior to the onset of her symptoms, she was lying in bed because of a migraine headache. Episodes of similar chest pain usually resolved after a couple of minutes. She has smoked one pack of cigarettes daily for 20 years. Her only medication is sumatriptan. An ECG shows ST-segment elevations in the anterior leads. Serum troponins are negative on two successive blood draws and ECG shows no abnormalities 30 minutes later. Administration of which of the following is most likely to prevent further episodes of chest pain in this patient?

	Answer	Image
A	Ramipril	
B	Clopidogrel	
C	Aspirin	
D	Propranolol	
E	Diltiazem	

Hint

Typical anginal symptoms with reversible ST-elevations and negative troponin tests are typical for vasospastic angina (Prinzmetal angina). Known triggers for vasospastic episodes include cocaine, alcohol, exposure to cold, hyperventilation, and triptans (as seen in this patient, who likely took sumatriptan for her migraine).

Correct Answer

A - Ramipril

Explanation Why

ACE inhibitors such as [ramipril](#) are commonly prescribed in cases of [Takotsubo cardiomyopathy](#), which also presents with [angina](#) and [ECG](#) abnormalities such as [ST elevations](#). However, [Takotsubo cardiomyopathy](#) typically causes elevations of [troponin T](#) and I. ACE inhibitors do not play a role in the treatment of [vasospastic angina](#).

B - Clopidogrel

Explanation Why

[Vasospastic angina](#) is not treated with [clopidogrel](#) ([P2Y12 receptor antagonists](#)) because vasospasms, not abnormal [platelet aggregation](#), are causing the transient [ischemic](#) symptoms seen here. [P2Y12 receptor antagonists](#) are used in the treatment of [acute coronary syndrome](#) (usually in combination with [aspirin](#)) and to prevent thrombosis following stent placement. [ACS](#) presents with [angina](#) and [ECG](#) abnormalities such as [ST elevations](#), which are both present in this patient. However, the [ECG](#) of a [STEMI](#) would not return to normal 30 minutes later. Additionally, a [STEMI](#) would cause elevations of [troponin T](#) and I, which are absent in this case.

C - Aspirin

Explanation Why

[Aspirin](#) is used for both the initial management and long-term prevention of [acute coronary syndrome](#), as it inhibits [platelet aggregation](#). [ACS](#) can present with [angina](#) and [ST elevations](#), which are seen in this patient. However, in [STEMI](#), the [ECG](#) would not return to normal 30 minutes later, nor would serum levels of [troponin](#) be normal. High-dose [aspirin](#) inhibits the synthesis of [prostacyclins](#), potentially even inducing vasospasms, and therefore should be avoided in [vasospastic angina](#).

D - Propranolol

Explanation Why

[Nonselective beta blockers](#) such as [propranolol](#) should not be used in [vasospastic angina](#) because beta blockade prevents relaxation of the vascular [smooth muscle cells](#). [Smooth muscle cell](#) contraction is the root cause of coronary spasms, which means that [beta blockers](#) would worsen [vasospastic angina](#). [Beta blockers](#) are indicated, however, for the treatment of [acute coronary syndrome](#) to reduce [heart rate](#) and subsequent cardiac oxygen consumption. [ACS](#) can present with [angina](#) and [ST elevations](#), which are seen in this patient. However, in [STEMI](#), the [ECG](#) would not return to normal after 30 minutes, nor would serum levels of [troponin](#) be normal.

E - Diltiazem

Explanation But

The presence of [vasospastic angina](#) can be confirmed during [coronary angiography](#) with provocative tests (usually using [acetylcholine](#); less commonly ergonovine or [hyperventilation](#)).


Explanation Why

[Calcium channel blockers](#) (CCBs) such as [diltiazem](#) are the first-line treatment for [vasospastic angina](#). They are effective for both acute attacks and prophylaxis. [Triptans](#) should also be discontinued and the patient should be advised to stop smoking, which is a known [risk factor](#) for [vasospastic angina](#). [Nitroglycerin](#) is often used for the acute abortive treatment of [vasospastic angina](#). Long-acting [nitrates](#) (e.g., [isosorbide mononitrate](#)) are only used as second-line agents for controlling vasospastic episodes because their long-term use is limited by the development of [nitrate](#) tolerance limits.

Question # 18

A 51-year-old man comes to the emergency department because of a 3-day history of shortness of breath, fever, and chills. He has no history of serious illness. His temperature is 39.5°C (103.1°F). Physical examination shows a grade 4/6, holosystolic, blowing murmur over the apex that radiates to the axilla. Crackles are heard in both lower lung fields. Examination of the extremities shows several nontender, nonblanching, erythematous macules on the palms and soles. Histopathologic examination of these macules is most likely to show which of the following?

	Answer	Image
A	Capillary wall rupture with fibrin thrombus formation	
B	Microabscesses with neutrophil infiltration of capillaries	
C	Epithelioid cells with surrounding multinucleated giant cells	

	Answer	Image
D	Cleft-like vacuoles within the arterioles	
E	Spiral-shaped bacteria with axial filaments in the epidermis	

Hint

This patient presents with high fever, dyspnea, and a mitral regurgitation murmur, which suggests acute infective endocarditis of the mitral valve, complicated by pulmonary edema. The nontender, erythematous macules on the palms and soles of this man are most likely Janeway lesions.

Correct Answer

A - Capillary wall rupture with fibrin thrombus formation

Image



Explanation Why

[Roth spots](#) are the fundoscopic finding of white-centered retinal hemorrhages that are characterized histologically by microthrombi that form at the site of retinal [capillary](#) damage. They can be seen with [infective endocarditis](#), as well as with leukemia, anoxic brain injury, [carbon monoxide poisoning](#), and [HIV](#). In contrast, [Janeway lesions](#) do not affect the [eye](#) and hemorrhage is not believed to be a primary component of [Janeway lesions](#).

B - Microabscesses with neutrophil infiltration of capillaries

Image



Explanation Why

Microabscesses with [neutrophil](#) infiltration of [capillaries](#) and areas of hemorrhage are characteristic of the [Janeway lesions](#) seen in patients with [infective endocarditis](#). They are thought to be caused by [septic microemboli](#) from valve vegetations and, among other vascular phenomena, constitute a minor clinical criterion for the modified [Duke criteria](#) in [infective endocarditis](#). [Janeway lesions](#) are more common in [acute endocarditis](#) than in [subacute endocarditis](#).

C - Epithelioid cells with surrounding multinucleated giant cells


Explanation Why

Epithelioid cells with surrounding [multinucleated giant cells](#) are characteristic of [granulomatous inflammation](#), which is associated with numerous conditions, including [tuberculosis](#), systemic fungal

infection, [sarcoidosis](#), and [granulomatosis with polyangiitis](#). However, none of these conditions present with high [fever](#) and a new-onset [heart murmur](#). [Janeway lesions](#) are not caused by [granulomatous inflammation](#).

D - Cleft-like vacuoles within the arterioles

Image



Explanation But

The vacuole appearance of [cholesterol embolism](#) occurs due to [cholesterol](#) dissolution when exposed to tissue fixation.

Explanation Why

Cleft-like vacuoles within [arterioles](#) are the histological hallmark of [cholesterol embolism](#). The most common [skin](#) manifestations of [cholesterol](#) embolization include [livedo reticularis](#), [gangrene](#), [cyanosis](#), [skin ulcer](#), [purpura](#) or [petechiae](#), and firm, painful [erythematous](#) nodules. [Janeway lesions](#) have a different appearance on [histology](#) than [cholesterol emboli](#).


E - Spiral-shaped bacteria with axial filaments in the epidermis

Explanation Why

Spiral-shaped bacteria with axial filaments on microscopic examination of a specimen would be consistent with [secondary syphilis](#), which is caused by an infection with the [spirochete *Treponema pallidum*](#). Patients with [secondary syphilis](#) present with a disseminated, [maculopapular rash](#) involving the entire trunk and extremities (including the palms and soles). However, this patient does not have any other clinical features of [secondary syphilis](#), such as [generalized lymphadenopathy](#), myalgia, or [condylomata lata](#).

Question # 19

A previously healthy 30-year-old man is brought to the emergency department 30 minutes after collapsing at home. Cardiopulmonary resuscitation is attempted for 10 minutes without success. An autopsy is performed one hour after the death of the patient. During the autopsy, examination of the heart shows pliable valves without any calcification. A photograph of sections of the heart obtained at autopsy is shown. The patient's condition is most likely associated with which of the following pathophysiologic changes?



	Answer	Image
A	Decreased duration of the myocardial action potential	
B	Significantly reduced myocardial contraction force	

	Answer	Image
C	Fixed obstruction of the left ventricular outflow tract	The diagram illustrates the pathophysiology of fixed obstruction of the left ventricular outflow tract (LVOT) in hypertrophic cardiomyopathy (HCM). It is divided into three panels: Normal, Nonobstructive type, and Obstructive type. The Normal panel shows a heart with a standard-sized interventricular septum and a clear LVOT. The Nonobstructive type panel shows a heart with a thickened interventricular septum but no obstruction of the LVOT. The Obstructive type panel shows a heart with a significantly thickened interventricular septum and the mitral valve being pushed into the LVOT during systole, causing obstruction. An inset diagram provides a closer look at the SAM mechanism, labeling the interventricular septum hypertrophy and the systolic anterior motion (SAM) of the mitral valve.
D	Marked noncompliance of the left ventricular wall	
E	Severe narrowing of the left anterior descending artery	

Hint

The severe hypertrophy of the left ventricle seen in the autopsy images of this young man who died of sudden cardiac death confirms a diagnosis of hypertrophic cardiomyopathy.

Correct Answer

A - Decreased duration of the myocardial action potential

Explanation Why

Significantly reduced duration of [myocardial action potential](#) (due to a defect in the voltage-gated sodium channel) is a pathophysiologic change in [Brugada syndrome](#), an important cause of [sudden cardiac death](#) in young adults. However, [Brugada syndrome](#) would not be associated with ventricular [hypertrophy](#), which is seen in the photograph. In patients with ventricular [hypertrophy](#), an increase in the duration of the [myocardial depolarization](#) would be expected.


B - Significantly reduced myocardial contraction force

Explanation Why

Significantly reduced [myocardial](#) contraction force ([systolic dysfunction](#)) is seen in [dilated cardiomyopathy \(DCM\)](#), which can cause [sudden cardiac death](#). However, [DCM](#) more commonly presents with [palpitations](#), exertional [dyspnea](#), and/or features of [congestive heart failure](#). A gross pathology specimen of [DCM](#) shows dilated cardiac chambers with thinned ventricular walls as opposed to the severe [hypertrophy](#) of the [left ventricle](#) seen here. In this patient's condition, a reduction in [myocardial](#) contraction force would not be expected.

C - Fixed obstruction of the left ventricular outflow tract

Image



Explanation Why

Fixed obstruction of the left ventricular outflow tract is caused by [aortic stenosis](#) (AS), which can cause [sudden cardiac death](#) in severe cases. AS can manifest in young adults as a result of calcification of a congenital [bicuspid valve](#) (in high-income countries) or as a sequela to [rheumatic heart disease](#) (in low-income countries). However, patients with severe AS typically experience [syncope](#), [angina](#), and [dyspnea](#). This man was previously healthy; in a young, asymptomatic individual with pliable, noncalcified [heart valves](#), [hypertrophic cardiomyopathy](#) is a more likely cause of [sudden cardiac death](#). Unlike AS, [hypertrophic cardiomyopathy](#) causes dynamic (not fixed) left ventricular outflow tract obstruction.

D - Marked noncompliance of the left ventricular wall

Explanation But

Chamber noncompliance is also seen in cases of [restrictive cardiomyopathy](#); however, this condition is not commonly associated with [sudden cardiac death](#) or severe thickening of the [left ventricle](#).

Explanation Why

Marked reduction in compliance (physiology) of the [left ventricular](#) wall and reduced chamber volume due to [left ventricular hypertrophy](#) are key features of [hypertrophic cardiomyopathy \(HCM\)](#). These changes prevent ventricular relaxation and decrease the volume of blood that enters the [left ventricle](#) during [diastole](#) ([diastolic dysfunction](#)). The morphology of LV [hypertrophy](#) in [HCM](#) is highly variable; patients typically develop asymmetrical septal [hypertrophy](#) but some patients develop symmetric, concentric LV [hypertrophy](#), as seen here, or apical LV [hypertrophy](#). Interventricular septum [hypertrophy](#) may narrow the left ventricular outflow tract, which increases the velocity of blood flowing past it and decreases pressure ([Venturi effect](#)). This pulls the [mitral valve](#) leaflet down during [systole](#) ([systolic anterior motion](#)) instead of allowing it to snap shut, which causes dynamic [left ventricular](#) outflow obstruction, [mitral regurgitation](#), and, in severe cases, hemodynamic collapse and death.

E - Severe narrowing of the left anterior descending artery

Explanation Why

Severe narrowing ($\geq 90\%$ stenosis) of the [left anterior descending artery](#) (LAD) can cause [myocardial infarction \(MI\)](#), which is an important cause of [sudden cardiac death](#). Since the earliest gross changes of [MI](#) (e.g., a dark mottling) are only seen 12 hours after death, an [MI](#) cannot be ruled out from the specimen in the photograph, which was obtained one hour after the patient's death. This patient's condition, i.e., [hypertrophic cardiomyopathy \(HCM\)](#), can result in [MI](#) due to increased oxygen demand in the [hypertrophied](#) ventricle and impaired coronary vessel perfusion (due to impaired ventricular [diastole](#) and obstruction of blood flow to the aorta), but [HCM](#) is not associated with narrowing of the large [coronary arteries](#) such as the LAD. Instead, severe narrowing of the large [coronary arteries](#) is a feature of atherosclerotic disease, which would not be expected in this young patient.

Question # 20

An autopsy of a patient's heart who recently died in a motor vehicle accident shows multiple nodules near the line of closure on the ventricular side of the mitral valve leaflet. Microscopic examination shows that these nodules are composed of immune complexes, mononuclear cells, and thrombi interwoven with fibrin strands. These nodules are most likely to be found in which of the following patients?

	Answer	Image
A	A 54-year-old male who recently underwent dental surgery	
B	A 71-year-old male with acute-onset high fever and nail bed hemorrhages	
C	A 41-year-old female with a facial rash and nonerosive arthritis	
D	A 28-year-old HIV-positive female intravenous drug user	
E	A 62-year-old male with <i>Cardiobacterium hominis</i> bacteremia	
F	A 6-year-old female with subcutaneous nodules and erythema marginatum	

Hint

Nonbacterial verrucous thrombi on the undersurface of the mitral valve are suggestive of Libman-Sacks endocarditis.

Correct Answer

A - A 54-year-old male who recently underwent dental surgery

Explanation Why

Patients with congenital [heart](#) defects or [prosthetic heart valves](#) are [prone](#) to *S. viridans* [bacteremia](#) after dental surgery. *S. viridans* is the most common cause of subacute [infectious endocarditis](#). [Viridans streptococci](#) typically infect [platelet-fibrin](#) aggregates on [heart valves](#) with pre-existing damage. The microscopic exam in this deceased patient showed no evidence of bacteria, which rules out [infectious endocarditis](#).

B - A 71-year-old male with acute-onset high fever and nail bed hemorrhages

Explanation Why

[Splinter hemorrhages](#) in the setting of high [fever](#) are suspicious for acute [infectious endocarditis](#). *S. aureus* is the most common cause of acute [infectious endocarditis](#), which often affects previously normal valves. Unlike the nonbacterial [thrombi](#) found on autopsy, [infectious endocarditis](#) would show bacteria on microscopic examination.

C - A 41-year-old female with a facial rash and nonerosive arthritis

Explanation But

Other diseases with systemic [inflammation](#), such as advanced cancer or [rheumatoid arthritis](#), can also precipitate [Libman-Sacks endocarditis](#).

Explanation Why

A facial rash and symmetric, nonerosive polyarthritis in a middle-aged female is highly suspicious for [systemic lupus erythematosus](#). Systemic [inflammation](#) in [SLE](#) can lead to the deposition of circulating sterile [immune complexes](#) with [platelet thrombi](#) on either side of valve leaflets in what is

known as [Libman-Sacks endocarditis](#). Although the vegetations seen in [Libman-Sacks endocarditis](#) are usually asymptomatic, they are easily dislodged and can embolize, causing complications. Valvular damage is rare but can sometimes manifest as [mitral regurgitation](#) or [mitral stenosis](#).

D - A 28-year-old HIV-positive female intravenous drug user

Explanation Why

Injection drug use puts patients at high risk for [infectious endocarditis](#) and [HIV](#)-positive individuals are especially susceptible. The most common causal pathogen is [S. aureus](#). However, [infectious endocarditis](#) in intravenous drug users most often involves the [tricuspid valve](#) instead of the [mitral valve](#), as seen here. Furthermore, the microscopic exam in this deceased patient would have shown evidence of bacteria rather than nonbacterial [thrombi](#).

E - A 62-year-old male with *Cardiobacterium hominis* bacteremia

Explanation Why

Cardiobacterium hominis belongs to the [HACEK organisms](#), a class of bacteria that cause [infectious endocarditis](#). These organisms require longer times to grow in culture and can even result in negative cultures. [Infectious endocarditis](#) would present with bacteria on microscopic examination, as opposed to the nonbacterial (i.e., sterile) [thrombi](#) found on autopsy here.

F - A 6-year-old female with subcutaneous nodules and erythema marginatum

Explanation But

[Rheumatic heart disease](#) can rarely cause [nonbacterial thrombotic endocarditis](#) but only after years of disease. This 6-year-old presents with early symptoms of [acute rheumatic fever](#), and therefore, would not have such lesions.

Explanation Why

Subcutaneous nodules and [erythema marginatum](#) are manifestations of [acute rheumatic fever](#), which

results from the formation of [autoantibodies](#) after untreated group A [streptococcus](#) tonsillopharyngitis. Unlike the isolated [endocarditis](#) found on autopsy, the nonspecific carditis seen in [acute rheumatic fever](#) tends to involve the [pericardium](#), [epicardium](#), [myocardium](#), and [endocardium](#). [Acute rheumatic fever](#) is caused by a [type II hypersensitivity reaction](#), whereas the [immune complex](#) deposits seen on this patient's autopsy are indicative of a [type III hypersensitivity reaction](#). Also, microscopic examination would show [myocardial granulomas](#) ([Aschoff bodies](#)) and [Anitschkow cells](#).

Question # 21

A 3-year-old boy is brought to the physician because of recurrent nosebleeds and fatigue for the past 2 months. He also frequently complains his head hurts. The patient has met all motoric milestones for his age but does not like to run because his legs start to hurt if he does. He is at the 40th percentile for both height and weight. His temperature is 37.0°C (98.6°F), pulse is 125/min, respirations are 32/min, and blood pressure in the right arm is 130/85 mm Hg. A grade 2/6 systolic murmur is heard in the left paravertebral region. Further evaluation of this patient is most likely to show which of the following findings?

	Answer	Image
A	Inferior rib notching	
B	Interarm difference in tissue oxygenation	
C	Pulmonary valve stenosis	
D	Left-axis deviation on ECG	
E	Delayed pulse in lower extremities	

Hint

This patient's general fatigue, hypertension, low-grade systolic ejection murmur, lower-extremity claudication, headaches, and recurrent epistaxis together suggest coarctation of the aorta.

Correct Answer

A - Inferior rib notching

Explanation Why

Notching of the [ribs](#), an [x-ray](#) finding in patients with [coarctation of the aorta](#), results from compensatory collateral blood flow involving the intercostal vessels. Those vessels may dilate to bypass the left ventricular outflow tract obstruction. Although this patient may eventually develop visible [rib notching](#), it is generally a sign of chronic disease and usually does not become apparent in children before the age of 5 years.

B - Interarm difference in tissue oxygenation

Explanation Why

Interarm differences in tissue oxygenation and blood pressure can be a finding of [coarctation of the aorta](#) in the rare case that the narrowing is [proximal](#) to the [left subclavian artery](#). In such patients, blood pressure in the right arm would be markedly higher than in the left arm and [oxygen saturation](#) would be lower in the left arm. However, [coarctation of the aorta](#) most commonly presents with differences in blood pressure between the upper and the lower extremities.

C - Pulmonary valve stenosis

Explanation Why

Congenital [pulmonary valve stenosis](#) is a characteristic feature of [tetralogy of Fallot](#). A [systolic ejection murmur](#) can be present and is usually best heard at the left sternal border, rather than the left paravertebral region. Patients usually present with (mild) [cyanosis](#) and intermittent [hypoxic](#) episodes that improve with squatting, none of which are seen here. [Coarctation of the aorta](#) is characterized by an outflow obstruction of the [left ventricle](#).

D - Left-axis deviation on ECG

Explanation Why

Left-axis deviation and increased R wave amplitude in V5–V6 on [ECG](#) are signs of [left ventricular hypertrophy](#). While [myocardial hypertrophy](#) is one of the compensatory mechanisms to counter the [aortic narrowing](#) in patients with [coarctation of the aorta](#), such as this 3-year-old boy, the changes in [ECG](#) vary with age and degree of stenosis, and it usually takes several years before the [hypertrophy](#) becomes apparent. In older children and adults, [ECG](#) may show signs of [left ventricular hypertrophy](#) in the left [precordial leads](#) (e.g., positive [Sokolow-Lyon](#) criteria).

E - Delayed pulse in lower extremities

Explanation Why

In patients with [coarctation of the aorta](#), delayed or even absent lower extremity pulses are expected [distal](#) to the stenotic portion of the aorta. Since the post-stenotic blood supply is limited, the resulting lower-extremity [hypoxia](#) becomes evident, especially during exertion, which is a strong sign of [intermittent claudication](#), as in this patient. Doppler [echocardiography](#) is the [confirmatory test](#) of choice and can identify the location and extent of the stenosis.

Question # 22

Two days after being admitted for acute myocardial infarction, a 61-year-old man has sharp, substernal chest pain that worsens with inspiration and improves when leaning forward. Cardiac examination shows a scratchy sound best heard over the left sternal border. Histopathological examination of the affected tissue is most likely to show which of the following findings?

	Answer	Image
A	Neutrophilic infiltration	
B	Normal myocardium	
C	Hypercontracted myofibrils	
D	Collagenous scar tissue	
E	Granulation tissue with macrophages	

Hint

This patient has symptoms consistent with acute pericarditis, including a pericardial friction rub on auscultation and sharp, pleuritic chest pain that worsens with inspiration and improves with sitting and leaning forward. Given his history of myocardial infarction (MI) 2 days ago, it is most likely a subtype known as fibrinous pericarditis.

Correct Answer

A - Neutrophilic infiltration

Explanation Why

Neutrophilic infiltration is seen 1–3 days after a [myocardial infarction \(MI\)](#) on histopathological examination. After the [myocardium](#) undergoes [coagulative necrosis](#), it is surrounded by an infiltrate of acute inflammatory cells. Release of inflammatory mediators (e.g., [TNF- \$\alpha\$](#) , [IL-2](#), [IL-6](#)) from [neutrophils](#) causes exudation of [fibrin](#) and [leukocytes](#) into the [pericardial space](#), resulting in fibrinous pericarditis.

B - Normal myocardium

Explanation Why

A normal [myocardium](#) is seen on histopathological examination within the first 4 hours after an [MI](#) but not 2 days later.

C - Hypercontracted myofibrils

Explanation Why

Hypercontracted [myofibrils](#) are seen on histopathological examination up to 24 hours after an [MI](#) if there is [reperfusion injury](#); they would not be seen 2 days later.

D - Collagenous scar tissue

Explanation Why

Collagenous [scar](#) tissue does not occur 2 days post-[MI](#) but is seen on histopathological examination 2 weeks to several months after an [MI](#). In this time frame, the [myocardium](#) has completely formed a contraction [scar](#), and complications such as [congestive heart failure \(CHF\)](#), [arrhythmias](#), and true [ventricular aneurysms](#) can develop.

E - Granulation tissue with macrophages

Explanation Why

[Granulation tissue](#) with [macrophages](#) does not occur 2 days post-[MI](#) but is seen 3–14 days after an [MI](#) on histopathological examination. During this time period, [macrophage](#)-mediated degradation of tissue makes the [myocardium](#) more vulnerable to [cardiac tamponade](#), [papillary muscle rupture](#), left-to-right shunt (due to interventricular septal rupture), and left ventricular pseudoaneurysm formation.

Question # 23

A 61-year-old woman dies unexpectedly. She had not seen a physician in 10 years. Examination of the heart during autopsy shows hypertrophy of the left ventricular wall and enlargement of the left ventricular chamber. Microscopic examination shows lengthening of individual muscle fibers due to duplication of sarcomeres in series. Which of the following is the most likely underlying cause of these findings?

	Answer	Image
A	Increased pulmonary arterial pressure	
B	Decreased left ventricular contractility	
C	Decreased left ventricular compliance	
D	Increased systemic vascular resistance	
E	Increased left ventricular end diastolic volume	

Hint

Cardiac hypertrophy with an associated increase in chamber size due to sarcomeres being added in series indicates eccentric left ventricular hypertrophy (LVH).

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

Correct Answer

A - Increased pulmonary arterial pressure

Explanation Why

Increased [pulmonary artery](#) pressure, which can be seen in [pulmonary hypertension](#) and/or [left ventricular heart failure](#), causes an increase in [right ventricular afterload](#) and can result in RV [concentric hypertrophy](#). It does not affect the [left ventricle](#) and thus would not result in the LV [eccentric hypertrophy](#) seen here.

B - Decreased left ventricular contractility

Explanation Why

Decreased cardiac contractility typically occurs secondary to direct [myocardial](#) damage or cardiac remodeling (e.g., [eccentric hypertrophy](#)) but does not itself cause [eccentric hypertrophy](#). Instead, [eccentric hypertrophy](#) is caused by a process that initially increases rather than decreases cardiac contractility.

C - Decreased left ventricular compliance

Explanation Why

Decreased ventricular compliance is characteristic of [restrictive cardiomyopathy](#). [Restrictive cardiomyopathy](#) results in decreased [diastolic](#) filling (decreased [end-diastolic volume](#)) with atrial congestion and subsequent atrial enlargement, which ultimately leads to severe [diastolic dysfunction](#). [Left ventricular systolic](#) function is usually preserved and [eccentric hypertrophy](#) would not be expected.

D - Increased systemic vascular resistance

Explanation Why

An increase in [systemic vascular resistance](#) (e.g., [arterial hypertension](#)) results in an increase in LV [afterload](#), which increases the [myocardial](#) wall tension and [stroke volume](#) via the [Frank-Starling mechanism](#). Ultimately, it triggers cardiac remodeling, which results in [concentric hypertrophy](#) (addition of [sarcomeres](#) in parallel) and [hypertrophic cardiomyopathy](#) rather than [eccentric hypertrophy](#).


E - Increased left ventricular end diastolic volume


Explanation Why

Increased [left ventricular end-diastolic volume](#) ([preload](#)) results in dilation of the [left ventricle](#) in an attempt to increase [myocardial contractility](#) to maintain [cardiac output](#) ([Frank-Starling mechanism](#)). This compensatory mechanism ultimately fails, and the continuous strain of increased blood volume in the LV triggers myocardial remodeling that results in [eccentric hypertrophy](#) (addition of [sarcomeres](#) in series), eventually leading to a [dilated cardiomyopathy](#) and reduced [ejection fraction](#). Most cases of [dilated cardiomyopathy](#) are idiopathic, but common identifiable causes include exposure to toxic substances (alcohol), metabolic agents, and infectious pathogens that cause structural damage to the [myocardium](#) or valves.

Question # 24

A 13-year-old boy is brought to the emergency department by ambulance after suddenly losing consciousness while playing in a soccer tournament. The patient has had 2 episodes of syncope without a discernable trigger over the past year. He has been otherwise healthy. His father died suddenly at the age of 37. He reports lightheadedness and suddenly loses consciousness when physical examination is attempted. Radial pulses are not palpable. An ECG shows ventricular tachycardia with peaks of the QRS twisting around the isoelectric line. Which of the following is the most likely underlying cause of this patient's condition?

	Answer	Image
A	Anomalous origin of the left coronary artery	
B	Accessory atrioventricular conduction pathway	
C	Myofibrillar disarray of cardiac septum	
D	Bicuspid aortic valve	
E	Defect in interatrial septum	

	Answer	Image
F	Mutation of myocardial potassium channels	 The image is a 12-lead ECG tracing on a pink grid. The leads are arranged in a standard format: I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. The tracing shows a prolonged QT interval, which is a characteristic finding in certain mutations of myocardial potassium channels. The text 'Courtesy of Jason E. Roediger, CCT, CRAT' is visible at the top of the tracing.

Hint

Torsades de pointes arrhythmia on ECG (peaks of the QRS twisting around the isoelectric line), a history of unexplained syncope, and a suspected family history of sudden cardiac death are most concerning for Romano-Ward syndrome.

Correct Answer


A - Anomalous origin of the left coronary artery

Explanation Why

[Anomalous origin of the left coronary artery from the pulmonary artery \(ALCAPA\)](#) is a congenital anomaly that typically manifests in [infants](#) with pallor, irritability, and [diaphoresis](#) after feeding and can progress to [congestive heart failure](#). Rarely, some [infants](#) outgrow these symptoms and are asymptomatic until adulthood, when they present with [syncope](#) or [sudden cardiac death](#). [ALCAPA](#) is unlikely to cause [torsades de pointes](#) and cannot explain this patient's [family history](#) of [sudden cardiac death](#).

B - Accessory atrioventricular conduction pathway

Image



Explanation Why

An accessory atrioventricular conduction pathway ([bundle of Kent](#)) is the pathophysiology of [Wolff-Parkinson-White syndrome \(WPW\)](#), which can present with [syncope](#). However, [ECG](#) of a patient with [WPW syndrome](#) would be unlikely to reveal [torsades de pointes](#). Instead, it typically shows a decreased [PR interval](#), delta wave (i.e., slurring slow rise of the initial portion of the QRS), and QRS prolongation. Furthermore, [WPW syndrome](#) is typically sporadic and only follows an [autosomal dominant](#) pattern of inheritance in rare cases.

C - Myofibrillar disarray of cardiac septum

Explanation Why

Myofibrillar disarray of the cardiac septum is a feature of [hypertrophic cardiomyopathy \(HCM\)](#), which is a common cause of [sudden cardiac death](#) in young athletes, and can present with [dyspnea](#), [dizziness](#), and [syncope](#), as seen in this patient. Furthermore, it should be suspected in a patient with a [family history](#) of [sudden cardiac death](#), as it follows an [autosomal dominant](#) pattern of inheritance in approx. 50% of cases. However, [HCM](#) is unlikely to cause [torsades de pointes arrhythmia](#).

D - Bicuspid aortic valve

Explanation Why

A [bicuspid aortic valve](#) can cause congenital or acquired [aortic stenosis \(AS\)](#), which can present with [syncope](#) on exertion. However, [syncope](#) typically occurs in the severe form of AS, and patients would usually have gradual onset of symptoms such as exertional dyspnea, exertional [chest pain](#), easy fatigability, and exercise intolerance. Furthermore, AS is typically sporadic (unlikely to explain this patient's [family history](#) of [sudden cardiac death](#)), and [torsades de pointes arrhythmia](#) is not a typical feature of AS.


E - Defect in interatrial septum

Explanation Why

A defect in the [interatrial septum](#) (i.e., [atrial septal defect](#)) is sometimes seen in atrial [arrhythmias](#) such as [atrial fibrillation](#) and/or [atrial flutter](#). However, ASD would not explain this patient's [torsades de pointes arrhythmia](#), nor would it account for his [family history](#) of [sudden cardiac death](#).

F - Mutation of myocardial potassium channels

Image



Explanation But

Jervell and Lange-Nielson (JLN) syndrome follows an [autosomal recessive](#) pattern of inheritance and is associated with congenital deafness. [Romano-Ward syndrome](#) follows an [autosomal dominant](#) pattern of inheritance and is not associated with deafness.

Explanation Why

[Torsades de pointes](#) is most often due to [long QT syndrome](#), which may be acquired or congenital. [Congenital long QT syndrome](#) can be caused by mutations of [genes](#) coding for cardiac potassium, sodium, or [calcium ion](#) channels. These mutations cause ventricular [action potentials](#) to be prolonged, leading to a lengthened [QT interval](#) on [ECG](#). The two most common long QT syndromes are [Romano-Ward syndrome](#) and Jervell and Lange-Nielson (JLN) syndrome.

Question # 25

A 42-year-old man comes to the physician because of a 6-month history of progressively worsening shortness of breath with exertion. He was diagnosed with systemic sclerosis 5 years ago. Vital signs are within normal limits. Physical examination shows puffy, taut skin over the fingers. Pulmonary examination is unremarkable. There is no jugular venous distention. An x-ray of the chest shows enlargement of the pulmonary vessels and a prominent right heart border. Cardiac catheterization shows elevated right ventricular pressures and a mean pulmonary arterial pressure of 55 mm Hg. Treatment with tadalafil is begun. The expected beneficial effect of this drug is most likely due to which of the following actions?

	Answer	Image
A	Blockade of endothelin-1 binding at endothelin receptors	
B	Decreased smooth muscle sensitivity to norepinephrine	
C	Increased activation of protein kinase A	<pre> graph TD subgraph Extracellular_space [Extracellular space] Ligand end subgraph Cytosol [Cytosol] GPCR[GPCR] Gs[Gs] Gi[Gi] Adenylate_cyclase[Adenylate cyclase] ATP[ATP] cAMP[cAMP] PKA[PKA] Enzyme[Enzyme] Phosphodiesterase[Phosphodiesterase] AMP[AMP] end Ligand --> GPCR GPCR --> Gs Gs --> Adenylate_cyclase Adenylate_cyclase --> cAMP ATP --> cAMP cAMP --> PKA PKA --> Enzyme cAMP --> Phosphodiesterase Phosphodiesterase --> AMP </pre>
D	Reduced activity of angiotensin converting enzyme	

	Answer	Image
E	Enhanced activity of nitric oxide	The diagram illustrates the signaling pathway for nitric oxide (NO) in a smooth muscle cell. It shows the endothelium producing NO from L-arginine via eNOS. NO then enters the smooth muscle cell, where it activates guanylyl cyclase, converting GTP to cGMP. cGMP then activates protein kinase G, which inhibits phospholipase C and the sarcoplasmic reticulum Ca²⁺ release channel. This leads to a decrease in intracellular Ca²⁺ concentration, resulting in smooth muscle relaxation and vasodilation. The diagram also shows the breakdown of cGMP by PDE5 and the effects of nitrates and PDE5 inhibitors. Legend: - Effects of cGMP - Breakdown of cGMP - Physiological NO production

Hint

This patient has features of pulmonary arterial hypertension (PAH; group 1 pulmonary hypertension), including progressive dyspnea, enlarged pulmonary vessels, enlarged right side of the heart, and a history of systemic sclerosis (a cause of PAH). Elevated right ventricular and pulmonary arterial pressures confirm the diagnosis. Tadalafil is a phosphodiesterase type 5 inhibitor (PDE5 inhibitor) that has been shown to improve outcomes in patients with PAH.

Correct Answer

A - Blockade of endothelin-1 binding at endothelin receptors

Explanation Why

As endothelins are potent vasoconstrictors, blockade of endothelin-1 binding at the endothelin receptor causes [vasodilation](#). This is the mechanism of action of [endothelin receptor antagonists](#) (e.g., [bosentan](#), ambrisentan), a class of drugs used to treat [pulmonary arterial hypertension](#). However, this is not the mechanism of action of [tadalafil](#).


B - Decreased smooth muscle sensitivity to norepinephrine

Explanation Why

Decreasing [smooth muscle](#) sensitivity to [norepinephrine](#) is the mechanism of action of alpha 1 antagonists, not [tadalafil](#). Although alpha-1 antagonists cause [vasodilation](#), they are not used in the treatment of [pulmonary hypertension](#).

C - Increased activation of protein kinase A

Image



Explanation Why

Increased activation of [protein kinase A](#) is the mechanism of action of [prostacyclin analogs](#), a class of drugs used to treat [pulmonary hypertension](#). [Prostacyclin analogs](#) bind to [prostacyclin](#) receptors and activate [G protein](#), which in turn stimulates [adenylyl cyclase](#), resulting in increased intracellular [cAMP](#). Intracellular [cAMP](#) activates [protein kinase A](#), which causes [vasodilation](#) by inhibiting [myosin light chain kinase](#), the structure responsible for [smooth muscle](#) contraction. However, this is not the mechanism of action of [tadalafil](#).

D - Reduced activity of angiotensin converting enzyme


Explanation Why

Reduced activity of [angiotensin-converting enzyme](#) is the mechanism of action of [angiotensin-](#)

[converting enzyme](#) (ACE) inhibitors, a class of drugs used to treat [hypertension](#). ACE inhibitors decrease the conversion of [angiotensin I](#) to [angiotensin II](#), resulting in [vasodilation](#), reduced secretion of [aldosterone](#), and reduced breakdown of [bradykinin](#). However, this is not the mechanism of action of [tadalafil](#), and ACE inhibitors are not indicated for the treatment of [pulmonary hypertension](#).

E - Enhanced activity of nitric oxide

Image



Explanation But

[PDE5 inhibitors](#), including [tadalafil](#), are contraindicated in patients receiving [nitrates](#) due to a cumulative effect that greatly increases the risk of profound [hypotension](#). Drugs of this class may also cause [orthostatic hypotension](#) in patients taking [alpha-adrenergic](#) antagonists (e.g., [BPH](#)).

Explanation Why

Enhanced activity of [nitric oxide](#) (NO) is the mechanism of action of [PDE5 inhibitors](#), including [tadalafil](#). NO acts on [guanylyl cyclase](#) to increase the synthesis of [cGMP](#), which in turn causes [vasodilation](#) by causing a reduction in intracellular Ca^{2+} concentration and reducing the sensitivity of the vascular [smooth muscle cells](#) to calcium. [PDE5 inhibitors](#) prevent the breakdown of [cGMP](#) and thus prolong the effect of NO-mediated [vasodilation](#). Pulmonary [vasodilation](#) decreases [pulmonary](#)

[vascular resistance](#), which decreases [pulmonary hypertension](#). Because of their [vasodilatory](#) effect, [PDE5 inhibitors](#) are also used to treat [erectile dysfunction](#).

Question # 26

A 11-year-old girl comes to the physician for evaluation of recurrent nosebleeds since childhood. She has multiple, small dilated capillaries on the lips, nose, and fingers on exam. Her father has a similar history of recurrent nosebleeds. Which of the following conditions is this patient at increased risk for?

	Answer	Image
A	Gastrointestinal polyps	
B	High-output heart failure	
C	Pheochromocytoma	
D	Lymphoma	
E	Glaucoma	
F	Renal cell carcinoma	

Hint

This autosomal dominant condition that is causing this patient's telangiectasias is associated with arteriovenous malformations.

Correct Answer

A - Gastrointestinal polyps

Explanation Why

The primary symptom of [Peutz-Jeghers syndrome](#) is [hematochezia](#) due to bleeding from hamartomatous [colonic polyps](#), rather than [epistaxis](#) due to [telangiectasias](#) as seen in this patient. [Peutz-Jeghers](#) also causes changes in bowel habits such as [constipation](#) or [diarrhea](#), but this patient does not have any gastrointestinal symptoms. Finally, although most patients with [Peutz-Jeghers](#) have mucocutaneous [hyperpigmentation](#) on their [lips](#), buccal mucosa, palms, and soles, they do not classically have true [telangiectasias](#) or [arteriovenous malformations](#).

B - High-output heart failure

Explanation Why

This patient has recurrent episodes of [epistaxis](#) and [telangiectasias](#) of the [lips](#), nose, and fingers, symptoms that are classic for [hereditary hemorrhagic telangiectasia](#) (also known as [Osler-Weber-Rendu syndrome](#)). Patients with this syndrome may also present with pulmonary or hepatic arteriovenous shunts, which may lead to [high-output cardiac failure](#). Other complications associated with this syndrome include [brain abscesses](#), [stroke](#) due to [paradoxical emboli](#) (emboli passing through a pulmonary arteriovenous shunt into [systemic circulation](#), bypassing the [lungs](#)), and [anemia](#) due to chronic [gastrointestinal bleeding](#) from [telangiectasias](#) in the [GI tract](#).

C - Pheochromocytoma

Explanation Why

[Pheochromocytomas](#) are seen in multiple hereditary disorders, including [multiple endocrine neoplasias](#) types 2A and 2B ([MEN 2A](#), [MEN 2B](#)), [von Hippel-Lindau disease](#) ([VHL](#)), and [neurofibromatosis](#) type 1 ([NF1](#)). The cutaneous findings of [NF1](#) are cutaneous [neurofibromas](#), café-au-lait spots, and [Lisch nodules](#). Patients with [MEN 2A](#) present with [medullary thyroid cancer](#) and parathyroid [hyperplasia](#), while [MEN 2B](#) causes [medullary thyroid cancer](#), oral/intestinal [neuromas](#),

and [marfanoid habitus](#). [VHL](#) also presents with [hemangioblastomas](#) of the [retina](#) or [CNS](#), as well as renal/[pancreatic](#) cysts. Furthermore, they are at an increased risk for [renal cell carcinoma](#). None of these conditions are associated with mucocutaneous vascular [malformations](#) or [epistaxis](#).

D - Lymphoma

Explanation Why

Decreased immunocompetence with increased susceptibility to infections and likelihood of [lymphoma](#) and leukemia are seen in patients with [ataxia-telangiectasia](#) (AT). This condition can also cause [telangiectasias](#) of the face, as seen in this patient, but these are usually on the [skin](#) or [conjunctivae](#) rather than the [lips](#) or mucous membranes of the nose (leading to [epistaxis](#)). In addition, progressive [cerebellar ataxia](#) is a hallmark symptom of AT that this patient does not have.

E - Glaucoma

Explanation Why

Early-onset [glaucoma](#) is seen in patients with [Sturge-Weber syndrome](#). These patients also present with vascular anomalies, but they take the form of a [port wine stain](#) (usually unilateral in the V1–V2 facial [dermatomes](#)) and vascular [malformations](#) of the [choroid](#) and [CNS](#) (e.g., [leptomeningeal angioma](#), usually ipsilateral to the [port wine stain](#)). The [CNS](#) vascular [malformations](#) lead to [epilepsy](#), [intellectual disability](#), and [glaucoma](#) due to increased intraorbital pressure. [Sturge-Weber](#) is not associated with [telangiectasias](#) or recurrent [epistaxis](#).

F - Renal cell carcinoma

Explanation Why

Patients with [von Hippel-Lindau](#) syndrome ([VHL](#)) have [hemangioblastomas](#) of the [retina](#) or [CNS](#). These patients also have renal and [pancreatic](#) cysts, [pheochromocytomas](#), and are at an increased risk for [renal cell carcinoma](#). This patient does not have any of the symptoms associated with these conditions. In addition, facial [telangiectasias](#) and [epistaxis](#) are not consistent with a diagnosis of [VHL](#).

Question # 27

A 59-year-old man with angina pectoris comes to the physician because of a 6-month history of shortness of breath on exertion that improves with rest. He has hypertension and hyperlipidemia. Current medications include aspirin, metoprolol, and nitroglycerine. Echocardiography shows left ventricular septal and apical hypokinesis. Cardiac catheterization shows 96% occlusion of the left anterior descending artery. The patient undergoes angioplasty and placement of a stent. The patient's shortness of breath subsequently resolves and follow-up echocardiography one week later shows normal regional contractile function. Which of the following is the most accurate explanation for the changes in echocardiography?

	Answer	Image
A	Unstable angina pectoris	
B	Stress cardiomyopathy	
C	Hibernating myocardium	
D	Myocardial scarring	
E	Cardiac remodeling	
F	Myocardial stunning	

Hint

The patient had signs and symptoms of chronic left ventricular dysfunction that were reversed by revascularization.

Correct Answer

A - Unstable angina pectoris

Explanation Why

[Unstable angina pectoris](#) is characterized by the sudden new onset of symptoms of [acute coronary syndrome](#) at rest or on minimal exertion, or rapidly worsening symptoms ([crescendo angina](#)). However, this patient has long-standing [dyspnea](#), which likely occurs on exertion and improves with rest and would therefore be most consistent with [stable angina pectoris](#).

B - Stress cardiomyopathy

Explanation Why

[Stress cardiomyopathy](#) ([Takotsubo cardiomyopathy](#)) describes an acute decline in left-ventricular function following an emotionally or physically stressful event (e.g., unexpected death of a relative, natural disaster, or respiratory failure, infection). Although [dyspnea](#) can be present, it would not be limited to exertion, as in this patient. A hallmark finding on [echocardiography](#) is apical ballooning of the [left ventricle](#), which is not seen here. In contrast to this patient who experienced rapid symptom relief and functional improvement after revascularization, patients with [stress cardiomyopathy](#) recover spontaneously and only require symptomatic treatment with [beta blockers](#) or ACE inhibitors.

C - Hibernating myocardium

Explanation Why

[Hibernating myocardium](#) is characterized by persistently impaired [myocardial contractility](#) at rest (on [echocardiography](#) or radionuclide myocardial perfusion imaging) that is partially or completely reversible when adequate oxygen supply is restored (e.g., after angioplasty or [coronary artery bypass surgery](#)). In response to decreased perfusion, hibernating [myocardial cells](#) enter a state of decreased metabolic activity. In contrast, [myocardial stunning](#) occurs after an episode of acute [ischemia](#) and is characterized by transiently impaired [myocardial contractility](#).

D - Myocardial scarring

Explanation Why

[Myocardial scar](#) tissue indicates [myocardial necrosis](#), most often induced by irreversible [myocardial ischemia](#) due to [myocardial infarction](#) (MI). The patient could have undergone a silent MI, but, since [myocardial](#) scars are marked by complete replacement with fibrous, non-contractile tissue, there would be permanent [dyskinesia](#) on [echocardiography](#) even after revascularization. [Scar](#) tissue can also lead to a true [ventricular aneurysm](#), which would then manifest as an outward bulge of the affected segment during [myocardial](#) contraction.

E - Cardiac remodeling

Explanation Why

Cardiac remodeling is a chronic process in which the [myocardium](#) undergoes permanent changes in structure (i.e., mass, shape, dimensions) and/or function to compensate for cardiac injury or an increased hemodynamic load that does not rapidly improve after treatment (here within one week). Reduced contractile function would likely persist even after successful revascularization.


F - Myocardial stunning

Explanation Why

In [myocardial stunning](#), affected segments undergo short but significant acute [ischemia](#) ("stunning"), but regain their baseline contractile functionality because of the short-term nature of the event. This patient's [myocardial](#) dysfunction is similar in that function is regained after revascularization. However, this patient's symptoms suggest a prolonged, chronic history of [myocardial](#) flow reduction and [ischemia](#).

Question # 28

A 36-year-old man comes to the physician for a follow-up examination. Two weeks ago, he was diagnosed with an ischemic stroke of the right middle cerebral artery. He was treated with thrombolytics and does not have any residual symptoms. His pulse is 82/min and regular. Cardiovascular examination shows no abnormalities. Echocardiography shows a reproducible, transient, low-volume, right-to-left shunt through the atrial septum during coughing. Which of the following conditions is caused by failure of an embryologic process similar to that responsible for this patient's heart condition?

	Answer	Image
A	Midgut volvulus	
B	Transposition of the great vessels	 a Interventricular foramen b Muscular interventricular septum c Aorticopulmonary septum d Membranous interventricular septum
C	Hirschsprung disease	
D	Syndactyly	
E	Thyroglossal duct cyst	
F	Hypospadias	

Hint

A transient right-to-left atrial shunt that occurs during coughing, which increases right atrial pressure, confirms that this patient has a patent foramen ovale (PFO). PFO is caused by a failure of the septum primum and septum secundum to fuse after birth.

Correct Answer


A - Midgut volvulus

Explanation Why

[Midgut volvulus](#) is caused by malrotation of the intestines during [fetal development](#). Malrotation is not the embryologic process responsible for [PFO](#).

B - Transposition of the great vessels

Image



Explanation Why

[Transposition of the great vessels](#) is caused by a failure of spiraling in the [aorticopulmonary septum](#) during [heart morphogenesis](#), resulting in a defect in which the aorta leaves the [right ventricle](#) and the [pulmonary trunk](#) leaves the left. Failure of spiraling is not the embryologic process responsible for

[PFO](#).

C - Hirschsprung disease

Explanation Why

[Hirschsprung disease](#) is caused by failure of [neural crest cell](#) migration during intestinal development. Failure of [neural crest cell](#) migration also plays a role in some conotruncal cardiac abnormalities, including [transposition of the great vessels](#), [tetralogy of Fallot](#), and [persistent truncus arteriosus](#). However, it is not the embryologic process responsible for [PFO](#).

D - Syndactyly

Explanation Why

[Syndactyly](#) is caused by a failure of [apoptosis](#), which normally destroys tissue between the digits during limb development, leaving them unfused and individually mobile. Failure of [apoptosis](#) is not the embryologic process responsible for [PFO](#).

E - Thyroglossal duct cyst

Explanation Why

[Thyroglossal duct cysts](#) are caused by failure of the [thyroglossal duct](#) to obliterate during [thyroid](#) development. Failure of obliteration is also the mechanism behind both [branchial cleft cyst](#), in which a [branchial cleft](#) persists, and [Meckel diverticulum](#), in which the [vitelline duct](#) persists. Failure of obliteration is not the embryologic process responsible for [PFO](#).

F - Hypospadias

Explanation Why

[Hypospadias](#) is caused by a failure of the urethral folds to fuse, in the same underlying embryologic process responsible for [PFO](#). Other defects caused by failure of fusion include [uterus bicornis](#), in which the [paramesonephric ducts](#) fail to fuse, as well as [cleft lip](#) and [cleft palate](#), in which the maxillary prominence fails to fuse with the [medial](#) nasal prominence. Most patients with [PFO](#) do not require treatment. However, sometimes, as was likely the case here, patients develop [paradoxical emboli](#) resulting in [ischemic stroke](#).

Question # 29

A 66-year-old man comes to the emergency department because of shortness of breath. His temperature is 37.2°C (99°F) and pulse is 105/min. When the blood pressure cuff is inflated to 140 mm Hg, the patient's pulse is audible and regular. However, upon inspiration, the pulse disappears and does not reappear until expiration. Only when the blood pressure cuff is inflated to 125 mm Hg is the pulse audible throughout the entire respiratory cycle. Which of the following underlying conditions is most likely responsible for this patient's physical examination findings?

	Answer	Image
A	Mitral regurgitation	
B	Lobar pneumonia	
C	Hypertrophic cardiomyopathy	
D	Congestive heart failure	
E	Asthma	

Hint

Arterial systolic pressure normally falls during inspiration, although the magnitude of decrease usually does not exceed 10 mm Hg. A pathological decrease in arterial systolic pressure > 10 mm Hg during inspiration (in this case, the difference is 15 mm Hg) is characteristic of pulsus paradoxus.

Correct Answer

A - Mitral regurgitation

Explanation Why

Chronic [mitral regurgitation](#) ([MR](#)) can manifest as [dyspnea](#) due to pulmonary venous congestion. [Physical examination](#) findings associated with [MR](#) include [holosystolic](#) high-pitched [murmur](#) heard best at the apex and [crackles](#) on [lung auscultation](#). [Pulsus paradoxus](#) is not associated with [mitral regurgitation](#).

B - Lobar pneumonia

Explanation Why

[Physical examination](#) findings associated with [lobar pneumonia](#) include dullness to percussion, [crackles](#) or [rales](#) on auscultation, [egophony](#), [bronchophony](#), whispering pectoriloquy, and [pleural friction rub](#). [Pulsus paradoxus](#) is not associated with [lobar pneumonia](#).

C - Hypertrophic cardiomyopathy

Explanation Why

[Physical examination](#) findings associated with [hypertrophic cardiomyopathy](#) ([HOCM](#)) include the presence of an S4, harsh [systolic](#) crescendo-decrescendo [murmur](#) heard at the apex, and a brisk carotid upstroke. Rarely, it is also associated with [reverse pulsus paradoxus](#), which refers to an inspiratory increase rather than a decrease in [systolic](#) and [diastolic](#) pressures. [Pulsus paradoxus](#), as seen in this patient, is not associated with [HOCM](#).

D - Congestive heart failure

Explanation Why

[Congestive heart failure](#) (CHF) does not cause [pulsus paradoxus](#), but is associated with [pulsus alternans](#) (alternating strong and weak pulses) on [physical exam](#) due to alterations in [cardiac output](#). Additional [physical examination](#) findings that may be seen with CHF include peripheral [edema](#), [crackles](#) on [lung auscultation](#), the presence of an [S3](#), and distended jugular [veins](#).


E - Asthma

Explanation Why

[Pulsus paradoxus](#) most commonly occurs in patients with constrictive cardiac disease (e.g., [cardiac tamponade](#), [constrictive pericarditis](#), [tension pneumothorax](#)) or severe [obstructive lung disease](#) (e.g., [COPD](#), [asthma](#)). During inspiration, there is increased systemic [venous return](#) and increased volume in the right [heart](#), which is compensated for by expansion into the [pericardial space](#) in a healthy [heart](#). However, in a pathologically constricted [heart](#), this increase in [right ventricle](#) volume causes bowing of the RV free wall into the [left ventricle](#), resulting in decreased LV filling, decreased [cardiac output](#), and subsequent drop in [systolic](#) blood pressure. [Pulsus paradoxus](#) in patients with [obstructive lung disease](#) (e.g., [asthma](#)), on the other hand, occurs due to the amplification of intrathoracic pressure variation during respiration, which is then transmitted to the [distal arteries](#).

Question # 30

A 57-year-old woman comes to the physician because of a 2-month history of intermittent dyspnea and dizziness. She has a history of mitral valve stenosis. Her pulse is 125/min and irregularly irregular, and blood pressure is 102/66 mm Hg. A transthoracic echocardiogram shows doming of the anterior mitral valve leaflet during systole. Which of the following elements is most likely to be absent from this patient's jugular venous pressure waveform?



	Answer	Image
A	Area 1	
B	Area 2	
C	Area 3	

	Answer	Image
D	Area 4	
E	Area 5	

Hint

This patient's intermittent symptoms and tachycardia with an irregularly irregular heartbeat are classic features of symptomatic atrial fibrillation with rapid ventricular response. Mitral stenosis is a risk factor for atrial fibrillation.

Correct Answer

A - Area 1

Explanation But

A prominent [a wave](#) may be seen in patients with [right ventricular hypertrophy](#) or [pulmonic stenosis](#) (due to increased pressure gradient across the right side of the [heart](#)).

Explanation Why

Area 1 corresponds to the [a wave](#) of the [JVP](#) waveform and corresponds with atrial contraction. Right atrial contraction leads to increased RA pressure and subsequent retrograde blood flow back to the jugular [vein](#), therefore manifesting with a peak in the [JVP](#) tracing. The [a wave](#) is absent in [atrial fibrillation](#), the most likely etiology of this patient's abnormal rhythm, due to a lack of unified contraction of the [right atrium](#) during fibrillation.

B - Area 2

Explanation Why

Area 2 corresponds to the [c wave](#) of the [JVP](#) waveform. It represents [right ventricular](#) contraction against a closed [tricuspid valve](#), which causes the valve to bulge into the [right atrium](#). This creates a second peak in the [JVP](#). Because [atrial fibrillation](#) does not affect [right ventricular](#) contraction or [tricuspid valve](#) function, this patient should have a normal [c wave](#).

C - Area 3

Explanation Why

Area 3 corresponds to the [x descent](#) of the [JVP](#) waveform. It represents atrial relaxation with continued ventricular contraction in which there is inferior displacement of the closed [tricuspid valve](#), resulting in a drop in [JVP](#). The [x descent](#) may be absent in [tricuspid regurgitation](#) in which

retrograde blood flow during ventricular [systole](#) prohibits a drop in right atrial pressure. This patient has [atrial fibrillation](#), but no known tricuspid disease. Although [atrial fibrillation](#) may reduce the degree of [x descent](#) secondary to impaired atrial relaxation, it should not be absent.

D - Area 4

Explanation Why

Area 4 corresponds to the [v wave](#) of the [JVP](#) waveform, which represents the increase in right atrial pressure during passive atrial filling against a closed [tricuspid valve](#). A reduced [v wave](#) may be seen in cases of decreased [preload](#) (e.g., [hypovolemia](#), venous [vasodilation](#)), while a prominent [v wave](#) can be seen in [tricuspid regurgitation](#) and [right heart failure](#). This patient's [v wave](#) should not be significantly changed.


E - Area 5

Explanation Why

Area 5 corresponds to the [y descent](#) of the [JVP](#) waveform. It represents [tricuspid valve](#) opening, with rapid blood flow from the [right atrium](#) into the [right ventricle](#), with a resulting decrease in right atrial pressure. A reduced [y descent](#) can be seen in cases of tricuspid stenosis and [cardiac tamponade](#). This patient's [y descent](#) should not be significantly changed.

Question # 31

A previously healthy 28-year-old man comes to the emergency department because of dizziness and palpitations for 2 days. Prior to the onset of the symptoms, he attended a bachelor party where he lost several drinking games. An ECG is shown. Which of the following is the most likely diagnosis?



	Answer	Image
A	Hypertrophic obstructive cardiomyopathy	
B	Dilated cardiomyopathy	
C	Paroxysmal atrial fibrillation	
D	Brugada syndrome	

	Answer	Image
E	Ventricular tachycardia	
F	Sick sinus syndrome	

Hint

The ECG shows an irregular rhythm with narrow QRS complexes and no discernable p-waves. Other common triggers of this patient's ECG findings include elevated sympathetic activity (e.g., during stress), hyperthyroidism, caffeine consumption, and mitral valve disease.

Correct Answer

A - Hypertrophic obstructive cardiomyopathy

Explanation Why

[Hypertrophic obstructive cardiomyopathy](#) is a hereditary disease of the [myocardium](#) that is characterized by asymmetric [hypertrophy](#) of the [left ventricle](#). Although it is usually asymptomatic and often an incidental finding, it can cause [syncope](#) and/or lethal [arrhythmias](#) that lead to [sudden cardiac death](#), typically during periods of physical exertion. In some cases, it can cause [palpitations](#) and [dizziness](#) as in this patient. However, [ECG](#) findings would show abnormally deep Q waves and signs of [left ventricular hypertrophy](#) (according to the [Sokolow-Lyon](#) Index).

B - Dilated cardiomyopathy

Explanation Why

[Dilated cardiomyopathy](#) is a unilateral or global enlargement of the [myocardium](#), which leads to limited contractility and a subsequent decrease in pumping function. Although the condition could explain this patient's [arrhythmia](#), [palpitations](#), and [dizziness](#), other specific signs of [right ventricular failure](#) (e.g. peripheral [pitting edema](#), signs of increased central venous pressure) or [left ventricular failure](#) (e.g. [pulmonary edema](#), [dyspnea](#)) are not present, making this condition very unlikely.

C - Paroxysmal atrial fibrillation

Explanation Why

[Atrial fibrillation](#) is a [supraventricular tachyarrhythmia](#) caused by uncoordinated atrial activation. Although usually asymptomatic, it can also cause symptoms such as this patient's [dizziness](#) and [palpitations](#). In particular, acute alcohol consumption often triggers paroxysmal A-fib (also known as “[holiday heart syndrome](#)”), which subsides within 7 days without intervention. As in this patient, an [ECG](#) shows irregular RR intervals, indiscernible [p waves](#), and [tachycardia](#) with narrow [QRS complexes](#). To prevent future episodes, this patient should abstain from alcohol in the future.

D - Brugada syndrome

Explanation Why

[Brugada syndrome](#) is a genetically inherited disorder that causes a [repolarization](#) abnormality and occurs more commonly in males of Asian descent. Although it is usually asymptomatic, it can cause [palpitations](#) and [dizziness](#), as in this patient, which may also be triggered by [substance abuse](#) or alcohol consumption. However, an [ECG](#) would show a persistent [ST elevation](#) in V1–V2 with intact [p waves](#).

E - Ventricular tachycardia

Explanation Why

[Ventricular tachycardia](#) is a potentially life-threatening [arrhythmia](#) originating in the ventricles and is associated with structural heart disease or [myocardial infarction](#). Although it can present with [palpitations](#), which may also be triggered by [substance abuse](#) or alcohol consumption, [ECG](#) findings would show AV dissociation and broad QRS complexes with a [heart rate](#) > 120 bpm.

F - Sick sinus syndrome



Explanation Why

[Sick sinus syndrome](#) is due to a dysfunctional [sinoatrial node](#) that can cause [bradyarrhythmias](#) and/or [supraventricular tachyarrhythmias](#). Although it can present with an irregular rhythm with narrow QRS complexes and no discernable [p-waves](#), [ECG](#) findings usually also show sinoatrial pauses or “[tachycardia-bradycardia](#)” cycling. In addition, [sick sinus syndrome](#) is associated with [coronary artery](#) or structural heart diseases. This patient's age, history of a single event, and [ECG](#) findings make [sick sinus syndrome](#) an unlikely explanation for his symptoms.

Question # 32

An 81-year-old man comes to the physician because of increased exertional dyspnea and dizziness over the past 8 weeks. He has hypertension for which he takes lisinopril. He has smoked one pack of cigarettes daily for the past 50 years. Physical examination shows weak peripheral pulses. Cardiac examination is shown. Which of the following is the most likely diagnosis?

[Video](#)

	Answer	Image
A	Pulmonary stenosis	
B	Mitral regurgitation	 The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The ECG shows a normal sinus rhythm with a narrow QRS complex. The PCG shows a high-pitched, early diastolic murmur that decays rapidly, characteristic of mitral regurgitation. The waveforms are synchronized with a timing diagram at the top indicating Diastole and Systole phases. Labels S1 and S2 are present at the bottom of the PCG trace.
C	Aortic stenosis	 The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The ECG shows a normal sinus rhythm with a narrow QRS complex. The PCG shows a high-pitched, early diastolic murmur that decays rapidly, characteristic of mitral regurgitation. The waveforms are synchronized with a timing diagram at the top indicating Diastole and Systole phases. Labels S1 and S2 are present at the bottom of the PCG trace.

	Answer	Image
D	Mitral valve prolapse	Diastole Systole Diastole Systole Diastole Electrocardiogram (ECG) Mid-systolic click Phonocardiogram S1 S2 S1 S2
E	Mitral stenosis	Diastole Systole Diastole Systole Diastole Electrocardiogram (ECG) OS Phonocardiogram S1 S2 S1 S2
F	Pulmonary regurgitation	
G	Tricuspid stenosis	
H	Aortic regurgitation	Diastole Systole Diastole Systole Diastole Electrocardiogram (ECG) Phonocardiogram S1 S2 S1 S2

	Answer	Image
I	Tricuspid regurgitation	

Hint

Auscultation shows a harsh crescendo-decrescendo, systolic ejection murmur best heard at the 2nd right intercostal space (ICS) that radiates to the carotids. These features are suggestive of the most common valvular heart disease in industrialized countries.

Correct Answer


A - Pulmonary stenosis

Explanation Why

[Pulmonary stenosis](#) is mostly a congenital valve defect that can also manifest with exertional dyspnea. In adults, common causes include [carcinoid syndrome](#) and chronic [rheumatic fever](#). But auscultation yields an ejection [systolic murmur](#) best heard in the left 2nd ICS.

B - Mitral regurgitation

Image




Explanation Why

Chronic [mitral regurgitation](#) ([MR](#)) leads to pulmonary venous congestion, which, in turn, causes

[dyspnea](#), as seen in this patient. However, patients with [MR](#) would have normal peripheral pulses and a [holosystolic](#) high-pitched [murmur](#) heard best at the left [midclavicular line](#) of the 5th ICS that radiates to the left [axilla](#).

C - Aortic stenosis

Image




Explanation Why

This patient's auscultation findings are consistent with [aortic stenosis](#) (AS), which causes [left ventricular](#) outflow obstruction. The result is [dizziness](#) and exacerbation of [dyspnea](#) upon exertion, and weak, delayed peripheral pulses ([pulsus parvus et tardus](#)). Incomplete [left ventricular](#) emptying may also cause pulmonary venous congestion, which further exacerbates [dyspnea](#).

D - Mitral valve prolapse

Image




Explanation Why

[Mitral valve prolapse \(MVP\)](#) is typically asymptomatic but can also present with exertional dyspnea. However, auscultation of an [MVP](#) shows a high-frequency, midsystolic click with a late [systolic murmur](#) that is heard best at the left [midclavicular line](#) of the 5th ICS.

E - Mitral stenosis

Image



Explanation Why

[Mitral stenosis](#) (MS) leads to pulmonary venous congestion, which, in turn, causes [dyspnea](#), as seen in this patient. However, auscultation in MS yields a mid-[diastolic murmur](#) heard best at the midclavicular 5th left ICS (the apex), and is often preceded by an [opening snap](#).

F - Pulmonary regurgitation

Explanation Why

Pulmonary regurgitation (PR) is usually asymptomatic but can also manifest as exertional dyspnea. However, [features of right-sided heart failure](#) would be present, and auscultation in PR would show an early [diastolic](#) decrescendo [murmur](#) that is best heard parasternally in the left 2nd ICS.


G - Tricuspid stenosis

Explanation Why

Tricuspid stenosis (TS) is extremely rare and typically presents with abdominal discomfort (due to [hepatomegaly](#)) and other [features of right-sided heart failure](#), which are not present here. In addition, the typical [murmur](#) of TS is a soft mid-[diastolic murmur](#) heard best in the parasternal region of the left 4th ICS.

H - Aortic regurgitation

Image



Explanation Why

Patients with severe [aortic regurgitation](#) can present with [dizziness](#) and exertional dyspnea, like this patient. However, examination in AR would show a high [pulse pressure](#), a [water hammer pulse](#), and

a high-pitched early [diastolic murmur](#) heard best at the 2nd right ICS that does not radiate to the carotid [arteries](#).

I - Tricuspid regurgitation

Explanation Why

[Tricuspid regurgitation](#) (TR) is extremely rare and in severe cases presents with abdominal discomfort (due to [hepatomegaly](#)) and other [features of right-sided heart failure](#), which are not present here. In addition, the [murmur](#) of TR is [holosystolic](#) and heard best in the parasternal region of the left 4th ICS.

Question # 33

A 43-year-old man comes to the physician because of anxiety, difficulty focusing on tasks, and a 4.6-kg (10-lb) weight loss over the past 4 weeks. He is diaphoretic. His pulse is 100/min, respirations are 18/min, and blood pressure is 150/78 mm Hg. Physical examination shows warm, moist skin, goiter, and a resting tremor of both hands. Laboratory studies show a thyroxine (T4) concentration of 30 µg/dL and a thyroid-stimulating hormone concentration of 0.1 µU/mL. The patient is started on methimazole and atenolol. The latter agent predominantly affects which of the following?

	Answer	Image
A	Atrioventricular node activity	Class II
B	Vagal tone	
C	Effective refractory period of the cardiac action potential	
D	His-Purkinje conduction	

	Answer	Image
E	Phase 0 depolarization slope of the cardiac action potential	


Hint

Atenolol is a cardioselective beta blocker, which competitively antagonizes the positive chronotropic, inotropic, and dromotropic effect of catecholamines.

Correct Answer

A - Atrioventricular node activity

Image



Explanation Why

[\$\beta_1\$ adrenergic receptor](#) antagonists lower concentrations of [cAMP](#) and calcium in the cells of the [atrioventricular node](#). This decreases conduction and [automaticity](#) at the [atrioventricular node](#) by decreasing the slope of phase 4 [depolarization](#) of the nodal (pacemaker) [action potential](#). On [ECG](#) this effect would appear as a prolonged [PR interval](#). [Beta blockers](#) also have a negative [chronotropic](#) ($\downarrow$ risk of [heart failure](#)) and a negative [inotropic](#) ($\downarrow$ contractility) effect.


B - Vagal tone

Explanation Why

[Atenolol](#) acts on β_1 receptors to decrease [sympathetic](#) tone. It has no effect on [vagal](#) ([parasympathetic](#)) tone. [Digoxin](#) is an antiarrhythmic that can increase [vagal](#) tone.

C - Effective refractory period of the cardiac action potential

Image



Explanation Why

Prolongation of the [effective refractory period](#) of the [cardiac action potential](#) is caused by Class IA, IC, III, and IV [antiarrhythmic drugs](#). [Beta-blockers](#) ([class II antiarrhythmics](#)), however, do not significantly alter the [refractory period](#).

D - His-Purkinje conduction

Explanation Why

Class I [antiarrhythmics](#) can decrease conduction in extranodal tissue (e.g., His-Purkinje system). [Beta blockers](#) do not act on the His-Purkinje system.

E - Phase 0 depolarization slope of the cardiac action potential

Explanation Why


Phase 0 [depolarization](#) slope of the [cardiac action potential](#) is caused by the opening of fast [voltage-gated Na⁺ channels](#). Class I antiarrhythmics, which block these channels, would, therefore, affect phase 0. [Beta-blockers](#) ([class II antiarrhythmics](#)) do not affect Na⁺ channel activity and do not have any effect on phase 0. They instead decrease the slope of phase 4 [depolarization](#).

Question # 34

A 64-year-old man who recently immigrated to the United States from Haiti comes to the physician because of a 3-week history of progressively worsening exertional dyspnea and fatigue. For the past few days, he has also had difficulty lying flat due to trouble breathing. Over the past year, he has had intermittent fever, night sweats, and cough but he has not been seen by a physician for evaluation of these symptoms. His temperature is 37.8°C (100°F). An x-ray of the chest is shown. Further evaluation of this patient is most likely to show which of the following findings?



	Answer	Image
A	Elimination of S2 heart sound splitting with inspiration	
B	Head bobbing in synchrony with heart beat	

	Answer	Image
C	Painless erythematous lesions on palms or soles	
D	Prominent "a" wave on jugular venous pressure tracing	
E	Jugular venous distention on inspiration	
F	Crescendo-decrescendo systolic ejection murmur	

Hint

New-onset dyspnea, orthopnea, and fatigue and pericardial calcifications on x-ray indicate constrictive pericarditis. This patient who hails from Haiti and has a prolonged history of night sweats, fever, cough most likely has tuberculosis, which can cause constrictive pericarditis.

Correct Answer

A - Elimination of S2 heart sound splitting with inspiration

Explanation Why

[Splitting of the second heart sound](#) (S₂) is a physiological finding heard during inspiration. It is caused by the early closure of the [aortic valve](#) (A₂) in relation to the [pulmonary valve](#) (P₂) when intrathoracic pressure is decreased. In conditions such as [aortic stenosis](#), [VSD](#) with [Eisenmenger syndrome](#), and [LBBB](#), elimination of S₂ splitting can occur because A₂ is delayed and superimposed on P₂. Although severe, long-standing [aortic stenosis](#) can result in fatigue, [dyspnea](#), orthopnea, and [cough](#) (due to cardiogenic pulmonary edema), chronic AS is not associated with [fever](#) or night sweats but with symptoms such as [syncope](#), [palpitations](#), and [angina](#). Moreover, an [x-ray](#) would not show [pericardial](#) calcifications but an enlarged [cardiac silhouette](#).

B - Head bobbing in synchrony with heart beat

Explanation Why

Rhythmic nodding or bobbing of the head in synchronicity with the heartbeat is referred to as [de Musset sign](#), a typical feature of chronic [aortic regurgitation](#) (AR). AR may present with progressively worsening fatigue, [dyspnea](#), orthopnea, and [cough](#) (due to cardiogenic pulmonary edema). However, chronic AR is not associated with [fever](#) or night sweats, and an [x-ray](#) would not show [pericardial](#) calcifications, but rather a prominent aortic root and an enlarged [cardiac silhouette](#).

C - Painless erythematous lesions on palms or soles

Image



Explanation Why

Painless [erythematous](#) lesions on the palms or soles are called [Janeway lesions](#), a finding highly suggestive of [infective endocarditis](#), which can present with [dyspnea](#), fatigue, [cough](#), [fever](#), and night sweats. However, the symptoms of [infective endocarditis](#) have an acute or subacute onset (days to weeks), unlike this patient who had [fever](#), night sweats, and [cough](#) for a year before developing [dyspnea](#) and fatigue. Moreover, an [x-ray](#) would not show [pericardial](#) calcifications, and [infective endocarditis](#) is almost always associated with the development of a new [heart murmur](#).

D - Prominent "a" wave on jugular venous pressure tracing

Explanation Why

On a [jugular venous pressure](#) tracing, the "a" wave is caused by atrial contraction in late [diastole](#). A

prominent “a” wave is a sign of increased atrial contraction pressure due to restricted filling of the [right ventricle](#) (e.g., tricuspid valve stenosis, [right ventricular hypertrophy](#), [cardiac tamponade](#)). In [constrictive pericarditis](#), [right ventricular](#) filling is only impaired when the elastic limit of the [pericardium](#) is reached. As a result, all [right ventricular](#) filling rapidly occurs in early [diastole](#), producing a prominent and deep [x descent](#) and [y descent](#) on venous pressure tracing rather than a prominent “a” wave.

E - Jugular venous distention on inspiration

Explanation But

In [cardiac tamponade](#) without underlying [pericarditis](#) (e.g., traumatic [hemopericardium](#)), the [jugular venous pressure](#) is increased as a result of impaired [right ventricular](#) filling. However, the [right ventricle](#) is still compliant and a decrease in intrathoracic pressure can be transmitted to the [right ventricle](#) across the fluid-filled [pericardial sac](#). Therefore, [Kussmaul sign](#) would not be expected in traumatic [cardiac tamponade](#).

Explanation Why

Normally, the decrease in intrathoracic pressure during inspiration expands the compliant [right ventricle](#) resulting in a decrease in [jugular venous pressure](#). In [constrictive pericarditis](#), compliance of the [right ventricle](#) is decreased and the [right ventricle](#) fails to expand during inspiration. Therefore, the [jugular venous pressure](#), which might already be increased in [constrictive pericarditis](#) due to restricted [right ventricular](#) filling during late [diastole](#), would increase further during inspiration. This phenomenon, which is called [Kussmaul sign](#), can also be seen in patients with [restrictive cardiomyopathy](#) and right atrial or ventricular tumors.

F - Crescendo-decrescendo systolic ejection murmur

Image





Explanation Why

A crescendo-decrescendo (diamond-shaped) [systolic ejection murmur](#) best heard in the second right [intercostal space](#) is a common finding in [aortic valve stenosis](#). Although severe, long-standing [aortic stenosis](#) can result in [chest pain](#), fatigue, [dyspnea](#), orthopnea, and [cough](#) (due to cardiogenic pulmonary edema), chronic AS is not associated with [fever](#) or night sweats but with symptoms such as [syncope](#), [angina](#), and [palpitations](#). Moreover, an [x-ray](#) would not show [pericardial](#) calcifications but an enlarged [cardiac silhouette](#).

Question # 35

A 38-year-old woman comes to the physician because of an 8-week history of shortness of breath and dull chest pain. She has a history of antiphospholipid syndrome. Physical examination shows jugular venous distention. Right heart catheterization shows a mean pulmonary arterial pressure of 30 mm Hg and a pulmonary capillary wedge pressure of 10 mm Hg. Further evaluation is most likely to show which of the following?

	Answer	Image
A	Constriction of the renal afferent arteriole	
B	Dilation of the coronary sinus	
C	Decreased left ventricular contractility	
D	Mitral valve leaflet thickening	

	Answer	Image
E	Hemosiderin-laden macrophages	

Hint

The presence of jugular venous distention suggests right heart failure, which can present with findings such as exertional shortness of breath and chest pain, as seen here. Antiphospholipid syndrome can cause pulmonary embolism, resulting in pulmonary hypertension and right heart failure. A mean pulmonary artery pressure ≥ 25 mm Hg at rest confirms the diagnosis of pulmonary hypertension, and a normal PCWP (4–12 mm Hg) excludes left heart failure as the cause.

Correct Answer


A - Constriction of the renal afferent arteriole

Explanation Why

In cases of systemic hypoperfusion, reduced renal perfusion results in dilatation of the [afferent arteriole](#) via [myogenic autoregulation](#) and the [tubuloglomerular feedback](#) mechanism, as well as constriction of the [efferent arteriole](#) via an increase in [angiotensin II](#) synthesis. [Afferent arteriole](#) dilation and [efferent arteriole](#) constriction then work in combination to maintain a constant [GFR](#) despite decreased [cardiac output](#). [Right heart failure](#) is a condition that can cause systemic hypoperfusion. For this reason dilatation (not constriction) of the renal [afferent arteriole](#) would rather be expected in this patient.

B - Dilation of the coronary sinus

Image



Explanation Why

[Right heart failure](#) due to [pulmonary hypertension](#) involves increased [afterload](#) of the [right ventricle](#), resulting in reduced [cardiac output](#) on the right side of the [heart](#). Reduced right-sided [cardiac output](#) leads to venous congestion in the vessels that drain into the right [heart](#), including the [coronary sinus](#), which can become dilated as a result.

C - Decreased left ventricular contractility

Explanation Why

Decreased [left ventricular contractility](#) is the hallmark of [systolic left heart failure](#). Although [left heart failure](#) can lead to [right heart failure](#), [PCWP](#) is typically elevated in [left heart failure](#) (> 18 mm Hg). This patient's normal [PCWP](#) makes [left heart failure](#) unlikely.


D - Mitral valve leaflet thickening

Explanation Why

[Mitral valve](#) leaflet thickening can occur in [mitral valve](#) diseases such as [mitral stenosis](#) (e.g., due to [rheumatic fever](#)). [Mitral valve](#) disease usually results in increased left atrial pressure and therefore increased [PCWP](#), whereas [PCWP](#) is normal in this patient.

E - Hemosiderin-laden macrophages

Image



Explanation Why

[Hemosiderin](#)-laden [macrophages](#) ([heart failure cells](#)) are typically seen in the alveoli of patients with chronic pulmonary congestion due to blood leakage into the alveolar spaces. Conditions that cause chronic pulmonary venous congestion include [left heart failure](#) and [mitral stenosis](#). [PCWP](#) is typically elevated in [left heart failure](#) and [mitral stenosis](#), whereas it is normal in this patient.

Question # 36

A 61-year-old man comes to the physician with several months of sharp, shooting pain in both legs. Twenty years ago, he had a painless ulcer on his penis that resolved without treatment. He has no history of serious illness. Examination shows small pupils that constrict with accommodation but do not react to light. Sensation to pinprick and light touch is decreased over the distal lower extremities. Patellar reflexes are absent bilaterally. His gait is unsteady and broad-based. This patient is at increased risk for which of the following complications?

	Answer	Image
A	Atrioventricular block	
B	Mitral valve regurgitation	
C	Penile squamous cell carcinoma	
D	Cerebral artery septic embolism	
E	Inflammatory polyarthritis	
F	Thoracic aortic aneurysm	

Hint

This patient has evidence of tabes dorsalis (ataxia, lower extremity pain, impaired sensation in the lower extremities) and Argyll Robertson pupil (impaired pupillary constriction), suggesting tertiary syphilis.

Correct Answer

A - Atrioventricular block

Explanation Why

Development of [atrioventricular block](#) can be caused by various etiologies (e.g., infection, [cardiomyopathy](#), [ischemia](#), etc); the most common infectious causes include [Lyme disease](#) and [infective endocarditis](#) (IE). This patient's [syphilis](#) infection would not increase his risk of developing [atrioventricular block](#).

B - Mitral valve regurgitation

Explanation Why

[Mitral valve regurgitation](#) is a potential complication of [rheumatic fever](#) (RF) or [rheumatic heart disease](#) (due to untreated RF). RF can also present with neurological sequelae (e.g., [Sydenham chorea](#)); however, [paresthesia](#) and pupillary involvement would not be expected. This patient's [syphilis](#) infection would not increase his risk of developing [mitral valve regurgitation](#).

C - Penile squamous cell carcinoma

Explanation Why

Penile [squamous cell carcinoma](#) is associated with infection from high-risk strains of [human papillomavirus](#) (i.e., [HPV-16](#), [HPV-18](#)). Anogenital lesions caused by [HPV](#) are usually raised, cauliflower-like, and exophytic, rather than the painless ulcers reported by this patient. This patient's [syphilis](#) infection would not increase his risk of developing [squamous cell carcinoma](#).

D - Cerebral artery septic embolism

Explanation Why

Cerebral [artery septic embolism](#) is almost exclusively seen in left-sided [infective endocarditis](#), especially with [S. aureus](#) and [Streptococcus bovis](#) infection, and can lead to the development of [mycotic aneurysm](#). This patient has evidence of [syphilis](#) infection, which is not known to cause embolization.

E - Inflammatory polyarthrititis

Explanation But

Patients with [DGI](#) may also present with a rash, but it is typically characterized by disseminated vesicular [dermatitis](#) (sometimes as part of an [arthritis-dermatitis syndrome](#)) that occurs concurrently with other symptoms.


Explanation Why

Inflammatory polyarthrititis can occur in patients with [disseminated gonococcal infection \(DGI\)](#) or [reactive arthritis](#), but not [syphilis](#) infection. While synovitis can develop in patients with [syphilis](#), it is more typically seen in [secondary syphilis](#), and would classically present within 10 weeks of primary infection; this patient's infection has been present for much longer.

F - Thoracic aortic aneurysm

Explanation Why

In addition to [tabes dorsalis](#) and [Argyll Robertson pupil](#), patients with [tertiary syphilis](#) are at risk of developing [thoracic aortic aneurysm \(TAA\)](#) from [aortitis](#). Treponemal-mediated destruction of the [vasa vasorum](#) classically affects the [ascending aorta](#) and can also lead to [aortic root dilatation](#) and [aortic regurgitation](#). These cardiovascular complications are late manifestations of [syphilis](#), and typically only occur after 15 years of untreated infection.



Question # 37

A 52-year-old man with a history of hypertension and hyperlipidemia comes to the physician because of a 10-month history of substernal chest pain on exertion that is relieved with rest. His pulse is 82/min and blood pressure is 145/82 mm Hg. He is prescribed a drug that acts by forming free radical nitric oxide. The patient is most likely to experience which of the following adverse effects as a result of this drug?

	Answer	Image
A	Pulsating headaches	
B	Digital vasospasm	
C	Hypertensive urgency	
D	Nonproductive cough	
E	Lupus-like syndrome	
F	Lower extremity edema	
G	Erectile dysfunction	

Hint

Nitrates work by forming free radical nitric oxide, which leads to smooth muscle relaxation and vasodilation.

Correct Answer

A - Pulsating headaches

Explanation But

Other adverse effects include [gastroesophageal reflux](#) and [reflex tachycardia](#), the latter of which can be treated with [beta blockers](#).

Explanation Why

[Headaches](#) are a common side effect of [nitrates](#) and affect approx. 10% of patients. These [headaches](#) are either mild-to-moderate and occur immediately after drug administration (due to [nitric oxide](#)-induced [vasodilation](#)) or [migraine](#)-like, occurring several hours after drug intake (due to [nitric oxide](#)-mediated release of [neurotransmitters](#) such as [calcitonin gene-related peptide](#) and [glutamate](#), and/or altered ion channel function). [Nitrate](#)-induced [vasodilation](#) decreases both [preload](#) and [afterload](#) and improves [myocardial](#) perfusion, but may also result in cutaneous flushing and [hypotension](#).

B - Digital vasospasm

Explanation Why

Digital vasospasm (secondary [Raynaud phenomenon](#)) is a potential side effect of [beta blockers](#). This side effect is mediated by the blockade of [beta blockers](#) on [β₂-receptors](#), which causes vasoconstriction. [Nitrates](#) do not cause digital vasospasm.

C - Hypertensive urgency

Explanation Why

[Nitrates](#) can cause [hypotension](#), rather than [hypertensive urgency](#). [Nitrates](#) are used to treat [hypertensive crisis](#) for short-term reduction of blood pressure and are contraindicated in hypotensive

patients.

D - Nonproductive cough

Explanation Why

Nonproductive [cough](#) is a side effect of ACE inhibitors. This side effect is due to decreased breakdown of [bradykinin](#), which increases production of [arachidonic acid](#) metabolites (e.g., [prostaglandins](#)). The accumulation of [prostaglandins](#) in the upper respiratory tract and the [lungs](#) results in nonproductive [cough](#). [Nitrates](#) do not cause [cough](#).

E - Lupus-like syndrome

Explanation Why

[Lupus](#)-like syndrome ([drug-induced lupus erythematosus/DILE](#)) is a side effect of [antihypertensives](#), such as [hydralazine](#) and [methyldopa](#). Other drugs that can lead to [DILE](#) include monoclonal [TNF- \$\alpha\$ inhibitors](#), such as [infliximab](#) or [etanercept](#), [procainamide](#), [penicillamine](#), [isoniazid](#), [minocycline](#), and [phenytoin](#). [Nitrates](#) do not cause [DILE](#).

F - Lower extremity edema

Explanation Why

Lower extremity [edema](#) is a common side effect of dihydropyridine [calcium channel blockers](#) ([CCBs](#)), particularly [amlodipine](#). [CCBs](#) cause [vasodilation](#), which increases the [hydrostatic pressure](#) within the blood vessels, causing extravasation of plasma into the [interstitial](#) space. Although [nitrates](#) can rarely cause lower extremity [edema](#), it is not the most common side effect associated with this drug class.

G - Erectile dysfunction

Explanation Why




[Erectile dysfunction](#) is a known side effect of [beta blockers](#), which effect [β2-receptors](#) in the vasculature. Patients taking [PDE-5 inhibitors](#) for [erectile dysfunction](#) have an increased risk of life-threatening [hypotension](#) if [nitrates](#) are taken within 24 hours of a [PDE-5 inhibitor](#). [Nitrates](#) do not cause [erectile dysfunction](#).









Question # 38


A 3-year-old boy is brought to the physician for a follow-up examination after having undergone extensive workup of a heart murmur that was detected on routine examination. Results of hemodynamic studies performed during cardiac catheterization show:

	Oxygen saturation (%)	Pressure (mm Hg)
Right atrium	74	6
Right ventricle	85	40/10
Truncus pulmonalis	85	35/20
Left atrium	98	10
Left ventricle	98	110/10
Ascending aorta	98	105/65

This patient's murmur is most likely caused by which of the following conditions?

	Answer	Image
A	Ventricular septal defect	1  2  3 

	Answer	Image
B	Atrial septal defect	1  2  3 
C	Coarctation of the aorta	1  2 
D	Patent ductus arteriosus	1  2  3 

	Answer	Image
E	Transposition of the great vessels	


Hint

This patient has an increased oxygen saturation and increased blood pressure in the right ventricle and truncus pulmonalis.

Correct Answer

A - Ventricular septal defect

Image



Explanation But


In larger [heart](#) defects, long-term complications such as [endocarditis](#), [arrhythmias](#), [aortic regurgitation](#), [right heart failure](#), and [Eisenmenger reaction](#) are possible.

Explanation Why

This patient's hemodynamic studies indicate a [ventricular septal defect \(VSD\)](#). Due to a [left-to-right shunt](#), oxygen-rich blood flows from the [left ventricle](#) into the [right ventricle](#) during [systole](#). This causes increased [oxygen saturation](#) and increased blood pressure in the [right ventricle](#) and [truncus pulmonalis](#). Since the shunt is at the level of the ventricles, the [right atrium](#) maintains normal oxygenation and pressure, as seen here. The classic [murmur](#) is a harsh, [holosystolic murmur](#) that is loudest at the left lower sternal border; it is typically louder with smaller defects. Patients are often asymptomatic and spontaneous closure of small [VSDs](#) frequently occurs. Surgical interventions are rarely necessary.

B - Atrial septal defect

Image




Explanation Why

[Atrial septal defects](#) (ASD) result in blood flow from the [left atrium](#) to the [right atrium](#). ASDs typically manifest with [systolic ejection murmurs](#) over the left second [intercostal space](#) and are usually asymptomatic during childhood and [adolescence](#). Although an ASD would also lead to increased blood pressure in the right [heart](#), the [right atrium](#) would be affected in addition to the [right ventricle](#) and [truncus pulmonalis](#). Additionally, the [oxygen saturation](#) would be increased in the [right atrium](#), [right ventricle](#), and truncus, while the [oxygen saturation](#) in this patient's [right atrium](#) was normal.

C - Coarctation of the aorta

Image




Explanation Why

Coarctation of the aorta results in higher resistance to left ventricular outflow. Patients with small defects can also be asymptomatic and present with a heart murmur in the left paravertebral region. However, the aortic outflow obstruction would increase the pressure in the left ventricle rather than in the right ventricle and truncus pulmonalis. Additionally, oxygen saturation throughout the heart would be normal, whereas this patient has an increased oxygen saturation in the right ventricle and truncus pulmonalis.

D - Patent ductus arteriosus

Image




Explanation Why

In a [patent ductus arteriosus \(PDA\)](#), blood shunts from the aorta to the [pulmonary arteries](#). Patients with [PDA](#) can also be asymptomatic and present with a [heart murmur](#), which is typically continuous (“[machinery murmur](#)”) and heard best over the left infraclavicular region. A [PDA](#) would also lead to increased [oxygen saturation](#) in the [truncus pulmonalis](#) because of the direct connection with the aorta. However, [oxygen saturation](#) of the [right ventricle](#) would be normal rather than increased. Moreover, the pressure in the [truncus pulmonalis](#) would be increased more than the [right ventricular](#) pressure, which is not the case here.

E - Transposition of the great vessels

Image



Explanation Why

Transposition of the great vessels (TGV) is caused by an anatomical reversal of the aorta and the pulmonary artery, which connects the right ventricle to the systemic circulation and the left ventricle to the pulmonary circulation. While the oxygen saturation in the left atrium, left ventricle, and truncus pulmonalis would be high, the aorta and entire right heart would be low in oxygen. This condition is typically evident at birth and requires immediate surgical repair; TGV is an unlikely diagnosis in an asymptomatic 3-year-old boy. Additionally, pressures would generally be higher in the right heart than in the left heart because of the connection to the systemic circulation.

Question # 39

A 73-year-old African American man comes to the physician for a routine health maintenance examination. He has type 2 diabetes mellitus, hypertension, hypercholesterolemia, and coronary artery disease. His current medications include metformin, hydrochlorothiazide, amlodipine, rosuvastatin, isosorbide mononitrate, aspirin, and dipyridamole. He is 180 cm (5 ft 11 in) tall and weighs 110 kg (242 lb); BMI is 34 kg/m^2 . His vital signs are within normal limits. His hemoglobin A_{1c} concentration is 6.7%. Serum lipid studies show:

Cholesterol, total	302 mg/L
HDL-cholesterol	39 mg/dL
LDL-cholesterol	197 mg/dL
Triglycerides	292 mg/dL

The physician prescribes a drug that inhibits intestinal cholesterol absorption. The addition of this drug is most likely to increase the risk of which of the following adverse effects?

	Answer	Image
A	Hepatotoxicity	
B	Hyperkalemia	
C	Hyperglycemia	
D	Cutaneous flushing	
E	Hyperuricemia	

	Answer	Image
F	Cholelithiasis	

Hint

The lipid-lowering drug that inhibits intestinal cholesterol absorption is ezetimibe.

Correct Answer

A - Hepatotoxicity

Explanation Why

[Ezetimibe](#) is a [second-line lipid-lowering agent](#) used either as monotherapy (in cases of intolerance or contraindication to [statins](#)) or in combination with a [statin](#) in cases of persistent [hypercholesterolemia](#), as seen here. Hepatotoxicity (typically manifesting with elevated [liver](#) enzymes) is a side effect of [ezetimibe](#), and a higher [incidence](#) of hepatotoxicity is seen in patients on concomitant [statin](#) therapy. Hepatotoxicity is usually self-limiting and rarely causes clinically apparent [liver](#) injury. Other adverse effects of [ezetimibe](#) include myalgias, abdominal [pain](#), flatulence, and [diarrhea](#).

B - Hyperkalemia

Explanation Why

Neither [ezetimibe](#) nor any of the other lipid-lowering agents are known to cause [hyperkalemia](#). Patients with multiple [risk factors for cardiovascular disease](#) often take polypharmacy medication involving antidiabetic, antihyperlipidemic, and antihypertensive agents. [Hyperkalemia](#) is an adverse effect of ACE-inhibitors, angiotensin-receptor blockers, [direct renin inhibitors](#), and [SGLT-2 inhibitors](#). This patient's medication regimen does not include any of these drugs.

C - Hyperglycemia

Explanation Why

[Hyperglycemia](#) is an adverse effect of [niacin](#), another [second-line lipid-lowering agent](#). This patient is at an increased risk of [hyperglycemia](#) because he takes [hydrochlorothiazide](#). However, [ezetimibe](#) is not known to cause this adverse effect.

D - Cutaneous flushing

Explanation Why

Cutaneous flushing is an adverse effect of [niacin](#), another [second-line lipid-lowering agent](#). Taking [NSAIDs](#) 30–60 minutes prior to [niacin](#) can minimize this [prostaglandin](#)-associated effect. This patient might experience flushing as an adverse effect of several of his medications ([amlodipine](#), [isosorbide mononitrate](#), and [dipyridamole](#)). However, [ezetimibe](#) is not known to cause this adverse effect.

E - Hyperuricemia

Explanation Why

[Hyperuricemia](#) and an increased risk of [gout](#) are adverse effects of [niacin](#), another [second-line lipid-lowering agent](#). This patient is at an increased risk of [hyperuricemia](#) because he takes [hydrochlorothiazide](#). However, [ezetimibe](#) is not known to cause this adverse effect.

F - Cholelithiasis


Explanation Why

[Cholelithiasis](#) is an adverse effect of [fibrates](#), another group of [second-line lipid-lowering agents](#). For this reason, [fibrates](#) are contraindicated in patients with [gall bladder](#) disease. However, [ezetimibe](#) is not known to cause this adverse effect.

Question # 40

A 55-year-old man is brought to the emergency department 12 hours after the sudden onset of shortness of breath and substernal chest pain at rest; the pain is increased by inspiration. He has also had a nonproductive cough, fever, and malaise for the past 5 days. He does not smoke or use illicit drugs. His temperature is 38°C (100.4°F), pulse is 125/min, respirations are 32/min, and blood pressure is 85/45 mm Hg. Physical examination shows distended neck veins. Auscultation of the chest discloses bilateral basilar rales and muffled heart sounds. An ECG shows sinus tachycardia, diffuse ST segment elevation, low voltage QRS complexes, and fluctuating R wave amplitude. Which of the following is the most likely diagnosis?

	Answer	Image
A	Post-infarction fibrinous pericarditis	
B	Kawasaki disease	
C	Rheumatic fever	
D	Dressler syndrome	
E	Infective endocarditis	

	Answer	Image
F	Cardiac tamponade	 The image shows a 12-lead ECG tracing on a standard grid. The leads are arranged in four rows: Row 1 (I, aVR, V1), Row 2 (II, aVL, V2), Row 3 (III, aVF, V3), and Row 4 (aVF, V4, V5). The tracing exhibits electrical alternans, where the amplitude of the QRS complexes alternates between high and low in a regular fashion across all leads. This is a classic ECG finding in cardiac tamponade.

Hint

A 12-hour history of dyspnea and inspiratory chest pain in the setting of a 5-day prodrome of subjective fever, malaise, and dry cough suggests an acute infection of the pericardium.

Correct Answer

A - Post-infarction fibrinous pericarditis

Explanation Why

[Post-infarction fibrinous pericarditis](#) typically occurs 1–3 days after a transmural [myocardial infarction](#) (MI) and can manifest with [fever](#), inspiratory [chest pain](#), dry [cough](#), and diffuse [ST elevation](#), as seen in this patient. However, this patient does not have a history of [coronary artery disease](#) or recent [MI](#), making [post-infarction fibrinous pericarditis](#) unlikely. Rather, his 5-day history of [cough](#), [fever](#), and malaise suggests an infectious cause for the [pericarditis](#).

B - Kawasaki disease

Explanation Why

[Kawasaki disease](#) manifests with [fever](#) and malaise, as seen in this patient. [Myocarditis](#) and [myocardial infarction](#) are known complications of this condition that can manifest with [chest pain](#) and features of cardiac decompensation (e.g., [dyspnea](#), bilateral [rales](#), distended neck [veins](#)). However, a diagnosis of [Kawasaki disease](#) requires further specific symptoms, such as [erythema](#) and [edema](#) of the hands and feet, polymorphous rash, [conjunctivitis](#), cervical [lymphadenopathy](#), and oropharyngeal mucositis. Moreover, this condition most commonly affects children under the age of 5 years, so it is an unlikely diagnosis in a 55-year-old man.

C - Rheumatic fever

Explanation Why

[Rheumatic fever](#) (RF) manifests with [fever](#) and malaise, as seen in this patient. Cardiac complications of RF include pancarditis and [valvular heart disease](#), which can cause symptoms of cardiac decompensation (e.g., [dyspnea](#), bilateral [rales](#), distended neck [veins](#)). However, the typical [ECG](#) finding in RF is a prolonged [PR interval](#), which is not seen here. Moreover, this patient does not have a history of untreated [bacterial tonsillopharyngitis](#) or other typical features of RF (e.g., migratory polyarthritis, subcutaneous nodules, [erythema marginatum](#), [Sydenham chorea](#)).

D - Dressler syndrome

Explanation Why

[Dressler syndrome](#) usually occurs 2–10 weeks after [myocardial infarction \(MI\)](#) due to immune complex-mediated damage in the [pericardium](#). While this patient does present with signs and symptoms of [acute pericarditis](#) (e.g., [fever](#), [pleuritic chest pain](#), dry [cough](#), and diffuse [ST elevation](#)), he does not have a history of [coronary artery disease](#) or past [MI](#), making [Dressler syndrome](#) unlikely. Rather, his 5-day history of [cough](#), [fever](#), and malaise suggests an infectious cause for the [pericarditis](#).


E - Infective endocarditis

Explanation Why

[Infective endocarditis \(IE\)](#) could explain this patient's [fever](#), malaise, [pleuritic chest pain](#), and features of cardiac decompensation (e.g., [dyspnea](#), bilateral [rales](#), distended neck [veins](#)). However, he presents with muffled [heart sounds](#) rather than a new [murmur](#), which is a key manifestation of IE, and he does not have extracardiac manifestations of IE (e.g., [splinter hemorrhages](#), [Janeway lesions](#), [Osler nodes](#), and [Roth spots](#)). Additionally, his [ECG](#) findings suggest another diagnosis.

F - Cardiac tamponade

Image



Explanation Why

[Cardiac tamponade](#), a complication of [acute pericarditis](#), is caused by the combination of increased secretion of fluid into [pericardial space](#) and decreased elasticity of the [pericardium](#) secondary to [inflammation](#). Once the [pericardial effusion](#) becomes sufficiently pronounced, either due to the rate of fluid accumulation or the amount of fluid, the [heart](#) can become compressed, which decreases [cardiac output](#) and ventricular [diastolic](#) filling. This patient presents with the classic symptoms of [cardiac tamponade](#): [hypotension](#), distended neck [veins](#), and muffled [heart sounds](#) (i.e., the [Beck triad](#)) as well as a characteristic [ECG](#) finding of low-voltage [QRS complexes](#) and fluctuating R wave amplitude (i.e., [electrical alternans](#)).

Question # 1

A 60-year-old man comes to the emergency department because of a 2-day history of sharp chest pain and a nonproductive cough. The pain worsens with deep inspiration and improves when he leans forward. Three weeks ago, the patient was diagnosed with an ST-elevation myocardial infarction and underwent stent implantation of the right coronary artery. His temperature is 38.4°C (101.1°F) and blood pressure is 132/85 mm Hg. Cardiac auscultation shows a high-pitched scratching sound during expiration. An x-ray of the chest shows enlargement of the cardiac silhouette and a left-sided pleural effusion. Which of the following is the most likely underlying cause of this patient's current condition?

	Answer	Image
A	Outpouching of ventricular wall	
B	Occlusion of coronary artery stent	
C	Pericardial infiltration with neutrophils	
D	Embolism to left pulmonary artery	
E	Immune response to cardiac antigens	
F	Rupture of interventricular septum	

Hint

Fever, pleuritic chest pain, pericardial friction rub (high-pitched scratching sound), and a pericardial and unilateral pleural effusion occurring 3 weeks after an MI suggest postmyocardial infarction syndrome (Dressler syndrome).

Correct Answer

A - Outpouching of ventricular wall

Explanation Why

In true [ventricular aneurysms](#), a portion of the ventricular free wall affected by the [MI](#) will outpouch, putting patients at risk for [cardiac arrhythmia](#), [heart failure](#), and thrombus formation. [Ventricular aneurysm](#) occurs between 2 weeks and several months after [MI](#), which fits the timeframe of this patient. On chest [x-ray](#), [aneurysms](#) may cause enlargement of the [cardiac silhouette](#). However, [ventricular aneurysm](#) would not cause the [pericardial friction rub](#) or [pleuritic chest pain](#) seen here. Furthermore, the vast majority of [ventricular aneurysm](#) form on the [anterior](#) or apical walls, due to occlusion of the left [anterior descending coronary artery](#); this patient's occlusion was in the [right coronary artery](#).

B - Occlusion of coronary artery stent

Explanation Why

[Stent thrombosis](#), which is an important consideration for any patient with recent coronary stenting, can cause a [myocardial infarction](#) and associated acute-onset [chest pain](#) and [tachypnea](#). However, the [pain](#) of [myocardial infarction](#) is typically described as retrosternal chest discomfort, tightness, or pressure rather than the [pleuritic chest pain](#) this patient is experiencing. Furthermore, [stent thrombosis](#) would not account for this patient's [pericardial friction rub](#) or the [pericardial](#) or [pleural effusions](#).

C - Pericardial infiltration with neutrophils

Explanation Why

Infiltration of the [pericardium](#) with [neutrophils](#) can occur as a complication of [myocardial infarction](#), in which case [neutrophils](#) and [fibrin](#) are found in the [pericardium](#), adjacent to the [necrotic myocardium](#) (early [infarct](#)-associated [pericarditis](#)). Although this patient is at risk for post-MI complications, early [infarct](#)-associated [pericarditis](#) is usually seen within the first week after a

[myocardial infarction](#). This patient is 3 weeks post-[MI](#) and therefore beyond the window for early [infarct](#)-associated [pericarditis](#).

D - Embolism to left pulmonary artery

Explanation Why

Embolism to the [pulmonary artery](#) is the mechanism of [pulmonary embolism](#). Many of this patient's symptoms are similar to those of [PE](#), including [pleuritic chest pain](#), [cough](#), and low-grade [fever](#). This patient's left-sided [pleural effusion](#) could be explained by [PE](#). However, [chest pain](#) that improves with leaning forward indicates a different process, and a [pericardial friction rub](#) and [pericardial effusion](#) are not [features of PE](#).

E - Immune response to cardiac antigens

Explanation Why

2–10 weeks after a [myocardial infarction](#), an autoimmune response against cardiac antigens can lead to immune complex-mediated damage ([type III hypersensitivity reaction](#)) to the [pericardium](#). While the exact mechanism of [immune complex](#) formation in [Dressler syndrome](#) is not fully understood, the release of neoantigens from infarcted [myocardium](#) is suspected to play a role. Patients typically present with features of [acute pericarditis](#) such as [fever](#), [tachypnea](#), [chest pain](#) that worsens during inspiration and is alleviated by leaning forward, dry [cough](#), [pericardial friction rub](#), increased [troponin](#) levels, diffuse [ST elevations](#), [pericardial effusion](#), and [pleural effusions](#).

F - Rupture of interventricular septum

Explanation Why

A [myocardial infarct](#) affecting the interventricular septum can result in its rupture, creating a [ventricular septal defect](#). This causes a [left-to-right shunt](#), resulting in [chest pain](#), new-onset [systolic murmur](#), and [heart failure](#) if the defect is sufficiently large. While septal rupture can happen as late as 2 weeks post-[MI](#), it usually happens at 3–5 days. This patient's symptoms began 3 weeks post-[MI](#), which places him beyond the window for rupture. Furthermore, his [cardiac examination](#) revealed a

[pericardial friction rub](#), not the [systolic ejection murmur](#) that would be expected in interventricular rupture. In addition, an interventricular septal rupture would not explain this patient's [pericardial effusion](#).

Question # 2

A 10-year-old boy is brought to the physician by his father, who is concerned because his son has been less interested in playing soccer with him recently. They used to play every weekend, but his son has started to tire easily and has complained of pain in the lower legs while running around on the soccer field. The boy has no personal or family history of serious illness. Cardiac examination shows a systolic ejection murmur best heard over the left sternal border that radiates to the left paravertebral region. An x-ray of the chest shows erosions of the posterior aspects of the 6th to 8th ribs. If left untreated, this patient is at greatest risk for which of the following?

	Answer	Image
A	Intracranial hemorrhage	
B	Central cyanosis	
C	Paradoxical embolism	
D	Abdominal aortic aneurysm	
E	Right heart failure	

Hint

Intermittent leg pain on exertion suggests inadequate perfusion of the lower extremities (intermittent claudication). The systolic ejection murmur described is likely due to aortic narrowing, and the erosions of the ribs suggest collateral circulation through the intercostal arteries (rib notching). Together, these findings indicate aortic coarctation.

Correct Answer

A - Intracranial hemorrhage

Explanation Why

Patients with untreated [aortic coarctation](#) usually develop [secondary hypertension](#), resulting in an increased risk of [intracranial hemorrhage](#). Moreover, [aortic coarctation](#) is associated with [intracranial aneurysm](#) formation (due to [hypertension](#) and/or genetic [risk factors](#)), which increases the risk of hemorrhage due to [aneurysm](#) rupture. Other complications of [aortic coarctation](#) include [aortic dissection](#), [coronary artery disease](#), [ischemic stroke](#), [endocarditis](#), and [heart failure](#).

B - Central cyanosis

Explanation Why

Central [cyanosis](#) is seen in patients with congenital [heart](#) defects that lead to [right-to-left shunting](#), such as [tetralogy of Fallot](#) and [transposition of the great vessels](#). It is not typically seen in [aortic coarctation](#), which is considered an [acyanotic heart defect](#). [Cyanotic heart defects](#) usually become evident in the [newborn](#) period, often manifesting with acute, life-threatening episodes of [cyanosis](#). If left untreated, most defects are fatal within the first year of life.

C - Paradoxical embolism

Explanation Why

[Paradoxical embolism](#) is a possible complication of shunts that allow a [thrombus](#) to pass from venous to arterial circulation, such as [patent foramen ovale](#) and [atrial septal defect](#) (ASD). Since both of these conditions are usually asymptomatic, [intermittent claudication](#) would not be expected. Moreover, unlike this patient's [murmur](#), ASD manifests with a mid-[systolic](#) pulmonary flow or ejection [murmur](#) accompanied by wide, fixed [splitting of S₂](#).

D - Abdominal aortic aneurysm

Explanation Why

Risk factors for abdominal aortic aneurysm (AAA) include smoking, advanced age, atherosclerosis, and arterial hypertension. Although aortic coarctation often leads to secondary hypertension, blood flow in the vessels distal to the constriction at the aortic isthmus is actually decreased, resulting in hypotension of the lower extremities. Therefore, the risk of AAA is not increased. However, increased blood flow proximal to the constriction increases the risk of dilation of the aorta and subsequent dissection.

E - Right heart failure

Explanation Why

Left ventricular outflow tract obstruction caused by aortic coarctation results in compensatory left ventricular hypertrophy and arterial hypertension, either of which can eventually result in left-sided heart failure. Although right-sided heart failure can occur secondary to left-sided heart failure, this patient is at a higher risk of a different complication.

Question # 3

A 67-year-old man with stable coronary artery disease comes to the physician for a follow-up examination. Aside from occasional exertional chest pain on mowing the lawn or prolonged jogging, he feels well. He goes jogging for 20 minutes once a week and takes a tablet of sublingual nitroglycerine prior to his run to prevent anginal chest pain. The patient would like to run longer distances and asks the physician whether he could increase the dose of the drug prior to running. Administration of higher dosages of this drug is most likely to result in which of the following?

	Answer	Image
A	Rebound angina	
B	Development of tolerance	
C	Reflex sympathetic activity	
D	Anaphylactic reaction	
E	Coronary artery vasospasm	
F	Symptoms of withdrawal	

Hint

Administration of metoprolol would help to counteract this adverse effect!

Correct Answer

A - Rebound angina

Explanation Why

Rebound [angina](#) is the increase in [anginal](#) symptoms due to decreased drug levels during the [nitrate](#)-free intervals in intermittent therapy. It is usually associated with the use of transdermal [nitroglycerin](#) patches, not with sublingual administration. Moreover, this patient's once-weekly usage puts him at low risk for rebound [angina](#).

B - Development of tolerance

Explanation Why

[Nitrate](#) tolerance often develops after 2–3 days of continuous administration and is more likely with high dosages and long-acting preparations. In order to minimize this side effect, daily [nitrate](#)-free intervals of at least 8 hours are recommended. Since this patient only has an estimated once-weekly usage with adequate medication-free intervals, a higher dosage will not increase his risk of developing tolerance.

C - Reflex sympathetic activity

Explanation Why

Low-dose [nitrate](#) improves [myocardial](#) perfusion primarily by acting as a venodilator that decreases ventricular wall tension via [preload](#) reduction. In addition, high-dose [nitrate](#) dilates coronary and peripheral [arteries](#), thereby preventing coronary spasms and reducing ventricular [afterload](#) along with subsequent myocardial oxygen demand. However, peripheral arterial dilation also causes systemic [hypotension](#), which induces a reflex [sympathetic](#) activity leading to [tachycardia](#) and increased [myocardial contractility](#). As a result, myocardial oxygen demand paradoxically increases. Simultaneous administration of [\$\beta\$ -blockers](#) has a synergetic anti-[ischemic](#) effect and can help prevent [reflex tachycardia](#).

D - Anaphylactic reaction

Explanation Why

[Anaphylactic reactions](#) occur in < 1% of patients treated with [nitrates](#). Furthermore, this patient has been taking this drug regularly, making an [allergic reaction](#) to an increase in dosage unlikely.

E - Coronary artery vasospasm

Explanation Why

[Nitrates](#) dilate [coronary arteries](#) and thus prevent rather than cause [coronary artery](#) spasms. Therefore, they are regularly used in the long-term treatment of [Prinzmetal angina](#).


F - Symptoms of withdrawal

Explanation Why

Symptoms of withdrawal can occur after abrupt cessation of use of an addictive recreational or therapeutic substance. [Nitrates](#) are not known to cause addiction.

Question # 4

A 63-year-old man who recently immigrated to the United States from Indonesia comes to the physician because of worsening shortness of breath and swollen extremities for the past 3 months. He has had a 4-kg (8.8-lb) weight loss and intermittent fevers over the last 6 months. Examination shows pitting pedal edema and abdominal distension. Abdominal pressure over the right upper quadrant produces persistent distention of the jugular veins. An x-ray of the chest shows cavernous infiltrates in the left and right-upper lobes and a calcified cardiac silhouette. Cardiovascular examination is most likely to show which of the following?

	Answer	Image
A	Absent jugular venous pulse y descent	
B	Friction rub	
C	Pericardial knock	
D	Fourth heart sound	

	Answer	Image
E	Pulsus parvus et tardus	
F	Fixed split second heart sound	


Hint

Dyspnea, peripheral pitting edema, abdominal distension, and a positive hepatojugular reflux in the context of a calcified cardiac silhouette suggest constrictive pericarditis.

Correct Answer

A - Absent jugular venous pulse y descent

Image



Explanation Why

The [y descent](#) of the [jugular venous pulse](#) waveform corresponds to the emptying of the [right atrium](#) following the opening of the [tricuspid valve](#). Absent [y descent](#) would indicate resistance to all phases of [diastolic](#) filling, which is seen in [cardiac tamponade](#). In [constrictive pericarditis](#), the early phase of [diastolic](#) filling is unchanged, but the mid-late phase is met with resistance. This, coupled with the high [JVD](#) of chronic venous congestion, produces a prominent [y descent](#) that is accentuated in the early phase as filling drops rapidly from a higher [peak pressure](#).

B - Friction rub

Explanation Why

In [acute pericarditis](#), [inflammation](#) of the [pericardial](#) layers can cause friction as they move during the [cardiac cycle](#), creating an audible friction rub on [cardiac auscultation](#). While a friction rub is characteristic of the acute phase of [pericarditis](#), a different [heart sound](#) is pathognomonic for chronic [constrictive pericarditis](#), in which the ventricles are working against a calcified and thickened [pericardium](#).

C - Pericardial knock

Explanation But

This patient's history of immigration from Indonesia as well as his clinical presentation of [fever](#), weight loss, and cavernous lesions in the upper [lung](#) lobes make [tuberculosis](#) the most likely cause of his [constrictive pericarditis](#). In developed countries, prior cardiac surgery, [radiation therapy](#) to the chest, and viral infection are common causes.

Explanation Why

A [pericardial knock](#) is often heard on auscultation in [constrictive pericarditis](#) in which [diastolic](#) ventricular relaxation and filling are abruptly halted by the rigid [pericardial sac](#). A [pericardial knock](#) is a high-pitched, early-[diastolic](#) sound that sounds like a premature [S3](#). In [constrictive pericarditis](#), reduced [ventricular filling](#) leads to manifestations of reduced [cardiac output](#) (forward failure), such as fatigue, [dyspnea](#), as well as those of fluid overload (backward failure), such as [JVD](#), positive [Kussmaul sign](#), [hepatomegaly](#), and the clinical features seen in this patient.

D - Fourth heart sound

Explanation Why

A [fourth heart sound](#) (S4) is caused by a late [diastolic](#) contraction of the [atria](#) and is seen in

[restrictive cardiomyopathy](#), [left ventricular hypertrophy](#), and [ischemic heart disease](#). [Restrictive cardiomyopathy](#) is an important differential diagnosis of [constrictive pericarditis](#) because it can present similarly. However, it does not cause calcification of the [cardiac silhouette](#). An S4 is not a [physical exam](#) finding of [constrictive pericarditis](#), most likely because the rigid [pericardial sac](#) impairs atrial contraction.

E - Pulsus parvus et tardus

Explanation Why

Pulsus parvus et tardus is characteristic of the prolonged [left ventricular](#) total ejection time seen in [aortic stenosis](#). [Aortic stenosis](#) would cause signs and symptoms of [left heart failure](#), but not the [symptoms of right heart failure](#) seen in this patient. Furthermore, calcification in [aortic stenosis](#) is limited to the [aortic valve](#) and does not involve the entire [cardiac silhouette](#), as seen here.

F - Fixed split second heart sound

Explanation Why

Fixed [splitting of S₂](#) is pathognomonic for [atrial septal defect](#). Although patients with [constrictive pericarditis](#) may occasionally have wide [splitting of S₂](#) during inspiration due to a reduction in the [left ventricular](#) ejection time, they do not have fixed [splitting of S₂](#).

Question # 5

A 51-year-old woman comes to the physician because of worsening chest pain on exertion. She was diagnosed with coronary artery disease and hyperlipidemia 3 months ago. At the time of diagnosis, she was able to walk for 15 minutes on the treadmill until the onset of chest pain. Her endurance had improved temporarily after she began medical treatment and she was able to walk her dog for 30 minutes daily without experiencing chest pain. Her current medications include daily aspirin, metoprolol, atorvastatin, and isosorbide dinitrate four times daily. Her pulse is 55/min and blood pressure is 115/78 mm Hg. Treadmill walking test shows an onset of chest pain after 18 minutes. Which of the following is most likely to improve this patient's symptoms?

	Answer	Image
A	Increase dose of daily metoprolol	
B	Avoid isosorbide dinitrate at night	
C	Discontinue atorvastatin therapy	
D	Add tadalafil to medication regimen	
E	Perform coronary angiography	
F	Decrease amount of aerobic exercise	

Hint

This patient experiences stable anginal chest pain which initially improved with medication, but now the patient's symptom level is similar to the level prior to the initiation of medication. This is likely due to development of tolerance to one of her drugs.

Correct Answer

A - Increase dose of daily metoprolol

Explanation Why

[Beta blockers](#) such as [metoprolol](#) are a first-line therapy for [stable angina](#), as their effect of lowering [heart rate](#) and contractility reduces [ischemia](#). However, this patient already has a low [heart rate](#), indicating that [metoprolol](#) is still effective. Increasing the [metoprolol](#) dose further could cause hemodynamic instability.

B - Avoid isosorbide dinitrate at night

Explanation Why

Development of tolerance is a typical side-effect of chronic [nitrate](#) therapy. The mechanism underlying [nitrate](#) tolerance is not completely understood but is thought to involve decreased sensitivity of the vascular [endothelium](#) to [nitrates](#). The most effective strategy to prevent this is intermittent therapy with [nitrate](#)-free intervals of at least 8 hours, which would be accomplished by avoiding [isosorbide dinitrate](#) at night.

C - Discontinue atorvastatin therapy

Explanation Why

[Statins](#) are commonly used in the treatment of both [hyperlipidemia](#) and [coronary artery disease](#). Discontinuation may be necessary in the setting of intolerable side effects or [statin-induced myopathy](#) which would present with symptoms of muscle [pain](#) and weakness. This patient, however, is experiencing [drug tolerance](#), not drug toxicity.

D - Add tadalafil to medication regimen

Explanation Why

[Phosphodiesterase type-5 inhibitor](#) such as [tadalafil](#) are contraindicated in patients taking [nitrates](#), because the synergistic [vasodilation](#) can result in extreme [hypotension](#) and [syncope](#). Furthermore, [tadalafil](#) is indicated for the treatment of [erectile dysfunction](#), [BPH](#), and [pulmonary hypertension](#) but not [angina pectoris](#).

E - Perform coronary angiography

Explanation Why

Invasive tests such as [coronary angiography](#) are reserved for patients with [stable angina](#) who are either refractory to optimal medical management or belong to a high-risk demographic. This patient was asymptomatic after initial pharmacotherapy but is now exhibiting signs of [drug tolerance](#), indicating that medical management is sub-optimal. Furthermore, angiography alone would not treat this patient's symptoms since it is a diagnostic test, not a therapeutic intervention.

F - Decrease amount of aerobic exercise


Explanation Why

Patients with [stable angina pectoris](#) should be encouraged to engage in regular aerobic exercise, since this can result in a lower oxygen requirement of the [myocardium](#) for a given workload, thereby improving exercise tolerance and reducing symptoms. Furthermore, this patient has experienced improvement from her original 15-minute exercise tolerance which indicates that her capacity for aerobic exercise has not decreased.


Question # 6

An otherwise healthy 25-year-old man comes to the physician because of a 2-day history of chills, palpitations, and worsening shortness of breath. The patient is 183 cm (6 ft) tall and weighs 70 kg (154 lbs); BMI is 21 kg/m². His temperature is 39.5°C (103.1°F), pulse is 110/min, and blood pressure is 140/60 mm Hg. Examination of the extremities shows several nontender, nonblanching, erythematous macules on the palms and soles and painful nodules on the pads of the fingers and toes. Coarse crackles are heard over both lungs. Cardiac examination is shown. Results of initial blood cultures are pending. Echocardiography is most likely to show which of the following abnormalities?

[Video](#)

	Answer	Image
A	Aortic regurgitation	 The image displays a comparison between an Electrocardiogram (ECG) and a Phonocardiogram (PCG) for aortic regurgitation. The ECG (top trace, blue) shows a normal sinus rhythm with a narrow QRS complex. The PCG (bottom trace, orange) shows aortic regurgitation, characterized by a high-pitched, decelerating diastolic murmur. The background is divided into Diastole and Systole phases, with S1 and S2 markers indicating the timing of the heart sounds. The PCG shows a prominent diastolic murmur following S2, which is characteristic of aortic regurgitation.
B	Pulmonary regurgitation	
C	Tricuspid stenosis	

	Answer	Image
D	Mitral stenosis	The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The background is divided into alternating Diastole (grey) and Systole (white) phases. The ECG shows a normal sinus rhythm. The PCG shows a low-intensity S1, a high-intensity S2, and a prominent, long-duration diastolic murmur (OS) that starts at the second heart sound and continues through diastole. The labels S1 and S2 are positioned below the corresponding heart sounds.
E	Aortic stenosis	The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The background is divided into alternating Diastole (grey) and Systole (white) phases. The ECG shows a normal sinus rhythm. The PCG shows a low-intensity S1, a high-intensity S2, and a prominent, long-duration systolic murmur that starts at the first heart sound and continues through systole. The labels S1 and S2 are positioned below the corresponding heart sounds.
F	Mitral valve prolapse	The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The background is divided into alternating Diastole (grey) and Systole (white) phases. The ECG shows a normal sinus rhythm. The PCG shows a low-intensity S1, a high-intensity S2, and a prominent, long-duration systolic murmur that starts at the first heart sound and continues through systole. A label 'Mid-systolic click' points to a sharp, high-frequency sound occurring mid-systole. The labels S1 and S2 are positioned below the corresponding heart sounds.

	Answer	Image
G	Mitral regurgitation	 The image displays a comparison between an Electrocardiogram (ECG) and a Phonocardiogram (PCG) for Mitral regurgitation. The ECG (top trace, blue) shows a normal sinus rhythm with labeled Diastole and Systole phases. The PCG (bottom trace, orange) shows a normal S1 (first heart sound) and a significantly prolonged and decelerating S2 (second heart sound), which is characteristic of mitral regurgitation. The background is divided into alternating light blue (Diastole) and light orange (Systole) vertical bands. The S1 and S2 labels are positioned below the PCG trace.
H	Pulmonary stenosis	
I	Tricuspid regurgitation	


Hint

This patient most likely has acute infective endocarditis (IE), which is indicated by a high fever, chills, palpitations, dyspnea, Janeway lesions (nontender, nonblanching, erythematous macules on the palms and soles), and Osler nodes (painful nodules on pads of the fingers and toes). The carotid pulsations heard in the audio clip indicate ventricular systole. There is a high-pitched, nonradiating, blowing, decrescendo early diastolic murmur, heard best at the left sternal border.

Correct Answer

A - Aortic regurgitation

Image



Explanation Why

The auscultatory findings in this patient are characteristic of [aortic regurgitation](#) (AR), which causes [left ventricular end-diastolic volume](#) overload as the [left ventricle](#) receives blood from the [left atrium](#) and the regurgitant aorta. This overload results in hyperdynamic circulation, which manifests as [palpitations](#) (especially in the left [lateral](#) decubitus position, when the enlarged [left ventricle](#) is closest to the [chest wall](#)), a widened [pulse pressure](#), and [dyspnea](#) due to [pulmonary edema](#) resulting from [left heart failure](#), as seen here. The most common cause of acute valvular AR is IE, which this patient has. AR due to valvular disease is heard best along the left sternal border, as seen in this patient, while AR due to aortic root disease (e.g., in [Marfan syndrome](#)) is heard best along the right sternal border. Other common causes of AR include [aortic dissection](#), [acute rheumatic fever](#), and congenital [bicuspid aortic valve](#).

B - Pulmonary regurgitation

Explanation Why

The [murmur](#) characteristic of pulmonary regurgitation (PR) is very similar to the one heard in the audio clip (i.e., an early [diastolic](#) decrescendo [murmur](#) heard best at the left sternal border). Although IE can cause PR, the pulmonic valve is only rarely affected. Furthermore, PR is usually asymptomatic until it leads to [right heart failure](#), manifesting with [dyspnea](#) on exertion and fatigue rather than [palpitations](#) and [pulmonary edema](#). Severe PR is most commonly [iatrogenic](#); other causes include [carcinoid syndrome](#), congenital [heart](#) defects, and [rheumatic heart disease](#).


C - Tricuspid stenosis

Explanation Why

The [murmur](#) characteristic of tricuspid stenosis (TS) is very similar to the [diastolic murmur](#) heard in the audio clip. TS causes a soft mid-[diastolic murmur](#) heard best in the parasternal region of the left 4th intercostal space. TS is, however, extremely rare, and symptomatic TS typically manifests with [features of right-sided heart failure](#) rather than the [palpitations](#) and symptoms of [pulmonary edema](#) seen here.

D - Mitral stenosis

Image




Explanation Why

[Mitral stenosis](#) (MS) can manifest as [dyspnea](#) (due to pulmonary venous congestion) and [palpitations](#) (if [atrial fibrillation](#) develops), both of which are seen here. Although the [murmur](#) that is characteristic of MS also occurs during [diastole](#), as heard in this clip, it is most often a mid-[diastolic](#) decrescendo [murmur](#) preceded by an [opening snap](#) and is heard best at the cardiac apex. This patient, however, has an early [diastolic murmur](#) heard best at the left sternal border.

E - Aortic stenosis

Image




Explanation Why

The characteristic triad of symptomatic [aortic stenosis](#) (AS) is exertional dyspnea, [stable angina](#), and exertional [dizziness/syncope](#) caused by [left ventricular](#) outflow obstruction. [Palpitations](#), which are seen in this patient, are not typical of AS. AS causes weak peripheral pulses ([pulsus parvus et tardus](#)) and low/narrow [pulse pressure](#), in contrast to this patient's findings. Moreover, the characteristic [murmur](#) of AS is a harsh crescendo-decrescendo late [systolic ejection murmur](#) heard best over the right 2nd intercostal space, which radiates to the carotids.

F - Mitral valve prolapse

Image



Explanation Why

[Mitral valve prolapse \(MVP\)](#) is typically asymptomatic but can manifest with [palpitations](#) and [dyspnea](#), which are seen here. [MVP](#) is one of the most common valvular abnormalities in the US, especially in young adults, and a [risk factor](#) for IE, which this patient has. However, widened [pulse pressure](#), also seen here, is not caused by [MVP](#). Typical cardiac auscultatory findings in [MVP](#) would also include a high-frequency, midsystolic click with a late [systolic murmur](#) that is heard best at the cardiac apex (left [midclavicular line](#) of the 5th intercostal space).

G - Mitral regurgitation

Image



Explanation Why

[Mitral regurgitation](#) ([MR](#)) can manifest with [dyspnea](#) (due to pulmonary venous congestion) and [palpitations](#) (if [atrial fibrillation](#) develops), both of which are seen here. In addition, [MR](#) is the most common [murmur](#) resulting from IE, which this patient has. However, [MR](#) manifests with a [holosystolic murmur](#) heard best at the cardiac apex (left 5th intercostal space), in contrast to this patient's [diastolic murmur](#).

H - Pulmonary stenosis

Explanation Why

[Pulmonary stenosis](#) (PS) is usually a congenital [heart](#) defect. Acquired PS is rare but can occur in patients with [carcinoid syndrome](#) or [rheumatic heart disease](#). It would typically manifest with

[features of right-sided heart failure](#) rather than the [dyspnea](#) and [palpitations](#) seen in this patient. The typical [murmur](#) of PS is an ejection [systolic murmur](#), which is heard best parasternally at the left 2nd intercostal space, often with a widely [split S₂](#). This patient's [diastolic murmur](#) rules out PS as a diagnosis.

I - Tricuspid regurgitation

Explanation Why

[Tricuspid regurgitation](#) (TR) is a rare valvular abnormality that is typically asymptomatic and can be caused by IE, which this patient has. However, right-sided [valvular heart diseases](#) are more common in individuals with particular [risk factors](#) (e.g., IV drug use), which are not seen here. Chronic and/or severe TR manifests with [features of right-sided heart failure](#), not with [dyspnea](#) and [palpitations](#), which are [features of left-sided heart failure](#) and seen in this patient. In addition, TR causes a [holosystolic murmur](#) heard best at the 4th intercostal space of the left sternal border rather than a [diastolic murmur](#).

Question # 7

A 61-year-old man is brought to the emergency department by ambulance because of severe retrosternal chest pain and shortness of breath for 30 minutes. Paramedics report that an ECG recorded en route to the hospital showed ST-segment elevation in I, aVL, and the precordial leads. On arrival, the patient is unresponsive to painful stimuli. Examination shows neither respiration nor pulse. Despite appropriate lifesaving measures, he dies 10 minutes later. Which of the following is the most likely cause of death in this patient?

	Answer	Image
A	Cardiac septal rupture	
B	Left ventricular failure	
C	Ventricular aneurysm	
D	Hemorrhagic stroke	
E	Cardiac free wall rupture	
F	Ventricular fibrillation	

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

Hint

The patient's chest pain and dyspnea combined with the finding of ST-segment elevation in a proximal left anterior descending distribution on ECG are highly suggestive of myocardial infarction (MI). The patient's subsequent death within an hour of symptom onset is known as sudden cardiac death (SCD).

Correct Answer

A - Cardiac septal rupture

Explanation Why

[MI](#) that affects the interventricular septum has the potential to rupture, creating a [left-to-right shunt](#) and [right heart failure](#). While this would also present with [shortness of breath](#), it would usually not occur until 3–5 days after [MI](#).

B - Left ventricular failure

Explanation Why

Depending on the extent and distribution of a [MI](#), contractility of the [left ventricle](#) can decrease to the point of [left ventricular failure](#). In addition, [MI](#) can cause [papillary muscle rupture](#) resulting in [mitral valve regurgitation](#) and [left ventricular failure](#). While acute [left heart failure](#) secondary to [MI](#) does present with [shortness of breath](#) and [chest pain](#), it typically occurs several hours after [MI](#) and is thus not the most common cause of SCD.

C - Ventricular aneurysm

Explanation Why

In [ventricular aneurysm](#), a portion of the ventricular free wall affected by the [MI](#) will outpouch, putting patients at risk for [cardiac arrhythmia](#), [heart failure](#), and [cardiac tamponade](#), all of which can cause [sudden cardiac death](#). However, [ventricular aneurysm](#) happens between 2 weeks and several months after [MI](#).

D - Hemorrhagic stroke

Explanation Why

[Thromboembolism](#) can result from left [ventricular aneurysm](#) or new onset [atrial fibrillation](#) post-[MI](#) and can cause [hemorrhagic stroke](#). However, these complications typically take days to weeks to develop, and [stroke](#) is not a common cause of [sudden cardiac death](#) in patients with [MI](#).

E - Cardiac free wall rupture

Explanation Why

[MI](#) can cause weakness of the ventricular free wall, which makes it [prone](#) to rupture under the high pressures of [systole](#). This results in [cardiac tamponade](#) and is often fatal. However, ventricular free wall rupture occurs usually within the first 2 weeks of [MI](#), when [necrosis](#) has had time to sufficiently weaken the ventricular free wall tissue, not during first hour post-[MI](#).

F - Ventricular fibrillation

Explanation Why

[Ventricular fibrillation](#) is a [cardiac arrhythmia](#) in which multiple [ectopic foci](#) of electrical activity within the ventricle cause disorganized [depolarization](#) and diminished [cardiac output](#). Patients with [ventricular fibrillation](#) require rapid [defibrillation](#) and [CPR](#), otherwise hemodynamic instability can lead to loss of consciousness and SCD, which is the leading cause of death post-[MI](#). SCD after [MI](#) is usually due to [ventricular tachyarrhythmias](#) such as [ventricular tachycardia](#) or [ventricular fibrillation](#).

Question # 8

A 2-week-old male newborn is brought to the physician because his mother has noticed her son has occasional bouts of "turning blue in the face" while crying. He also tires easily and sweats while feeding. He weighed 2150 g (4 lb 11 oz) at birth and has gained 200 g (7 oz). The baby appears mildly cyanotic. Examination shows a 3/6 systolic ejection murmur heard over the left upper sternal border. A single S₂ is present. An echocardiography confirms the diagnosis. Which of the following factors is the main determinant of the severity of this patient's cyanosis?

	Answer	Image
A	Right ventricular outflow obstruction	
B	Left ventricular outflow obstruction	
C	Right ventricular hypertrophy	
D	Ventricular septal defect	
E	Overriding aorta	
F	Atrial septal defect	

Hint

The presence of (intermittent) cyanosis, fatigue, and sweating during periods of exertion in a neonate with a systolic ejection murmur and a single S₂ is suggestive of tetralogy of Fallot (TOF).

Correct Answer

A - Right ventricular outflow obstruction

Explanation But

Older children with TOF tend to squat during spells of [hypoxia](#), which relieves symptoms by increasing [systemic vascular resistance](#) and, in turn, LV outflow resistance. This creates higher LV pressures, so that blood in the RV will be directed through the pulmonary vasculature rather than the [VSD](#). The increased blood flow across the RV outflow obstruction during squatting is also associated with an increased intensity of the ejection [murmur](#).

Explanation Why

In patients with [tetralogy of Fallot](#) (TOF), the [right ventricular](#) (RV) outflow obstruction due to central [pulmonary valve stenosis](#) and [hypoplasia](#) determines the severity of [cyanosis](#) and the intensity of the [systolic heart murmur](#). The higher the degree of RV outflow obstruction, the more (unoxygenated) blood will bypass [pulmonary circulation](#) and shunt straight through the [VSD](#) to the [left ventricle](#) (LV). The lower the degree of RV outflow obstruction, the more blood flows through the [lungs](#) and becomes oxygenated before reaching the LV, resulting in a reduction of [cyanosis](#). In TOF, the single S₂ can result from an attenuated pulmonary component (P₂) that is often not heard on auscultation.

B - Left ventricular outflow obstruction

Explanation Why

[Left ventricular](#) outflow obstruction (e.g., [aortic stenosis](#)) is not one of the four features of TOF. However, if [aortic stenosis](#) (or any form of increased [left ventricular](#) outflow resistance) were present, this would decrease [right-to-left shunting](#) by creating higher left-sided pressures and thus reduce, rather than increase, [cyanosis](#).

C - Right ventricular hypertrophy

Explanation Why

Although [right ventricular hypertrophy \(RVH\)](#) is one of the four features of TOF, it is not the main determinant of [cyanosis](#). [RVH](#) is a compensatory mechanism against the outflow tract obstruction, which helps to reduce [cyanosis](#) by increasing flow through the stenotic pulmonary vasculature, so that deoxygenated blood does not flow through the [VSD](#) and into the [systemic circulation](#). However, [RVH](#) can only compensate to a small degree to improve normal circulatory flow and is, therefore, only a minor determinant of [cyanosis](#).

D - Ventricular septal defect

Explanation Why

A [ventricular septal defect \(VSD\)](#) is one of the four features of TOF but it is not the main determinant of [cyanosis](#). Rather, it is the pressure differences across the [VSD](#) that determine the degree of [cyanosis](#). Specifically, higher [right ventricular](#) pressures and/or lower [left ventricular](#) pressures are associated with increased right-to-left shunting and, thus, with increased [cyanosis](#).

E - Overriding aorta

Explanation Why

Although an overriding aorta is one of the four features of TOF, it is not the main determinant of [cyanosis](#). In patients with TOF, the aorta is displaced to the [right ventricle](#), over the [VSD](#), allowing blood from both ventricles to be pumped into the aorta. This causes deoxygenated blood to enter the [systemic circulation](#), contributing to [cyanosis](#). However, another factor has a greater impact on the degree of [cyanosis](#).


F - Atrial septal defect

Explanation Why

[Atrial septal defect](#) (ASD) is not one of the four features of TOF. However, in a patient with TOF, the presence of a concomitant ASD could lead to greater right-to-left shunting and thus exacerbate [cyanosis](#).

Question # 9

A 31-year-old man comes to the emergency department for acute tearing chest pain that radiates to the back. Despite appropriate therapy, the patient dies. Autopsy shows an increase in mucoid extracellular matrix and loss of smooth muscle cell nuclei in the media of large arteries. Which of the following additional findings is most likely in this patient?

	Answer	Image
A	Nasal septum perforation	
B	Inferonasal lens dislocation	
C	Pes cavus with hammer toes	
D	Polycystic kidneys	
E	Pectus carinatum	Marfan syndrome Epidemiology Prevalence: 1-5/10,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15; autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissections) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups) 

	Answer	Image
F	Condylomata lata	

Hint

Acute tearing chest pain with radiation to the back is consistent with aortic dissection, a life-threatening condition. Muroid extracellular matrix accumulation and loss of smooth muscle cell nuclei from cystic medial degeneration of the large arteries predisposes to aortic dissection, and, in young patients, is indicative of connective tissue disorders such as Marfan syndrome.

Correct Answer

A - Nasal septum perforation

Explanation Why

[Nasal septum perforation](#) can be seen in patients with a history of repeated nasal inhalation of [cocaine](#). Although repeated [cocaine](#) use can cause acute [aortic dissection](#) in a young patient (particularly in those with underlying [hypertension](#)), it would not result in histopathologic changes of [cystic medial degeneration](#), as seen in this patient.

B - Inferonasal lens dislocation

Explanation Why

Inferonasal (down and inward) [lens dislocation](#) can occur in patients with underlying [homocystinuria](#). These patients typically have a [marfanoid habitus](#) and an increased risk of developing [coronary artery disease](#) and [stroke](#) because [hyperhomocysteinemia](#) induces [atherosclerosis](#) and a [hypercoagulable state](#). They do not have an increased risk of developing an [aortic dissection](#) or [cystic medial degeneration](#) of the large [arteries](#).

C - Pes cavus with hammer toes

Image



Explanation Why

[Pes cavus](#) with [hammer toes](#) is a characteristic finding in [Friedreich ataxia](#). Although many patients with this condition die at a young age due to cardiovascular complications, the major cause of death is often [heart failure](#) resulting from underlying secondary [hypertrophic cardiomyopathy](#), not [aortic dissection](#).

D - Polycystic kidneys


Explanation Why

[Autosomal dominant polycystic kidney disease](#) increases the risk of developing cerebral [berry aneurysms](#). It does not directly affect the aorta or predispose to [aortic dissection](#). Cardiovascular complications of [ADPKD](#) include [mitral valve prolapse](#), [left ventricular hypertrophy](#) (typically

asymptomatic), and [hypertension](#) (which can increase the risk of developing an [aortic dissection](#)). [Hypertension](#) would not cause the histopathologic changes of [cystic medial degeneration](#) seen in this patient.

E - Pectus carinatum

Image



Explanation But


Patients with [Marfan syndrome](#) classically have [marfanoid habitus](#), [joint](#) hypermobility, a high-arched [palate](#), [arachnodactyly](#), [skin](#) hyperelasticity, and superotemporal [lens subluxation](#).

Explanation Why

Deformity of the [chest wall](#) ([pectus carinatum](#) or [pectus excavatum](#)) is a common finding in patients with [Marfan syndrome](#). This condition is strongly associated with aortic pathology, including [aortic dissection](#) (as seen in this patient), [aortic valve regurgitation](#), [aortic aneurysm](#), and [aortic root dilation](#). [Marfan syndrome](#) is caused by a mutation in the [fibrillin-1 gene](#), which encodes an [extracellular matrix](#) glycoprotein that forms a supportive sheath around [elastin](#). Defects in [fibrillin-1](#) make [elastin](#) fibers vulnerable to fragmentation and predispose to [mucoid extracellular matrix accumulation](#), [medial fibrosis](#), and loss of [smooth muscle cell](#) nuclei.

F - Condylomata lata

Image



Explanation Why

[Condylomata lata](#) are characteristic of [secondary syphilis](#), not [Marfan](#) syndrome. In addition to [condylomata lata](#), aortic involvement is possible in advanced syphilitic disease but typically does not arise until the final stage ([tertiary syphilis](#)) and would be an unlikely finding in this young patient. Furthermore, histopathologic findings of affected [arteries](#) in [tertiary syphilis](#) show degeneration of the [vasa vasorum](#) and vascular wall [atrophy](#) (not [cystic medial degeneration](#), as seen in this patient).

Question # 10

A previously healthy 35-year-old woman comes to the emergency department because of sudden weakness of her right arm and leg that started 3 hours ago. She returned from a business trip from Europe 3 days ago. She has smoked a pack of cigarettes daily for 10 years. Her only medication is an oral contraceptive. Her temperature is 38.0°C (100.4°F), pulse is 115/min and regular, and blood pressure is 155/85 mm Hg. Examination shows decreased muscle strength on the entire right side. Deep tendon reflexes are 4+ on the right. Babinski sign is present on the right. The left lower leg is swollen, erythematous, and tender to palpation. Further evaluation is most likely to show which of the following?

	Answer	Image
A	Patent foramen ovale	
B	Atrial myxoma	
C	Mitral valve vegetation	
D	Ventricular septal defect	
E	Atrial fibrillation	
F	Carotid artery dissection	

Hint

This patient has multiple risk factors for deep venous thrombosis (DVT), including a history of immobilization (business trip by plane), smoking, and oral contraceptive use. Now, she presents with clinical features of a left-sided stroke (e.g., right-sided weakness, hyperreflexia, positive Babinski), which in a young patient with signs of DVT (swollen, tender, erythematous left leg) should raise concern for paradoxical embolism.

Correct Answer

A - Patent foramen ovale

Explanation Why

[Paradoxical emboli](#) can pass through any [right-to-left shunt](#), but a [patent foramen ovale \(PFO\)](#) is the most common route. A [PFO](#) is considered an abnormal anatomical variant of the [heart](#) and is present in ~ 25% of the general population. In addition to [stroke](#) workup (i.e., noncontrast head CT, possible [thrombolysis](#), etc.), she should also undergo [compression ultrasonography](#) to confirm [DVT](#). Because of the risk of [pulmonary embolism](#) and recurrent cerebral emboli, this patient should be started on [anticoagulant](#) therapy once cerebral hemorrhage is ruled out and undergo [echocardiography](#) to evaluate for surgical closure of the [PFO](#).

B - Atrial myxoma

Explanation Why

Primary cardiac tumors (e.g., [atrial myxoma](#)) are extremely rare ([incidence](#) < 0.1%), and can sometimes embolize, causing [stroke](#). Most patients with primary cardiac tumors do not initially present with [stroke](#), but experience symptoms of [heart failure](#) (e.g., [dyspnea](#), [edema](#)) due to the [tumor](#) obstructing blood flow. Additionally, patients often experience constitutional symptoms (e.g., [fever](#), weight loss). While this patient has [stroke](#) and [fever](#), she lacks signs of heart failure. Her unilateral leg abnormalities are more suggestive of [DVT](#).

C - Mitral valve vegetation

Explanation Why

Embolism of [septic](#) vegetations from [infective endocarditis](#) (IE) that go to the brain can cause [stroke](#). Patients with IE can also present with [fever](#) and [tachycardia](#). However, IE is unlikely in patients without major [risk factors](#) for the disease (e.g., recent dental procedure, IV [drug abuse](#), active untreated bacterial infection). This patient's warm, swollen leg could be confused with [cellulitis](#),

which would increase her risk of IE, but without a possible entry point for pathogens as in this case, a noninfectious diagnosis is more likely.

D - Ventricular septal defect

Explanation Why

[Ventricular septal defects](#) are the most common pediatric septal defect and can cause [paradoxical embolism](#), which has likely occurred in this patient. However, most [VSDs](#) self-resolve during childhood (and the majority of others are surgically repaired). [VSDs](#) make up ~ 10% of congenital [heart](#) defects found in adults.

E - Atrial fibrillation

Explanation Why

[Atrial fibrillation](#) is a very common cause of [thromboembolic stroke](#) in the elderly and those with [heart](#) disease (e.g., [heart failure](#)). This patient is at low risk for [atrial fibrillation](#) because of her age and lack of [heart](#) disease. Also, she was in normal [heart](#) rhythm on examination, which decreases the likelihood of [atrial fibrillation](#) but does not rule it out.

F - Carotid artery dissection

Explanation Why

[Carotid dissection](#) can occur at any age and is a common cause of [stroke](#) in young patients (~ 20% of cases). [Cervical artery dissection](#) classically presents with [headache](#) and neck [pain](#) following exertion (e.g., exercise, sexual intercourse, or even sneezing). This patient's lack of [headache](#) or neck [pain](#) in the presence of symptoms suggesting [DVT](#) make another etiology more likely.

Question # 11

A previously healthy 20-year-old man is brought to the emergency department 15 minutes after collapsing while playing basketball. He has no history of serious illness. On arrival, there is no palpable pulse or respiratory effort observed. He is declared dead. The family agrees to an autopsy. Cardiac workup prior to this patient's death would most likely have shown which of the following findings?

	Answer	Image
A	Systolic anterior motion of the mitral valve	
B	Narrowing of the left main coronary artery	
C	Bicuspid aortic valve	

	Answer	Image
D	Ventricular septum defect	The image displays a comparison of ECG and Phonocardiogram (PCG) waveforms. The top trace is the ECG (blue) and the bottom trace is the PCG (red). The background is divided into Diastole (grey) and Systole (white) phases. The ECG shows normal QRS complexes. The PCG shows a normal aortic valve closure sound (S1) and aortic valve opening sound (S2). The PCG also shows a high-frequency, high-amplitude murmur (S3) during the Diastole phase, which is characteristic of a VSD.
E	Symmetric thickening of the left ventricle	
F	Increased mean pressure gradient across the aortic valve	

Hint

This patient likely had a left ventricular outflow tract obstruction and fatal arrhythmia that was exacerbated by exercise.

Correct Answer


A - Systolic anterior motion of the mitral valve

Explanation Why

The most likely cause of this young patient's death is [sudden cardiac arrest](#) due to [hypertrophic obstructive cardiomyopathy \(HOCM\)](#). The diagnosis can be confirmed by [echocardiography](#), which typically shows [systolic anterior motion of the mitral valve](#), asymmetrical septal [hypertrophy](#), and a septal wall thickness of >15 mm. Doppler studies show increased left ventricular outflow tract pressure gradients.

B - Narrowing of the left main coronary artery

Image




Explanation Why

Narrowing of the [left main coronary artery](#) is seen in [coronary artery disease](#) (CAD). CAD is a chronic disease process that typically manifests in older individuals with exertional [angina](#) or [dyspnea](#). Patients with severe CAD can present with [cardiac arrest](#) due to [myocardial ischemia](#). However, this patient has no known history of [angina](#), and the sudden collapse of a young patient with no other history of symptoms makes a diagnosis of premature, severe CAD very unlikely.

C - Bicuspid aortic valve

Image



Explanation Why

[Bicuspid aortic valve](#) is the most common cause of [aortic stenosis](#) (AS) in patients younger than 70. AS usually manifests with exertional dyspnea, [syncope](#), and, in severe cases, can lead to [sudden cardiac death](#). However, without any prior history of symptoms in this patient, [sudden cardiac death](#) secondary to severe [aortic stenosis](#) is very unlikely.

D - Ventricular septum defect

Image



Explanation Why

[Ventricular septal defect \(VSD\)](#), like any structural [heart](#) defect, increases the risk of [sudden cardiac death](#). While small defects are usually asymptomatic (and mostly benign), larger defects can lead to myocardial remodeling, [Eisenmenger syndrome](#), and [heart failure](#). The absence of any symptoms or medical problems prior to this patient's collapse makes [VSD](#) an unlikely cause of death in this case.

E - Symmetric thickening of the left ventricle

Explanation Why

Symmetric thickening of the [left ventricle](#) is seen in [restrictive cardiomyopathy \(RCM\)](#). It is often the result of chronic illness that leads to decreased elasticity of the [heart](#) (e.g., [amyloidosis](#), [hemosiderosis](#)), which, in the long run, can lead to [sudden cardiac death](#). However, prior to such

an event, patients typically present with a long-standing history of [palpitations](#), [dyspnea](#), and [syncope](#). [Sudden cardiac arrest](#) in a young patient is better explained by a condition that classically manifests with asymmetrical septal [hypertrophy](#) and can become symptomatic at any age.

F - Increased mean pressure gradient across the aortic valve

Explanation Why

An increased mean pressure gradient across the [aortic valve](#) is seen in [aortic stenosis](#) (AS), which can manifest with [syncope](#) and cardiac death. However, patients usually present at older ages with exertional dyspnea and, depending on the severity of the stenosis, [syncope](#). The age of presentation is typically older than 70 years in degenerative calcific [aortic stenosis](#) and older than 40 in congenital [bicuspid aortic valves](#). Without any prior symptomatic history, [sudden cardiac death](#) secondary to severe [aortic stenosis](#) is very unlikely. [Sudden cardiac arrest](#) in a young patient is better explained by a condition that classically manifests with asymmetrical septal [hypertrophy](#) causing increased left ventricular outflow tract pressure gradients below (not across) the level of the [aortic valve](#).

Question # 12

An otherwise healthy 15-year-old boy comes to the physician for a routine health maintenance examination. He feels well and is doing well in school. He has no history of serious illness. Vital signs are within normal limits. The lungs are clear to auscultation. Cardiac auscultation shows no murmur, but a wide-split S₂ that does not change with respiration. If left untreated, this patient is at increased risk for which of the following complications?

	Answer	Image
A	Sudden cardiac death	
B	Cerebral aneurysm	
C	Left ventricular hypertrophy	
D	Paradoxical embolism	
E	Infective endocarditis	

Hint

This patient's auscultation findings are likely caused by an atrial septal defect (ASD).

Correct Answer

A - Sudden cardiac death

Explanation Why

[Hypertrophic cardiomyopathy \(HCM\)](#) typically manifests with a [systolic ejection murmur](#) caused by left ventricular outflow tract obstruction (LVOT). [Hypertrophic myocardium](#), [fibrosis](#), and disarrangement of cardiac [myocytes](#) predispose to potentially lethal [ventricular arrhythmias](#), which can lead to [sudden cardiac death](#), usually during or after physical activity. If severe enough (and then likely symptomatic), [HCM](#) can also lead to the [paradoxical splitting of S₂](#), in which the [aortic valve](#) closes later than the [pulmonary valve](#) (P₂, then A₂). In an otherwise healthy patient, there is a more likely anatomic cause that explains the fixed, wide [splitting of S₂](#) in which the [pulmonary valve](#) closes later than the [aortic valve](#) (A₂, then P₂).

B - Cerebral aneurysm

Explanation Why

[Cerebral aneurysms](#) are associated with [coarctation of the aorta](#). Coarctation can manifest with a [systolic ejection murmur](#) over the left paravertebral area due to blood flowing through the [aortic narrowing](#) and/or a [continuous murmur](#) from collateral vessels heard between the shoulder blades. It does not manifest with a fixed, wide [splitting of S₂](#). [Coarctation of the aorta](#) may also be accompanied by other symptoms, such as [hypertension](#), lower extremity [cyanosis](#), lower extremity [claudication](#), cold feet, [dyspnea](#) on exertion, or [chest pain](#).

C - Left ventricular hypertrophy

Explanation Why

[Left ventricular hypertrophy](#) is the often consequence of volume overload or increased pressures in the [left ventricle](#) (e.g., [aortic regurgitation](#), [aortic stenosis](#)). ASD, [PFO](#) (seen here), and [VSD](#) increase the volume in the [right ventricle](#) and subsequently lead to [right ventricular hypertrophy](#). In the case

of large septal defects, untreated chronic volume overload in the right [heart](#) leads to severe [hypertrophy](#) and pressure reversal ([Eisenmenger syndrome](#)), which then shunts deoxygenated blood to the [left ventricle](#), leading to [cyanosis](#).

D - Paradoxical embolism

Explanation Why

[Paradoxical embolism](#) is associated with [atrial septal defects](#) (ASD, [PFO](#)) and [ventricular septal defects](#) ([VSD](#)), which allows right-sided [thromboemboli](#) to enter the left-sided circulation by bypassing the [pulmonary circulation](#). An ASD causes wide [splitting of S₂](#) that is fixed (does not change with respiration) due to increased volume within the [pulmonary circulation](#). A [VSD](#) can manifest with a [holosystolic murmur](#).

E - Infective endocarditis

Explanation Why

Congenital [heart](#) defects (e.g., [ventricular septal defect](#), [pulmonary stenosis](#)) and predamaged [heart valves](#) (e.g., [aortic stenosis](#), [mitral valve regurgitation](#)) can predispose to [infective endocarditis](#). However, this risk is negligible in ASDs and PFOs.

Question # 13

A 65-year-old man comes to the physician for a routine examination. He feels well. His pulse is 80/min and blood pressure is 140/85 mm Hg. Cardiac examination shows a holosystolic murmur in the 4th intercostal space along the left sternal border that gets louder during inspiration. The increase of this patient's murmur is best explained by which of the following hemodynamic changes?

	Answer	Image
A	Increased peripheral vascular resistance	
B	Increased right ventricular stroke volume	
C	Decreased pulmonary vessel capacity	
D	Decreased left ventricular venous return	
E	Increased systemic venous compliance	

Hint

A holosystolic murmur that becomes louder on inspiration hints at a right-sided murmur. The auscultation site suggests tricuspid valve regurgitation.

Correct Answer

A - Increased peripheral vascular resistance

Explanation Why

Increased [peripheral vascular resistance](#) ([afterload](#)) does not occur during inspiration. Asking a patient to squat can effectively increase [afterload](#), though. Increased [afterload](#) leads to enhanced [murmur](#) intensity in [mitral regurgitation](#), [aortic regurgitation](#), and [ventricular septal defect](#). In all these instances, increased [afterload](#) augments the pathological, retrograde blood flow across the [mitral valve](#), [aortic valve](#), or [ventricular septal defect](#), respectively. Increased blood flow is appreciated as a more intense [murmur](#).

B - Increased right ventricular stroke volume

Explanation But

As a diagnostic maneuver, inspiration increases the intensity of all [murmurs](#) involving the right side of the [heart](#). These also include [diastolic murmurs](#) like tricuspid stenosis and pulmonary regurgitation.

Explanation Why

A [holosystolic murmur](#) in the 4th [intercostal space](#) along the left sternal border that gets louder during inspiration is classic for [tricuspid regurgitation](#). [Tricuspid regurgitation](#), unlike other [holosystolic murmurs](#) (e.g., [mitral regurgitation](#) or [ventricular septal defects](#)), increases during inspiration. Inspiration causes a period of negative intrathoracic pressure, allowing for increased [venous return](#) to the [heart](#), ultimately producing increased [right ventricular stroke volume](#). This excess blood volume must be entirely ejected through the [tricuspid valve](#), producing the increased intensity of the [murmur](#). The [left ventricle stroke volume](#) is affected in the opposite manner with inspiration decreasing [venous return](#) from the [pulmonary veins](#), leading to a reduction in [left ventricular stroke volume](#). Consequently, the left-sided [holosystolic murmurs](#) of [mitral regurgitation](#) and [ventricular septal defects](#) do not augment with inspiration.

C - Decreased pulmonary vessel capacity

Explanation Why

During inspiration, the intrathoracic pressure is negative. This reduces the barometric compressive force on the outer surface of the pulmonary vessels, permitting an increase in the vessel capacitance. This increase in vessel capacity is part of the explanation for the augmented [murmur](#) intensity during inspiration.

D - Decreased left ventricular venous return

Explanation Why

Decreased [left ventricular venous return](#) occurs during inspiration but does not impact right-sided [heart murmurs](#) such as [tricuspid regurgitation](#). Left-sided [murmur](#) intensity, such as in [mitral regurgitation](#), is typically unaffected or minimally reduced by this phenomenon.

E - Increased systemic venous compliance

Explanation Why

Increased systemic [venous compliance](#), such as immediately following a change from [supine](#) to standing, leads to decreased [preload](#) as a result of blood pooling in the peripheral venous circulation. The majority of cardiac [murmurs](#) become softer in this scenario as a result of diminished blood flow across valves. The one notable exception is the [murmur](#) of [hypertrophic cardiomyopathy](#), which increases. Altered systemic [venous compliance](#) does not have a role in explaining this patient's examination findings.

Question # 14

A 65-year-old woman comes to the physician because of a 3-month history of intermittent palpitations and shortness of breath. Cardiopulmonary examination shows no other abnormalities. An ECG shows an absence of P waves, an oscillating baseline, and irregular RR intervals at a rate of approximately 95 beats per minute. The difference between atrial and ventricular rates in this patient is most likely due to which of the following?

	Answer	Image
A	Temporary inactivation of Na^+ channels in the AV node	
B	Inhibition of the Na^+/K^+ -ATPase pump in ventricular cells	
C	Transient activation of K^+ current in Purkinje fibers	
D	Stimulation of funny current channels in the SA node	
E	Limited speed of conduction through the left bundle branch	
F	Prolonged influx through voltage-gated Ca^{2+} channels in the bundle of His	

Hint

This patient's findings of absent P waves, oscillating baseline, and irregular RR intervals (indicating variable ventricular contractions) on ECG are diagnostic of atrial fibrillation (A-fib), which commonly presents with intermittent palpitations. While this patient's ventricular rate is ~ 95 beats per minute, her atrial contraction rate is likely > 300 beats per minutes.

Correct Answer

A - Temporary inactivation of Na^+ channels in the AV node

Explanation Why

The temporary inactivation of Na^+ channels in the [AV node](#) after Phase 0 of the [action potential](#) is what leads to a [refractory period](#) of the [AV node](#). A [refractory period](#) is the time it takes for the ventricles to recover and be available to conduct another impulse from the [atria](#). This patient's [atria](#) are contracting at > 300 bpm, but the [AV node refractory period](#) prevents this many impulses from being conducted from the [atria](#) to the ventricles so that the ventricular rate is only 95 bpm.

B - Inhibition of the Na^+/K^+ -ATPase pump in ventricular cells

Explanation Why

Inhibition of the [\$\text{Na}^+/\text{K}^+\$ -ATPase pump](#) is the mechanism of action of the drug [digoxin](#), which increases intracellular sodium levels, decreases activity of the sodium-calcium pump, and increases intracellular calcium (leads to overall increased contractility). It also increases [vagal](#) activity and has several extracardiac effects, but this mechanism does not cause a difference in atrial and ventricular contraction rates or lead to A-fib. In addition, [digoxin](#) is actually a second-line treatment option for patients with A-fib.

C - Transient activation of K^+ current in Purkinje fibers

Explanation Why

The transient activation of K^+ current in [Purkinje fibers](#) does not cause a dissociation between the rates of the [atria](#) and ventricles as in this patient with A-fib. Instead, it causes the overshoot of the [cardiac action potential](#) during Phase 1. This is when K^+ ions exit the cell, leaving the membrane slightly more negative and creating a 'notch' on the waveform before the plateau and [repolarization](#).

D - Stimulation of funny current channels in the SA node

Explanation Why

[Funny current](#) channels set the [pacemaker current](#) of the [SA node](#), which determines the atrial contraction rate. Their stimulation would increase the [heart rate](#) but not lead to a difference in atrial and ventricular contraction rates as in this patient's A-fib.

E - Limited speed of conduction through the left bundle branch

Explanation Why

Limited speed of conduction through the [left bundle branch](#) describes a [left bundle branch block](#). A [bundle branch](#) block is a form of wide-complex [tachyarrhythmia](#), as opposed to A-fib seen in this patient.

F - Prolonged influx through voltage-gated Ca^{2+} channels in the bundle of His

Explanation Why

A prolonged influx through voltage-gated Ca^{2+} channels in the [bundle of His](#) would lead to a wide-complex tachycardia rather than this patient's A-fib. It would not lead to a difference in atrial and ventricular rates.

Question # 15

A 3-year-old girl is brought to the emergency department by her mother 30 minutes after the sudden onset of shortness of breath, dizziness, abdominal pain, and urticaria. The symptoms began 5 minutes after she ate a peanut butter sandwich. Her respirations are 36/min and blood pressure is 84/50 mm Hg. There is stridor on auscultation of the chest. She is intubated and mechanical ventilation is begun. Intravenous fluid resuscitation and the appropriate pharmacotherapy are begun. The drug administered to this patient most likely has which of the following effects on cardiac pacemaker cells during an action potential?

	Answer	Image
A	Increased Ca^{2+} influx in phase 4	
B	Decreased Na^{+} influx in phase 0	
C	Increased Na^{+} influx in phase 3	
D	Decreased Ca^{2+} influx in phase 0	
E	Increased K^{+} efflux in phase 4	
F	Decreased K^{+} efflux during phase 3	


Hint

The onset of shortness of breath, hypotension, abdominal pain, urticaria, and angioedema shortly after ingestion of a commonly allergenic food (peanuts) is consistent with anaphylaxis. Epinephrine is the mainstay of pharmacotherapy for this condition.

Correct Answer

A - Increased Ca^{2+} influx in phase 4

Image



Explanation But

[Parasympathomimetics](#) and adenosine have opposite effects on the cardiac pacemaker AP. These drugs decrease the slope of phase 4 by reducing the depolarizing influx of Ca^{2+} and Na^+ and increasing the efflux of K^+ , thereby lowering the [heart rate](#).

Explanation Why

[Catecholamines](#) like [epinephrine](#) stimulate [beta 1 adrenergic receptors](#) in the [heart](#), thereby increasing Ca^{2+} influx through the sequential opening of [T-type calcium channels](#) and [L-type calcium channels](#) in the [sinoatrial node](#) and [atrioventricular node](#) in the late part of phase 4 of the [pacemaker action potential](#) (AP). [Catecholamines](#) also increase the flow of Na^+ through pacemaker channels ([funny channels](#); I_f). These effects lead to more rapid [depolarization](#), which increases [heart rate](#). Other effects of [epinephrine](#) are mediated through [beta 2 adrenergic receptors](#) (bronchial relaxation) and [alpha 1 adrenergic receptors](#) (vasoconstriction), which ease breathing and stabilize

blood pressure in [anaphylactic shock](#), respectively.

B - Decreased Na^+ influx in phase 0

Explanation Why

Decreased Na^+ influx in phase 0 (upstroke) of the [cardiac action potential](#) would slow [depolarization](#) in the cardiac [myocytes](#). It would not have an effect on pacemaker cells, as they rely on calcium (not sodium) channels during [depolarization](#). Decreasing Na^+ influx is the mechanism of action of class I antiarrhythmic agents, which are not used during [anaphylactic shock](#).

C - Increased Na^+ influx in phase 3

Explanation Why

During phase 3 of the cardiac [pacemaker potential](#) (the [repolarization](#) phase), the voltage-gated Ca^{2+} channels close and the K^+ channels remain open, allowing for K^+ efflux. Na^+ channels are normally inactive during this phase. Increased Na^+ influx during phase 3 slows [repolarization](#) in [cardiac pacemaker cells](#), which would lead to negative [chronotropy](#), the opposite of the desired effect in treatment of [anaphylactic shock](#).

D - Decreased Ca^{2+} influx in phase 0

Explanation Why

At the [threshold potential](#), the opening of additional [L-type \$\text{Ca}^{2+}\$ channels](#) augments the inward current of [calcium ions](#), resulting in the upstroke of phase 0 of the [pacemaker action potential](#). [Calcium channel blockers](#) block [L-type calcium channels](#), resulting in reduced Ca^{2+} influx during the late part of phase 4 and phase 0 of the [pacemaker action potential](#). However, [calcium channel blockers](#) do not have a role in treatment of [anaphylactic shock](#).

E - Increased K^+ efflux in phase 4

Explanation Why

Increased K^+ efflux in phase 4 leads to a decreased rate of spontaneous [depolarization](#) since cardiac [myocytes](#) become more electronegative and therefore require more cation influx to depolarize.

[Acetylcholine](#) and adenosine both increase K^+ efflux in phase 4 of the [action potential](#). However, neither of these drugs are used in the treatment of [anaphylactic shock](#).


F - Decreased K^+ efflux during phase 3

Explanation Why

Decreased K^+ efflux during phase 3 of the [cardiac action potential](#) leads to prolonged [repolarization](#), extending the [refractory period](#). This is the mechanism of action of [class III antiarrhythmic drugs](#), which are not used to treat [anaphylactic shock](#).

Question # 16

A 56-year-old man comes to the emergency department for chest pain that started 1 hour ago. He has a history of hypertension, hyperlipidemia, and type 2 diabetes mellitus. He appears diaphoretic and distressed. An ECG shows P waves independent of QRS complexes. The QRS interval is 0.11 seconds with normal morphology. Which of the following is the most likely trigger of this patient's ventricular contractions?

	Answer	Image
A	Sinoatrial node	
B	Bachmann bundle	
C	Bundle of His	
D	Purkinje fibers	
E	Left bundle branch	
F	Ectopic atrial focus	

Hint

An ECG that shows P waves independent of QRS complexes indicates complete heart block; the atria and ventricles are contracting independently of each other. The normally-shaped QRS complexes indicate that there is an ectopic pacemaker elsewhere along the conduction pathway.

Correct Answer

A - Sinoatrial node

Explanation Why

An electrical rhythm generated from the [SA node](#) (the natural pacemaker of the [heart](#)) results in normal narrow [QRS complexes](#), as seen in this [ECG](#). However, the [P waves](#) will always precede the [QRS complex](#) at regular, fixed intervals, which is not seen here.


B - Bachmann bundle

Explanation Why

The Bachmann bundle conducts electrical impulses from the [SA node](#) to the [left atrium](#); it is not part of the ventricular portion of the cardiac conduction pathway. The Bachmann bundle is not known to serve as an [ectopic pacemaker](#); however, electric activity originating there would theoretically pass through the [SA node](#) to the [AV node](#) and result in [P waves](#) that precede the [QRS complex](#) at regular, fixed intervals, which is not seen here.

C - Bundle of His

Image



Explanation Why

In [complete heart block](#), electrical impulses from the [SA node](#) are not conducted to the ventricles. [Myocytes](#) of the cardiac conduction pathway have the potential to spontaneously depolarize and behave as pacemakers when they do not receive signals from the [SA node](#), which results in escape rhythms. Depending on the shape of the [QRS complexes](#) of the escape rhythms, the site of these [ectopic pacemakers](#) can be determined. Normal, narrow [QRS complexes](#) as seen on this [ECG](#) are triggered by impulses anywhere from the [SA node](#) up to the [bundle of His](#) (i.e., the block is at or [proximal](#) to the [bundle of His](#)).

D - Purkinje fibers

Explanation Why

An [ectopic pacemaker](#) located in the [Purkinje fibers](#) results in wide [QRS complexes](#) because the rate of conduction of the electrical impulse to the ventricles is much slower than when it originates from the [SA node](#).

E - Left bundle branch

Explanation Why

An [ectopic pacemaker](#) located in either the left or the right [bundle branch](#) of the cardiac conduction pathway results in wide [QRS complexes](#) because the rate of conduction of the electrical impulse to the ventricles is much slower than when it originates from the [SA node](#).

F - Ectopic atrial focus

Explanation Why

An atrial [ectopic pacemaker](#) generates an escape rhythm characterized by an abnormal [P wave](#) that precedes every [QRS complex](#). The [QRS complexes](#) would be normal because the atrioventricular conduction is normal (i.e., via the His-Purkinje pathway). This patient's [ECG](#) shows normal [P waves](#) independent of the [QRS complexes](#).

Question # 17

A 62-year-old man with a history of coronary artery disease comes to the emergency department with substernal chest pain for several hours. An ECG shows no abnormalities. Troponin T test results are negative. The patient is admitted to the hospital and treated with intravenous nitroglycerin, with an initial resolution of his symptoms. After 6 hours of continuous infusion of nitroglycerin, he reports increasing chest pain. The underlying cause of this patient's recurrent symptoms is most likely to also occur in treatment with which of the following drugs?

	Answer	Image
A	Imatinib	
B	Levodopa	
C	Alprazolam	
D	Phenylephrine	
E	Methicillin	
F	Hydrocodone	

Hint

Continuous or repeated dosing of nitroglycerine leads to a depletion of nitric oxide in endothelial cells, leading to a decreased vasodilatory effect. This rapidly declining responsiveness due to the depletion of the body's endogenous mediators is known as tachyphylaxis.

Correct Answer

A - Imatinib

Explanation Why

[Imatinib](#) is a [BCR-ABL tyrosine kinase inhibitor](#) used in the treatment of [chronic myeloid leukemia](#). As with many other [chemotherapy drugs](#), patients may develop resistance to [imatinib](#). This process, however, is due to sporadic mutations in the [BCR-ABL fusion gene](#) of the [CML](#) cancer cells and not due to [tachyphylaxis](#).

B - Levodopa

Explanation Why

Long-term use of [levodopa](#) can result in its decreased effectiveness, as seen in the [end-of-dose akinesia](#) in patients with [Parkinson disease](#). This decreasing effect, however, is not due to [tachyphylaxis](#). Instead, it is believed to be due to the progression of the disease and/or the neurotoxic effects of [dopamine](#).

C - Alprazolam

Explanation Why

[Alprazolam](#), like other [benzodiazepines](#), can have decreasing effectiveness with chronic use. This effect, however, occurs via the long-term process of [drug tolerance](#) which likely involves the downregulation of [GABA](#) receptors, not via [tachyphylaxis](#), which is seen here.

D - Phenylephrine

Explanation Why

Repeated use of [phenylephrine](#) (an alpha-1 agonist) can cause downregulation of [alpha-1 receptors](#), resulting in [tachyphylaxis](#) that cannot be overcome by increasing the drug dose. This is seen in [rhinitis medicamentosa](#), a condition in which excessive use of [phenylephrine](#) nasal spray leads to rebound nasal congestion as the medication wears off. Other substances subject to [tachyphylaxis](#) include [niacin](#), [LSD](#), and [MDMA](#).

E - Methicillin

Explanation Why

Decreasing effectiveness of methicillin can be seen in the case of methicillin-resistant [Staphylococcus aureus](#). This process of [antibiotic](#) resistance, however, is caused by acquired mutations in [penicillin-binding proteins](#) rather than by [tachyphylaxis](#).

F - Hydrocodone

Explanation Why

Chronic use of [opiates](#) such as hydrocodone can reduce the therapeutic effect of a standard dose. However, the underlying mechanism here is not [tachyphylaxis](#) but [drug tolerance](#), which is a long-term process of desensitization and downregulation of [opiate](#) receptors.

Question # 18

A 74-year-old man comes to the physician because of a 2-month history of shortness of breath and leg swelling. He has trouble climbing one flight of stairs due to dyspnea. He has hypertension and was diagnosed with multiple myeloma 6 months ago. Current medications include hydrochlorothiazide. Cardiopulmonary examination shows jugular venous distention, bibasilar crackles, and pitting edema in the lower extremities. Transthoracic echocardiography shows a thickened myocardium with elevated diastolic filling pressures. Which of the following is the most likely underlying mechanism of this patient's cardiomyopathy?

	Answer	Image
A	Chronic vasoconstriction	
B	Myosin gene mutation	
C	Amyloid deposition	
D	Chronic hypercalcemia	
E	Iron deposition	The diagram illustrates the pathophysiology of hemochromatosis. On the left, 'Normal' iron metabolism is shown where the liver produces hepcidin (regulated by HFE protein) to bind and regulate ferroportin on macrophages and the liver, controlling iron export and maintaining normal plasma iron levels. On the right, 'Hemochromatosis' is shown where an HFE mutation (H63D or H63D) leads to a modified HFE protein, resulting in decreased hepcidin. This causes increased iron absorption from the gut and increased iron release from macrophages and the liver, leading to iron overload. The overload results in increased iron, ferritin, and transferrin saturation, which causes accumulation in organs, leading to free radical injury and subsequent liver cirrhosis, diabetes, and cardiomyopathy.

Hint

Multiple myeloma is characterized by the production of monoclonal immunoglobulins.

Correct Answer

A - Chronic vasoconstriction

Explanation Why

Chronic vasoconstriction can develop in long-term [cocaine](#) use and is due to alpha-1 receptor stimulation. [Cocaine](#) is toxic to the [myocardium](#), causes [myocardial](#) thinning, and can lead to [dilated cardiomyopathy](#) (usually seen on [echocardiography](#)). However, this patient's [TTE](#) did not support the diagnosis of [dilated cardiomyopathy](#) and he has no history of [substance abuse](#).

B - Myosin gene mutation

Explanation Why

[Myosin gene](#) mutation is an [autosomal dominant](#) mutation seen in patients with [hypertrophic cardiomyopathy \(HCM\)](#), and the obstructive type of this disease can cause [congestive heart failure](#). [Multiple myeloma](#), however, is not associated with [myosin gene](#) mutation. Moreover, [HCM](#) is usually seen in young male athletes with a positive [family history](#).

C - Amyloid deposition

Explanation Why

[Amyloid plaque](#) is a common sequela of [multiple myeloma](#) and forms as a result of excess production of monoclonal [immunoglobulins](#). [Immunoglobulins](#) can deposit in any organ, including the [heart](#), in which case they cause a specific type of [restrictive cardiomyopathy](#) known as [infiltrative cardiomyopathy](#). Thickening of the [myocardium](#) results in [diastolic dysfunction](#) and back-flow of blood causes [shortness of breath](#), [jugular venous distention](#), and [pulmonary edema](#), which are all seen here.


D - Chronic hypercalcemia

Explanation Why

Chronic [hypercalcemia](#) is often seen in patients with [multiple myeloma](#). [Hypercalcemia](#) can have a wide range of findings, including [constipation](#), bone pains, and psychiatric [mood disorders](#). [Cardiomyopathy](#), however, is not caused by [hypercalcemia](#).

E - Iron deposition

Image




Explanation Why

[Iron](#) deposition can lead to [restrictive cardiomyopathy](#) (although [dilated cardiomyopathy](#) is more common). [Hemochromatosis](#) is one condition that is associated with a high [iron](#) state. Patients with this condition can develop [cardiomyopathy](#) as a result of [iron](#) deposition, which presents with symptoms similar to those seen in this patient, including [shortness of breath](#) and [JVD](#). However,

other features typically associated with [hemochromatosis](#) (e.g., bronzed [skin](#), [diabetes](#), arthralgia) would be expected.

Question # 19

A 71-year-old man with hypertension comes to the physician for a follow-up examination. Cardiovascular exam shows the point of maximal impulse to be in the mid-axillary line. A transthoracic echocardiogram shows concentric left ventricular hypertrophy with a normal right ventricle. Which of the following is the most likely underlying mechanism of this patient's ventricular hypertrophy?

	Answer	Image
A	Accumulation of glycogen	
B	Accumulation of protein fibrils	
C	Deposition of endomyocardial collagen	
D	Accumulation of sarcomeres in parallel	 The diagram illustrates the structure of a sarcomere and a myofibril. The top part shows a single sarcomere with labels for the I band, A band, H zone, Z line, and M line. The bottom part shows a myofibril composed of multiple sarcomeres, with labels for the Z line, M line, Titin, and α-actinin.
E	Infiltration of T lymphocytes	

Hint

This patient's left ventricular hypertrophy (LVH) is due to a chronic increase in afterload.

Correct Answer

A - Accumulation of glycogen

Explanation Why

Accumulation of glycogen is seen in [type 2 glycogen storage disease \(Pompe disease\)](#). Clinical features include generalized muscle weakness, [hyperlipidemia](#), [hypoglycemia](#), and [ketosis](#), none of which are present in this patient. In addition, [Pompe disease](#) usually presents in [newborns](#), and death occurs within weeks of [birth](#).

B - Accumulation of protein fibrils

Explanation Why

The accumulation of protein fibrils within the [myocardium](#) is seen in [amyloidosis](#). [Amyloidosis](#) is caused by misfolded [proteins](#), which are most commonly derived from [immunoglobulin light chain](#), [amyloid precursor protein](#), and [serum amyloid A](#). [Amyloidosis](#) results in thickening of the [myocardium](#), as in this patient. However, the [myocardial](#) thickening in [amyloidosis](#) is usually biventricular.


C - Deposition of endomyocardial collagen

Explanation Why

Deposition of endomyocardial [collagen](#) is seen in [restrictive cardiomyopathy](#). It is caused by [proliferation](#) of [connective tissue](#), which results in decreased elasticity of the [myocardium](#) and severe [diastolic dysfunction](#). Although [restrictive cardiomyopathy](#) can cause atrial enlargement, it is commonly due to dilation, not [hypertrophy](#).

D - Accumulation of sarcomeres in parallel

Image



Explanation Why

This patient's [LVH](#) is due to accumulation of [sarcomeres](#) in parallel, likely caused by chronic [hypertension](#) (increased [afterload](#)). Chronic [hypertension](#) increases [myocardial](#) wall tension, which leads to changes in [gene expression](#) and subsequent duplication of [sarcomeres](#) in parallel and [diastolic dysfunction](#). In contrast, [left ventricular](#) dilation is caused by the accumulation of [sarcomeres](#) in series, which results in [systolic dysfunction](#).

E - Infiltration of T lymphocytes

Explanation Why

Infiltration of [T lymphocytes](#) is seen in acute [myocarditis](#). Clinical features of [myocarditis](#) include [flu-like symptoms](#), [dyspnea](#), [chest pain](#), [arrhythmias](#), [CHF](#), and [cardiogenic shock](#), all of which this

patient lacks. In addition, [echocardiography](#) in [myocarditis](#) is usually unremarkable but can show ventricular dilation, diffuse [hypokinesia](#), reduced [ejection fraction](#), impaired contractility, and regional wall motion abnormalities.

Question # 20

A 57-year-old man comes to the emergency department with fatigue and palpitations for several weeks. An ECG shows atrial fibrillation. Echocardiography shows thrombus formation in the left atrium. Which of the following organs is most likely to continue to function in the case of an embolic event?

	Answer	Image
A	Spleen	
B	Brain	
C	Kidney	
D	Liver	
E	Colon	

Hint

In the case of arterial emboli, most organs will infarct and cease to function. However, a dual blood supply may allow tissue to remain oxygenated and preserve function.

Correct Answer

A - Spleen

Explanation Why

The [spleen](#) primarily receives blood via the [splenic artery](#). Though splenic emboli are rare and usually quiescent clinically, the [spleen](#) is more susceptible to [infarction](#) and loss of function from emboli than an organ with dual blood supply.

B - Brain

Explanation Why

The [central nervous system](#) is especially vulnerable to [ischemia](#), from embolic occlusion or otherwise. This is due to its high metabolic rate and glucose consumption. In total, ~ 80% of embolism-related deaths in [atrial fibrillation](#) are from [stroke](#).

C - Kidney

Explanation Why

Each [kidney](#) has a single blood supply (e.g., respective [renal artery](#)). This makes the [kidneys](#) more susceptible to [infarction](#) than an organ with dual blood supply, particularly in the straight portion of the [proximal convoluted tubule](#) and [ascending loop of Henle](#) (due to their high energy demand and proximity to the corticomedullary junction, which has low blood flow). While A-fib is associated with a low [incidence](#) (0.01%) of renal [thromboembolism](#), 25% of patients who experienced an event had persistent [renal failure](#).

D - Liver

Explanation Why

The [liver](#) benefits from dual blood supply. The [hepatic portal vein](#) supplies ~ 75% of the [liver](#)'s blood, and the [hepatic artery proper](#) contributes the rest. The [hepatic portal vein](#) carries venous blood from the [spleen](#) and lower [gastrointestinal tract](#), while the hepatic [arteries](#) are branches of the [common hepatic artery](#) arising from the [celiac trunk](#). Oxygen is provided from both sources with about half of the [liver](#)'s oxygen demand met by the [hepatic portal vein](#). The dual blood and oxygen supply protects the [liver](#) against potential [infarct](#).

E - Colon

Explanation Why

While thrombosis in the bowel is usually less acute due to collateral circulation, embolic obstruction of the superior or [inferior mesenteric arteries](#) can still cause acute [ischemia](#) of the [colon](#), particularly in [watershed areas](#). Acute thromboembolic [mesenteric ischemia](#) due to A-fib has a low [incidence](#) but high [lethality](#) (~ 70%).

Question # 21

A 2-year-old boy with a history of recurrent respiratory infections is brought to the physician for a follow-up examination. His height and weight are both at the 20th percentile. Crackles are heard in both lower lung fields. Cardiac auscultation shows a grade 3/6 holosystolic murmur over the left lower sternal border and a diastolic rumble heard best at the apex. If left untreated, this patient is most likely to develop which of the following?

	Answer	Image
A	Myocardial ischemia	
B	Thrombocytosis	
C	Secondary hypertension	
D	Aortic dissection	
E	Digital clubbing	
F	Chronic kidney disease	

Hint

The presence of a holosystolic murmur (indicating flow through the defect) and a diastolic rumble (increased flow through the mitral valve during diastole) suggests a ventricular septal defect (VSD), the most common type of congenital heart defect. VSDs are associated with frequent respiratory infections secondary to congestion of the pulmonary vessels.

Correct Answer

A - Myocardial ischemia

Explanation Why

Congenital [heart](#) defects associated with an increased risk of [myocardial ischemia](#) are typically those that involve [coronary artery](#) abnormalities (e.g., anomalous course or origin, involving [fistulas](#)) or result in decreased coronary perfusion (e.g., in a large [PDA](#), the retrograde blood flow to the [left ventricle](#) can decrease [diastolic](#) pressure in the [proximal](#) aorta and thereby decrease coronary perfusion). Although a long-standing, uncorrected [VSD](#) can also lead to decreased blood pressure, there is no evidence that it increases the risk of [myocardial ischemia](#).

B - Thrombocytosis

Explanation Why

[Acyanotic heart defects](#), such as [VSDs](#), are not a known cause of increased [platelet counts](#). A long-standing, uncorrected [VSD](#) will eventually result in [Eisenmenger syndrome](#) with subsequent shunt reversal. In this case, [thrombocytopenia](#), not [thrombocytosis](#) may occur. [Thrombocytopenia](#) due to [Eisenmenger syndrome](#) is thought to result from [megakaryocytes](#) bypassing the pulmonary vasculature where they would normally be fragmented into [platelets](#).

C - Secondary hypertension

Explanation Why

[Secondary hypertension](#) can result from congenital [heart](#) defects such as [coarctation of the aorta](#). All aortic branches [proximal](#) to the coarctation receive increased blood flow, ultimately resulting in [intimal-medial](#) thickening and increased vascular stiffness. In a 2-year-old child, symptomatic coarctation would most likely present with [chest pain](#), cold feet, and lower-extremity [claudication](#) on physical exertion, none of which are seen in this patient. Furthermore, auscultation would reveal a [systolic ejection murmur](#) over the left paravertebral region and/or a [continuous murmur](#) below the left clavicle and between the shoulder blades.

D - Aortic dissection

Explanation Why

[Aortic dissection](#) is a dangerous complication of untreated [coarctation of the aorta](#) in adults. Chronic [hypertension proximal](#) to the area of coarctation increases intraluminal pressure and subsequently weakens the arterial wall, potentially resulting in [aortic dissection](#). Symptomatic coarctation in a 2-year-old child would most likely present with [chest pain](#), cold feet, and lower-extremity [claudication](#) on physical exertion, none of which are seen in this patient. Furthermore, auscultation would reveal a [systolic ejection murmur](#) over the left paravertebral region and/or a [continuous murmur](#) below the left clavicle and between the shoulder blades.

E - Digital clubbing

Explanation But

Chronic tissue [hypoxia](#) and [megakaryocyte](#) fragmentation are primary factors involved in the pathogenesis of [digital clubbing](#) in [cyanotic heart disease](#) (e.g., untreated [VSD](#)). In [Eisenmenger syndrome](#) (a complication of [VSDs](#)), blood can completely bypass the pulmonary vasculature, allowing whole [megakaryocytes](#) into the [systemic circulation](#). Once these intact cells reach [distal capillaries](#) in the hands and feet, they impact and subsequently fragment. This fragmentation causes the release of [platelet-derived growth factor \(PDGF\)](#) and [vascular endothelial growth factor \(VEGF\)](#), inducing [angiogenesis](#) and local mesenchymal growth and [proliferation](#), which clinically manifests as [digital clubbing](#).

Explanation Why

[Digital clubbing](#) is a sign of chronic [hypoxemia](#), which is a complication of an untreated [ventricular septal defect \(VSD\)](#). [VSDs](#) are [acyanotic congenital heart defects](#) that manifest as a [left-to-right shunt](#). An untreated [VSD](#) can lead to chronic volume overload of the [right ventricle \(RV\)](#) and subsequently increased blood flow through the pulmonary vasculature. This increased flow causes irreversible remodeling of the small pulmonary vessels, leading to increased [pulmonary vascular resistance](#) and RV [hypertrophy](#). RV pressure increases until it exceeds [left ventricular \(LV\)](#) pressure, at which point the shunt is reversed. This shunt reversal is known as [Eisenmenger syndrome](#). [Cyanosis](#) and consequently, chronic tissue [hypoxia](#), leads to the development of [digital clubbing](#).

F - Chronic kidney disease

Explanation Why

[Chronic kidney disease](#) ([CKD](#)) can result from a variety of conditions, including advanced [left heart failure](#). In individuals with low [cardiac output](#), advanced [left heart failure](#) may lead to decreased renal perfusion, decreased venous congestion, and eventually [CKD](#). Chronic, untreated [VSD](#), which is seen in this patient, causes constant volume overload of the [right ventricle](#) (RV) and subsequent RV damage. Advanced stages of [RV failure](#) manifest with [hepatomegaly](#) and peripheral [edema](#).

Question # 22

A 53-year-old man is brought to the emergency department because of wheezing and shortness of breath that began 1 hour after he took a new medication. Earlier in the day he was diagnosed with stable angina pectoris and prescribed a drug that irreversibly inhibits cyclooxygenase-1 and 2. He has chronic rhinosinusitis and asthma treated with inhaled β -adrenergic agonists and corticosteroids. His respirations are 26/min. Examination shows multiple small, erythematous nasal mucosal lesions. After the patient is stabilized, therapy for primary prevention of coronary artery disease should be switched to a drug with which of the following mechanisms of action?

	Answer	Image
A	Inhibition of vitamin K epoxide reductase	
B	Blockage of P2Y ₁₂ component of ADP receptors	
C	Direct inhibition of Factor Xa	
D	Sequestration of Ca ²⁺ ions	
E	Potentiation of antithrombin III	

Hint

The combination of asthma, aspirin sensitivity, and probable nasal polyposis (multiple small, erythematous nasal mucosal lesions) is known as Samter's triad. The patient develops signs of acute asthmatic attack (wheezing, dyspnea) shortly after initiating therapy with aspirin (an irreversible COX-1 and 2 inhibitor). The patient should be switched to another agent that is used along with aspirin prior to percutaneous coronary intervention.

Correct Answer

A - Inhibition of vitamin K epoxide reductase

Explanation Why

Competitive inhibition of [vitamin K epoxide reductase](#) is the mechanism of action of [warfarin](#). [Warfarin](#) is used for the treatment and prevention of thromboembolic disorders ([venous thromboembolism](#), [pulmonary embolism](#)) and embolic complications arising from [atrial fibrillation](#). [Warfarin](#) is not used for prevention of [coronary heart disease](#).

B - Blockage of P2Y₁₂ component of ADP receptors

Explanation Why

[P2Y₁₂ receptor antagonists](#) (e.g., [clopidogrel](#)) irreversibly inhibit [platelet aggregation](#) by preventing the ADP-mediated formation of the [GpIIb/IIIa receptor](#) complex. [Clopidogrel](#) is used for treatment of [stable angina pectoris](#) in patients with [aspirin allergy](#) as well as adjunctive therapy in acute coronary syndromes. Other indications include [secondary prevention](#) of recent [stroke](#), recent [myocardial infarction](#), and [peripheral arterial disease](#).

C - Direct inhibition of Factor Xa

Explanation Why

[Factor Xa inhibitors](#) (e.g., [apixaban](#), [rivaroxaban](#), [edoxaban](#)) are used for the prophylaxis and treatment of [venous thromboembolism](#), [pulmonary embolism](#) and non-valvular atrial fibrillation. Primary and [secondary prevention](#) of ischemic [heart](#) disease does not include the use of [factor Xa inhibitors](#) or any other new [oral anticoagulants](#) (NOACs).

D - Sequestration of Ca²⁺ ions

Explanation Why

[Citrate](#) is a calcium-binding salt with [anticoagulant](#) properties that is used for the preservation of blood-derived products. [Citrate](#) is also used for anticoagulation in patients undergoing dialysis and for the prevention of recurrent urinary calcium stones. [Citrate](#) is not used for the treatment or prevention of acute or stable coronary syndromes.

E - Potentiation of antithrombin III

Explanation Why

Potentiation of [antithrombin](#) (an endogenous [anticoagulant](#)) indirectly inhibits [thrombin](#) and [factor Xa](#) and is the mechanism of action for unfractionated [heparin](#) (UFH). [UFH](#) is used for anticoagulation in patients with [unstable angina](#) (UA), [ST elevation myocardial infarction](#) (STEMI), and non-[ST elevation myocardial infarction](#) (NSTEMI). However, the mechanism of this patient's stable [angina pectoris](#) involves a mismatch between the [myocardial](#) oxygen supply and myocardial oxygen demand, secondary to a fixed atherosclerotic occlusion without activating the [coagulation cascade](#). Thus, [anticoagulant](#) agents have no therapeutic effect on [stable angina pectoris](#).

Question # 23

A 70-year-old man comes to the physician for a follow-up examination of diffuse exertional chest pain which he has successfully been treating with sublingual nitroglycerin for the past year. The patient has been taking lisinopril daily for essential hypertension. His pulse is 75/min and regular, and blood pressure is 155/90 mm Hg. Cardiac and pulmonary examination show no abnormalities; there is no peripheral edema. A decrease of which of the following is the most likely explanation for the improvement of this patient's chest pain?

	Answer	Image
A	Ventricular compliance	
B	Peripheral arterial resistance	
C	Electrical conduction speed	
D	Venous pooling	
E	Coronary arterial resistance	
F	End-diastolic pressure	The diagram illustrates the mechanism of nitroglycerin-induced vasodilation. In the endothelium, stimuli like ACh, histamine, shearing forces, pulsatile strain, and high intracellular Ca^{2+} activate eNOS, which converts L-arginine to NO and L-citrulline. Nitrates also contribute to NO production. NO diffuses into the smooth muscle cell, where it activates guanylyl cyclase, converting GTP to cGMP. cGMP then activates protein kinase G, which inhibits phospholipase C and the sarcoplasmic reticulum Ca^{2+} release channel. This leads to a decrease in intracellular Ca^{2+}, which in turn inhibits the myosin light chain kinase (MLCK) pathway, resulting in smooth muscle relaxation and vasodilation. The diagram also shows that PDE5 inhibitors block the breakdown of cGMP by PDE5, thereby enhancing its effects. A legend indicates: red for effects of cGMP, yellow for breakdown of cGMP, and green for physiological NO production.

Hint

Exertional chest pain in this patient is likely caused by angina pectoris, a chest pain that occurs if myocardial oxygen demand cannot be met due to coronary heart disease. Nitroglycerin dilates smooth muscle cells in both arteries and veins, which in turn reduces myocardial preload, decreases myocardial oxygen demand, and subsequently provides symptom relief.

Correct Answer

A - Ventricular compliance

Explanation Why

Ventricular compliance describes the volume/fluid moved for a given change in pressure ($C = \Delta V / \Delta P$). It thus decreases in ventricular [hypertrophy](#) (increased pressure, with no change in volume moved) or any type of [restrictive cardiomyopathy](#) (pressure stays the same, volume decreases). [Nitroglycerin](#) actually improves ventricular compliance by reduction of [myocardial preload](#) and its effects.

B - Peripheral arterial resistance

Explanation Why

The decrease in peripheral arterial resistance is achieved through arterial [vasodilation](#), which is one of the effects of [nitroglycerin](#). It results in decreased [afterload](#), which in turn decreases myocardial oxygen demand to a certain extent. However, there is another, much more prominent [nitroglycerin](#)-mediated mechanism that is more likely to reduce this patient's symptoms. Arterial [vasodilation](#) is the primary effect of [calcium channel blockers](#), not that of [nitroglycerin](#).

C - Electrical conduction speed

Explanation Why

[Nitroglycerin](#) has no effect on electrical conduction. Reducing the electrical conduction speed, and thus [heart rate](#), is a mechanism to achieve a long-term reduction of myocardial oxygen demand. It can be achieved by various drugs, [metoprolol](#) being one of the most common ones. In later stages that are refractory to common treatment, [ranolazine](#) can be added to target late inward sodium channels within [myocardial](#) tissue and provide relief in some cases of chronic [angina](#).

D - Venous pooling

Explanation Why

Decreased venous pooling can result from the use of compression stockings or [sympathomimetic drugs](#) that cause vasoconstriction of arteriolar and venous vasculature. This increases [venous return](#) and therefore cardiac [preload](#), which in turn increases [myocardial](#) wall tension and reduces [myocardial](#) perfusion. [Nitroglycerin](#) has the opposite effect: [Smooth muscle](#) relaxation, mainly of the venous vessels, leads to venodilation and subsequent increased blood pooling in the venous system.

E - Coronary arterial resistance

Explanation But


[Nitrates](#) are also used in the treatment of [coronary artery](#) spasm (via muscle relaxation) in [vasospastic angina](#).

Explanation Why

[Nitroglycerin](#) does cause coronary arterial [vasodilation](#), in addition to [vasodilation](#) throughout the body. However, [nitroglycerin](#) predominantly targets the venous system and coronary dilation is not the primary mechanism of improving [myocardial](#) perfusion. In [angina](#), the [coronary arteries](#) are typically in a chronic state of dilation in an attempt to preserve perfusion of the [myocardial](#) tissue (= decreased coronary reserve). In these cases, [nitroglycerin](#) provides no additional increase in blood flow to the [coronary arteries](#) but can still achieve symptom relief via a different mechanism.

F - End-diastolic pressure

Image



Explanation But

Life-threatening [hypotension](#) may occur if [nitroglycerin](#) is taken within 24 hours of a [PDE-5 inhibitor](#) such as [sildenafil](#).

Explanation Why

The administration of [nitroglycerin](#) causes [smooth muscle](#) relaxation in the vascular system, predominantly in the venous system. It thus increases venous pooling which subsequently decreases [myocardial preload](#). This results in a decreased end-[diastolic](#) pressure, which decreases [myocardial](#) wall tension and thus improves [myocardial](#) perfusion while decreasing myocardial oxygen demand. The shift towards a more favorable [ratio](#) of perfusion to demand is the primary mechanism through which symptom relief is achieved.

Question # 24

A 70-year-old man comes to the physician because of a 6-month-history of worsening shortness of breath on exertion and bouts of coughing while sleeping. He has hypertension, hyperlipidemia, and type 2 diabetes mellitus. Current medications include lisinopril, simvastatin, and insulin. The patient appears tired but in no acute distress. His pulse is 70/min, blood pressure is 140/85 mm Hg, and respirations are 25/min. He has crackles over both lower lung fields and 2+ pitting edema of the lower extremities. An ECG shows T wave inversions in leads V1 to V4. Which of the following agents is most likely to improve the patient's long-term survival?

	Answer	Image
A	Furosemide	
B	Gemfibrozil	
C	Metoprolol	
D	Digoxin	
E	Amlodipine	
F	Dobutamine	

Hint

This patient shows the classic clinical signs of congestive heart failure (CHF) as evidenced by his worsening exertional dyspnea and paroxysmal nocturnal dyspnea. The physical exam findings (crackles on auscultation suggesting pulmonary edema and peripheral edema), and T wave inversions on ECG (suggesting structural myocardial damage) corroborate this diagnosis.

Correct Answer

A - Furosemide

Explanation Why

[Furosemide](#) is a [loop diuretic](#) used for symptomatic treatment in patients with [CHF](#). While this patient with worsening [shortness of breath](#), fine, bilateral [crackles](#) at the [lung](#) bases, and lower extremity [edema](#) would benefit from [furosemide](#) to improve his breathing, oxygenation, and volume overload status, [furosemide](#) has not been associated with improved survival in patients with [CHF](#).


B - Gemfibrozil

Explanation Why

[Gemfibrozil](#) is a fibrate that can be used as a second-line agent in patients with [hyperlipidemia](#) and is most effective in treating [hypertriglyceridemia](#). [Fibrates](#) as a drug class are not known to improve survival in [heart failure](#). Furthermore, this patient already takes [simvastatin](#); the addition of [fibrates](#) would greatly increase his risk of [myopathy](#) and [rhabdomyolysis](#).

C - Metoprolol

Image



Explanation Why

[Beta blockers](#) such as [metoprolol](#) provide symptomatic improvement and reduce [mortality](#) in patients with [CHF](#). The beneficial mechanisms of action in [heart failure](#) include the lowering of the [heart rate](#) and subsequent anti-[ischemic](#) effects, as well as curbing the effect of chronic exposure to [catecholamines](#), which would otherwise lead to systemic [hypertension](#) and myocardial remodeling. The use of [metoprolol](#) would be of particular benefit in this patient, given his residual [hypertension](#) on [lisinopril](#). However, detailed clinical evaluation prior to initiation is required, especially since [beta blockers](#) should not be immediately given to patients with potentially [decompensated heart failure](#) (warning signs: [edema](#) and progressively worsening [dyspnea](#)) because their negative [inotropic](#) and [chronotropic](#) effects could acutely worsen symptoms.

D - Digoxin

Explanation Why

[Digoxin](#) is a [cardiac glycoside](#) that can be used for symptomatic treatment in patients with [atrial fibrillation](#) and [CHF](#) with preserved [ejection fraction \(HFpEF\)](#). This patient has no evidence of A-fib on [ECG](#) and would not be a candidate for [digoxin](#) treatment. Furthermore, while [digoxin](#) has been shown to improve the rate of hospitalizations in patients with [CHF](#), it has not been shown to improve overall survival.

E - Amlodipine

Explanation Why

[Amlodipine](#) is a [dihydropyridine calcium channel blocker \(CCB\)](#) that can be added to the medication regimen of hypertensive patients who do not fully respond to initial therapy such as this man ([hypertension](#) despite [lisinopril](#)). However, [CCBs](#) have not been shown to have a direct effect or survival benefit in patients with [CHF](#).

F - Dobutamine

Explanation Why

[Dobutamine](#) is an [inotropic](#) agent that can be used temporarily in patients with severe [left ventricular systolic dysfunction](#) and low [cardiac output](#) resulting in organ hypoperfusion. It has a role in severely ill patients with [cardiogenic shock](#) as a bridge to definitive therapy of the underlying cardiac condition (coronary revascularization therapy for [MI](#), [heart transplantation](#), among others). [Dobutamine](#) is not used chronically for management of [CHF](#).

Question # 25

A 23-year-old man comes to the physician with a 1-week history of sharp, substernal chest pain that is worse with inspiration and relieved with leaning forward. He has also had nausea and myalgias. His father has coronary artery disease. His temperature is 37.3°C (99.1°F), pulse is 110/min, and blood pressure is 130/84 mm Hg. Cardiac examination shows a high-pitched rubbing sound between S1 and S2 that is best heard at the left sternal border. An ECG shows depressed PR interval and diffuse ST elevations. Which of the following is the most likely cause of this patient's symptoms?

	Answer	Image
A	Dressler syndrome	
B	Acute myocardial infarction	
C	Mycobacterium tuberculosis infection	
D	Systemic lupus erythematosus	
E	Acute viral infection	

Hint

Acute pericarditis is an inflammation of the pericardium often characterized by sharp substernal chest pain, pericardial friction rub, and serial ECG changes.

Correct Answer

A - Dressler syndrome

Explanation Why

[Dressler syndrome](#) is an autoimmune complication that presents as [pericarditis](#) weeks after a [myocardial infarction](#). This patient without [coronary artery disease](#) or past [myocardial infarction](#) is unlikely to develop this type of [pericarditis](#).

B - Acute myocardial infarction

Explanation Why

Although substernal [chest pain](#) should raise the concern for acute [myocardial infarction](#) ([MI](#)), this patient does not have [risk factors](#) for [coronary artery disease](#) (e.g., [diabetes](#), tobacco use) aside from [family history](#). Furthermore, [acute myocardial infarction](#) does not present with positional [chest pain](#), friction rub, or noncontiguous [ST elevations](#). [MI](#) may cause [pericarditis](#) within 1–3 days (i.e., post-[infarction](#) fibrinous pericarditis), but this patient has no history of recent [MI](#).

C - Mycobacterium tuberculosis infection

Explanation Why

[Tuberculosis](#) is the most common cause of [pericarditis](#) in developing countries. However, [tuberculosis pericarditis](#) is unlikely in a patient who has no characteristic symptoms ([cough](#), weight loss, night sweats). Furthermore, he has no demographic [risk factors](#) (travel to [endemic](#) countries, work in hospitals/nursing homes, incarceration) nor medical [risk factors](#) (congenital or acquired [immunodeficiency](#)) of [tuberculosis](#).

D - Systemic lupus erythematosus

Explanation Why

Cardiovascular involvement is common in [SLE](#) patients, including [pericarditis](#) and [endocarditis](#) ([Libman-Sacks](#)). [Pericarditis](#) in [SLE](#) patients is most commonly subclinical, noted only on [echocardiogram](#). This patient presents with symptomatic [pericarditis](#), has no signs of [SLE](#), and has no [family history](#) of autoimmune disease.

E - Acute viral infection

Explanation But

Idiopathic cases are presumed to be viral because the efforts to confirm etiology are not routinely pursued due to the expenses involved, the inaccessibility of [pericardial](#) tissue/fluid, and the time delay and inaccuracy of viral studies (eg, culture, [PCR](#), viral serology, etc). In most cases, clinical management is not changed by these results.

Explanation Why

The most common cause of acute [pericarditis](#) is idiopathic (assumed viral etiology) followed by cases of confirmed viral etiology. The most common causes of viral [pericarditis](#) are [coxsackievirus B](#) and [echovirus](#); other pathogens include hepatitis viruses, [influenza](#) virus, [measles](#) virus, [mumps](#) virus, [varicella](#) virus, and [HIV](#).

Question # 26

A previously healthy 52-year-old woman comes to the physician because of a 3-month history of chest pain on exertion. She takes no medications. Cardiopulmonary examination shows no abnormalities. Cardiac stress ECG shows inducible ST-segment depressions in the precordial leads that coincide with the patient's report of chest pain and resolve upon cessation of exercise. Pharmacotherapy with verapamil is initiated. This drug is most likely to have which of the following sets of effects?

	End-diastolic volume (EDV)	Blood pressure (BP)	Contractility	Heart rate (HR)
A	↓	↓	↓	↑
B	No change	no change	no change	no change
C	↓	↓	↓	no change
D	↓	↓	no change	↑
E	↑	↓	↓	↓

	Answer	Image
A	A	
B	B	
C	C	
D	D	

	Answer	Image
E	E	The diagram illustrates the cellular mechanisms of cardiac inotropic regulation. It shows a cell membrane with various receptors and transporters. Key components include: β-receptor: Activated by β-agonist (positive inotropic agent) and inhibited by β-blocker (negative inotropic agent). Adenylate cyclase: Activated by the β-receptor, converting ATP to cAMP. PDE3 inhibitor: Inhibits PDE3, which normally converts cAMP to AMP. Protein kinase A: Activated by cAMP, leading to the activation of Myosin activators. L-type Ca channel: Allows Ca²⁺ entry into the cell, which is inhibited by Ca channel blockers (negative inotropic agents). Na⁺/K⁺-ATPase: A pump on the membrane that moves Na⁺ out and K⁺ in, using energy from ATP. Digoxin: Inhibits Na⁺/K⁺-ATPase, leading to increased intracellular Na⁺ and Ca²⁺ levels (positive inotropic effect). Smooth ER: Contains SERCA (Sarcoplasmic Endoplasmic Reticulum Ca²⁺ ATPase) and Ryanodine receptor. Ca sensitizer: Increases the sensitivity of the Ryanodine receptor to Ca²⁺, leading to increased Ca²⁺ release from the Smooth ER. Contraction: The final result of increased intracellular Ca²⁺ levels. Legend: - Positively inotropic agent (Blue box) - Negatively inotropic agent (Pink box)

Hint

Verapamil is a nondihydropyridine calcium channel blocker that inhibits calcium ion entry in vascular smooth muscle and the myocardium during depolarization. This causes relaxation of coronary vascular smooth muscle, coronary vasodilation, and slows AV nodal conduction.

Correct Answer

A - A

Explanation Why

These changes would be consistent with the effects of [nifedipine](#), a [dihydropyridine CCB](#) and potent [vasodilator](#) with minimal [myocardial](#) depressant effect. [Vasodilation](#) decreases [systemic vascular resistance](#) and decreases BP, resulting in decreased [venous return](#) and consequent decrease in [EDV](#). The decrease in BP can cause a [reflex tachycardia](#) (which decreases ejection time). [Nifedipine](#) also causes coronary [vasodilation](#), which improves [myocardial](#) oxygenation. [Nifedipine](#) only has a mild effect on cardiac contractility compared to [non-dihydropyridine CCBs](#).

B - B

Explanation Why

These changes would be consistent with [ranolazine](#), an antianginal drug that decreases myocardial oxygen demand without altering the [heart rate](#), contractility, end-[diastolic](#) volume, or blood pressure. [Myocardial ischemia](#) increases the late sodium current of the [cardiac action potential](#) and impairs [myocardial](#) relaxation. [Ranolazine](#) blocks these sodium channels, thereby improving [myocardial](#) relaxation and perfusion, and decreasing MVO₂ and [anginal](#) symptoms.

C - C

Explanation Why

This combination of effects would be consistent with the concurrent use of a [beta-blocker](#) and [nitrate](#), a common combination used in the treatment of [angina](#). [Nitrates](#) cause [vasodilation](#) (venous > arterial), which decreases BP, resulting in [reflex tachycardia](#). However, the [beta-blocker](#) neutralizes this [reflex tachycardia](#) and therefore also mitigates some of the decrease in [EDV](#) seen with [nitrate](#) use. [Beta-blockers](#) cause a mild decrease in cardiac contractility, and both medications cause coronary [vasodilation](#), which increases [myocardial](#) perfusion and decreases [anginal](#) symptoms.


D - D

Explanation Why

These changes would be consistent with the effects of [nitrates](#), one of the first-line drugs for the treatment of [angina](#). [Nitrates](#) cause [vasodilation](#) (venous > arterial), which decreases BP. [Vasodilation](#) leads to decreased venous return, which decreases the [preload](#), [EDV](#), and ventricular wall stress, thereby decreasing [myocardial](#) demand. The relative hypotension induced by [nitrates](#) leads to a [reflex tachycardia](#) (and decreases [ventricular ejection](#) time). [Nitrates](#) have no effect on the cardiac contractility.

E - E

Image



Explanation But


[Beta blockers](#) have the same effect as [verapamil](#) and are also used for the treatment of [stable angina](#).

Explanation Why

[Verapamil](#) is used for the treatment of [angina](#) because of its inhibitory effects on calcium channels located on the [myocardium](#) and vascular [smooth muscle cells](#). It decreases cardiac contractility (negative [inotropic](#) action) and [heart rate](#) (negative [chronotropic](#) action), slows [AV node](#) conduction (negative [dromotropic](#) action), and dilates the [coronary arteries](#). The [heart](#) beats slower and less forcefully, thereby decreasing myocardial oxygen demand. The relative [bradycardia](#) leads to increased ventricular [diastolic](#) filling time and increased [end-diastolic volume](#). In addition, inhibition of peripheral vascular [smooth muscle](#) calcium channels also causes arteriolar and [capillary vasodilation](#), resulting in decreased blood pressure.

Question # 27

Prior to undergoing a total knee arthroplasty, a 62-year-old man with coronary artery disease undergoes diagnostic cardiac catheterization. The catheter is inserted via the femoral artery and then advanced to the ascending aorta. Pressure tracing of one cardiac cycle is shown. The notch indicated by the arrow is most likely caused by which of the following?



	Answer	Image
A	Opening of the pulmonic valve	
B	Opening of the mitral valve	
C	Right atrial relaxation	

	Answer	Image
D	Closure of the aortic valve	Pressure (mm Hg) 120 80 40 0 Time Systole Con- traction Ejection Relaxation Diastole Filling Aorta Left ventricle Left atrium 1 2 3 4
E	Right ventricular contraction	
F	Left atrial contraction	

Hint

The event that leads to the smaller peak, so-called the "incisura" or "dicrotic notch," is itself caused by the equalization of pressure between the left ventricle and the ascending aorta.

Correct Answer

A - Opening of the pulmonic valve

Explanation Why

Opening of the pulmonic valve occurs at the same time as the opening of the [aortic valve](#); these both occur at the end of ventricular [isovolumetric contraction](#) and the beginning of rapid blood ejection into the ventricular outflow tract. Because the blood flowing through the pulmonic valve enters the [pulmonary arteries](#) and not the aorta, the pressure change will not be recorded on an aortic tracing.

B - Opening of the mitral valve

Explanation Why

The opening of the [mitral valve](#) precedes the rapid [filling phase](#) of the [left ventricle](#). This results in increased volume but not significantly increased pressure in the [left ventricle](#). It is not reflected in the aortic pressure waveform.


C - Right atrial relaxation

Explanation Why

Right atrial relaxation occurs following the atrial contraction and accounts for rapid [right ventricular](#) filling. This process does not affect pressures in the aorta.

D - Closure of the aortic valve

Image



Explanation Why

Closure of the [aortic valve](#) is associated with a small backflow of blood toward the [left ventricle](#). This causes a slight decrease in aortic pressure as the blood refluxes toward the closed valve, followed by a transient increase in pressure as this blood is reflected back into the lumen of the aorta. This process accounts for the notch ([incisura](#) or [dicrotic notch](#)) and secondary upstroke in the pressure tracings.

E - Right ventricular contraction

Explanation Why

[Right ventricular](#) contraction is not seen in the arterial waveform. Contraction of the [right ventricle](#) occurs in tandem with contraction of and ejection from the [left ventricle](#). The contraction of the [right](#)

[ventricle](#) is captured as the "c wave" on the [jugular venous pulse](#) curve rather than the [dicrotic notch](#).

F - Left atrial contraction

Explanation Why

Left atrial contraction occurs during rapid [left ventricular](#) filing but prior to [isovolumetric contraction](#). This leads to increased LV volume but does not significantly increase pressure there. It is therefore not reflected in the arterial waveform. Furthermore, contraction of the [left atrium](#) does not move blood into the aorta so it would not be expected to have an effect on pressure readings there.

Question # 28

A 72-year-old woman is admitted to the hospital for treatment of unstable angina. Cardiac catheterization shows occlusion that has caused a 50% reduction in the diameter of the left circumflex artery. Resistance to blood flow in this vessel has increased by what factor relative to a vessel with no occlusion?

	Answer	Image
A	64	
B	16	
C	8	
D	128	
E	4	
F	32	

Hint

The corresponding increase in resistance to blood flow can be calculated by the Poiseuille equation:

$R = 8L\eta/(\pi r^4)$, where R = resistance to flow, L = length of the vessel, η = viscosity, and r = radius of the vessel.

Correct Answer

A - 64

Explanation Why

Although the resistance to blood flow increases when the [radius](#) of the vessel decreases (by 50% in this case), this answer is incorrect.

B - 16

Explanation Why

When the length of the vessel and viscosity of blood remain constant, the relationship between resistance and the [radius](#) of the vessel in the [Poiseuille equation](#) can be simplified to $R \sim 1/r^4$. The diameter (and thus the [radius](#)) of this patient's vessel has decreased by 50%. Therefore, $R \sim 1/(0.5 \times r)^4 \rightarrow 1/(0.0625 \times r^4) \rightarrow 1/0.0625 \times 1/r^4 \rightarrow 16 \times 1/r^4$. Thus, resistance to blood flow increases 16 times.

C - 8

Explanation Why

Although the resistance to blood flow increases when the [radius](#) of the vessel decreases (by 50% in this case), this answer is incorrect.

D - 128

Explanation Why

Although the resistance to blood flow increases when the [radius](#) of the vessel decreases (by 50% in this case), this answer is incorrect.

E - 4

Explanation Why

Although the resistance to blood flow increases when the [radius](#) of the vessel decreases (by 50% in this case), this answer is incorrect.

F - 32

Explanation Why

Although the resistance to blood flow increases when the [radius](#) of the vessel decreases (by 50% in this case), this answer is incorrect.

Question # 29

A 22-year-old man comes to the physician because of a 2-month history of episodes of shortness of breath, lightheadedness, and palpitations. During the examination, he reports the onset of one such episode. His pulse is 170/min and regular, respirations are 22/min, and blood pressure is 100/65 mm Hg. An ECG shows a regular narrow complex tachycardia; no P waves are visible. A common clinical maneuver to diagnose and/or relieve the patient's symptoms involves stimulation of which of the following nerves?

	Answer	Image
A	Trigeminal	
B	Recurrent laryngeal	
C	Facial	
D	Phrenic	
E	Glossopharyngeal	

Hint

Intermittent palpitations, lightheadedness, dyspnea, and ECG findings that show a narrow complex tachycardia without P waves are all features of atrioventricular nodal re-entrant tachycardia, a type of paroxysmal supraventricular tachycardia (PSVT). Several maneuvers that elicit an autonomic response can be used to diagnose and terminate this cardiac arrhythmia.

Correct Answer

A - Trigeminal

Explanation Why

The [parasympathetic](#) fibers of the [trigeminal nerve \(CN V\)](#) supply the salivary glands, the [lacrimal glands](#), and the nasal mucosa. They do not supply the [myocardium](#) and will therefore not help to diagnose or relieve this patient's [PSVT](#). In addition, the [trigeminal nerve](#) does not lie in close proximity to the [carotid sinus](#).

B - Recurrent laryngeal

Explanation Why

Although the [recurrent laryngeal nerves](#) lie in close proximity to the [carotid sinus](#) and branch out from the [vagus nerve](#), they do not participate in the autonomic response that would terminate this patient's [PSVT](#).

C - Facial

Explanation Why

The [parasympathetic](#) fibers of the [facial nerve \(CN VII\)](#) supply the submandibular and sublingual salivary glands and the [lacrimal glands](#). They do not supply the [myocardium](#) and will therefore not help to diagnose or relieve this patient's [PSVT](#). In addition, the [facial nerve](#) does not lie in close proximity to the [carotid sinus](#).

D - Phrenic

Explanation Why

Although the [phrenic nerves](#) lie in close proximity to the [heart](#) and provide sensory supply to the [pericardium](#), they do not provide autonomic supply to the [myocardium](#). Therefore, they will not help to diagnose or relieve this patient's [PSVT](#).

E - Glossopharyngeal

Explanation But



Other [vagal maneuvers](#) include the [Valsalva maneuver](#) and breath holding.




Explanation Why

The nerve of the [carotid sinus](#) ([nerve of Hering](#)) is a branch of the [glossopharyngeal nerve](#) (CN IX) that terminates in the vasomotor center of the [medulla oblongata](#) ([nucleus tractus solitarius](#)). [Carotid sinus massage](#) stimulates [baroreceptors](#), which transmits a signal to the [medulla oblongata](#) via [CN IX](#). The efferent signals from the medulla to the [myocardium](#) are conducted via the [vagus nerve](#) ([parasympathetic](#) supply), which results in ↓ [AV nodal](#) conduction, ↓ [heart](#) rate, ↓ contractility, and [vasodilation](#) (↓ blood pressure). This patient's symptoms are caused by [AV nodal](#) re-entry; slowing [AV nodal](#) conduction terminates the [arrhythmia](#).

Question # 30

A 23-year-old man comes to the emergency department because of palpitations, dizziness, and substernal chest pain for three hours. The day prior, he was at a friend's wedding, where he consumed seven glasses of wine. The patient appears diaphoretic. His pulse is 220/min and blood pressure is 120/84 mm Hg. Based on the patient's findings on electrocardiography, the physician diagnoses atrial fibrillation with rapid ventricular response and administers verapamil for rate control. Ten minutes later, the patient is unresponsive and loses consciousness. Despite resuscitative efforts, the patient dies. Histopathologic examination of the heart at autopsy shows an accessory atrioventricular conduction pathway. Electrocardiography prior to the onset of this patient's symptoms would most likely have shown which of the following findings?

	Answer	Image
A	Slurred upstroke of the QRS complex	 ECG tracing showing a normal QRS complex. The tracing includes leads I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. The QRS complex is narrow and has a sharp upstroke. The P wave is visible before the QRS complex. The T wave is visible after the QRS complex. The tracing is labeled with 'HR: 68', 'PR: 88 ms', 'PQ: 112 ms', 'QRS: 125 ms', 'QT: 400 ms', 'QTc: 418 ms', 'ST: 100 ms', 'Td: 174 ms', 'P ax: 64', 'QRS ax: 87', 'T ax: 71', and 'QRS-T ax(3D): 116'. The bottom of the tracing shows 'Cardiac V4 37.7 95-02-2021 13:38 1' and 'T: 0.00 - 0.05'.
B	Cyclic alteration of the QRS axis	 ECG tracing showing a wide QRS complex with a cyclic alteration of the QRS axis. The tracing includes leads I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. The QRS complex is wide and has a cyclic alteration of the QRS axis. The P wave is visible before the QRS complex. The T wave is visible after the QRS complex. The tracing is labeled with 'aVR', 'V1', 'V4', 'aVL', 'V2', 'V5', 'aVF', 'V3', and 'V6'.

	Answer	Image
C	Pseudo-right bundle branch block with ST elevation in V ₁₋₂	 The ECG tracing displays 12 leads: I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. There is a characteristic M-shaped ST segment with significant ST-segment depression in leads I, II, III, aVL, and aVF, and ST-segment depression in leads V4, V5, and V6. Conversely, there is ST-segment elevation in leads V1, V2, and V3. The QRS complex in leads I, II, III, aVL, and aVF shows a deep S wave, while in leads V1, V2, and V3, there is a deep R wave. This pattern is consistent with pseudo-right bundle branch block. 25mm/s 10mm/mV 40Hz 0050 128L 233 CID: 15 Courtesy of P.G. Postema, M.D., AMC ECG-JPEDIA.ORG
D	Epsilon wave following the QRS complex	
E	Prolongation of the QT interval	 The ECG tracing shows 12 leads: I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. The QT interval is significantly prolonged across all leads, which is a characteristic finding in long QT syndrome. 40 Hz 25.0 mm/s 10.0 mm/mV 2 kg 5s = 1 minute off MACRO 995A 11/18/01 14:02 ECG-JPEDIA.ORG
F	Positive Sokolow-Lyon index	 The ECG tracing displays 12 leads: I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. The Sokolow-Lyon index is positive, indicating a deep S wave in lead I (S_I) and a deep R wave in lead V5 (R_{V5}), which is a characteristic finding in left bundle branch block.


Hint

The autopsy finding of an accessory atrioventricular conduction pathway indicates Wolff-Parkinson-White (WPW) syndrome.

Correct Answer

A - Slurred upstroke of the QRS complex

Image




Explanation Why

A slurred upstroke of the [QRS complex](#) (i.e., delta wave) and shortened [PR interval](#) are the characteristic [ECG](#) findings of [WPW syndrome](#). These findings are caused by an accessory atrioventricular pathway (i.e., [bundle of Kent](#)) that bypasses the [AV node](#). [Atrial fibrillation](#) occurs in about 20% of patients with [WPW syndrome](#) and should be treated with rhythm control agents (e.g., [procainamide](#)). [AV nodal](#) blocking agents (e.g., adenosine, [beta-blockers](#), [nondihydropyridine calcium channel blockers](#) such as [verapamil](#)) are contraindicated since they increase conduction of atrial impulses through the [accessory pathway](#). This pathway has a short [refractory period](#), allowing frequent conduction to the ventricles and induction of fatal [arrhythmias](#) (e.g., [ventricular fibrillation](#)), which has occurred here.

B - Cyclic alteration of the QRS axis

Image




Explanation Why

Cyclic alteration of the QRS axis on [ECG](#) is characteristic of [torsades de pointes](#). Symptoms can include [palpitations](#), [syncope](#), [ventricular arrhythmias](#), and [sudden cardiac death](#). While [torsades de pointes](#) has many etiologies (e.g., [long QT syndrome](#), electrolyte abnormalities, and drugs), it is not caused by an accessory atrioventricular pathway.

C - Pseudo-right bundle branch block with ST elevation in V₁₋₂

Image



Explanation Why

Pseudo-right bundle branch block and ST elevations in leads V₁ and V₂ are characteristic ECG findings of [Brugada syndrome](#), which may cause [dizziness](#), [palpitations](#), and [atrial fibrillation](#). Additionally, [ventricular arrhythmias](#) may lead to [syncope](#) and [sudden cardiac death](#) in these patients. However, [Brugada syndrome](#) is caused by defective cardiac voltage-gated sodium channels rather than an accessory atrioventricular conduction pathway.

D - Epsilon wave following the QRS complex


Explanation Why

An epsilon wave following the QRS complex is highly specific for [arrhythmogenic right ventricular cardiomyopathy \(ARVC\)](#), which may cause [dizziness](#), [palpitations](#), [chest pain](#) and [sudden cardiac](#)

[death](#). [ARVC](#) is caused by mutations in [desmosomal proteins](#), which may lead to fatal [arrhythmias](#). However, an accessory atrioventricular pathway is inconsistent with [ARVC](#).

E - Prolongation of the QT interval

Image




Explanation Why

[Prolongation of the QT interval](#) is seen on [ECG](#) in patients with [long QT syndrome \(LQTS\)](#), which can manifest with [palpitations](#), [dizziness](#), and [sudden cardiac death](#). [LQTS](#) can cause [arrhythmias](#) (e.g., [torsade de pointes](#)) but does not typically cause [atrial fibrillation](#). Furthermore, [congenital LQTS](#) is a result of defective potassium channels in the [myocytes](#) rather than an accessory atrioventricular conduction pathway.

F - Positive Sokolow-Lyon index

Image



Explanation Why

A positive [Sokolow-Lyon](#) index on [ECG](#) indicates ventricular [hypertrophy](#). [Hypertrophic cardiomyopathy \(HCM\)](#) can manifest with [chest pain](#), [palpitations](#), [dizziness](#), and [sudden cardiac death](#) (due to [arrhythmias](#)) in a young patient. However, a rate control agent (e.g., [verapamil](#)) is a component of treatment for [HCM](#) and would not have the deleterious effect seen in this patient. Furthermore, [HCM](#) is caused by mutations in [sarcomere proteins](#) rather than an accessory atrioventricular pathway.

Question # 31

An 8-year-old girl is brought to the physician by her parents for the evaluation of an episode of unconsciousness while at the playground that morning. She was unconscious for about 15 seconds and did not shake, bite her tongue, or lose bowel or bladder control. Her grandfather died suddenly at the age of 29 of an unknown heart condition; her parents are both healthy. An ECG shows sinus rhythm and a QT interval corrected for heart rate (QTc) of 470 milliseconds. Laboratory studies are within normal limits. Which of the following is the most likely additional finding in this patient?

	Answer	Image
A	Oblique palpebral fissures	
B	Sensorineural hearing loss	
C	Skin folds between the mastoid process and acromion	
D	Brachial-femoral pulse delay	
E	Subvalvular ventricular outflow obstruction murmur	

Hint

This patient with a family history of sudden death presents with syncope associated with exercise and a prolonged QTc (> 440 milliseconds) on ECG, both of which indicate the diagnosis of long QT syndrome (LQTS). There is no indication of electrolyte or endocrine abnormalities that could explain this patient's prolonged QT interval, making a congenital rather than acquired LQTS the most likely diagnosis.

Correct Answer

A - Oblique palpebral fissures

Explanation Why

Oblique [palpebral fissures](#) is a characteristic finding of [Down syndrome](#). While [Down syndrome](#) is associated with cardiac anomalies including [endocardial cushion defects](#) and [Tetralogy of Fallot](#), it is not associated with [long QT syndrome](#).

B - Sensorineural hearing loss

Explanation Why

Congenital [long QT syndrome](#) can be divided into [Romano-Ward syndrome](#) and Jervell and Lange-Nielson syndrome; the latter is associated with [sensorineural deafness](#). Jervell and Lange-Nielson syndrome is [autosomal recessive](#) and more likely in this patient with an affected grandparent. This syndrome is caused by a mutation in the [gene](#) encoding outward-rectifying potassium channels, which are important in membrane [repolarization](#) in cardiac [myocytes](#) as well as [neurons](#) in the [inner ear](#). Romano-Ward syndrome is [autosomal dominant](#), and while it is more common, it is unlikely in a patient with unaffected parents.

C - Skin folds between the mastoid process and acromion

Explanation Why

[Webbed neck](#) is a characteristic finding of [Turner syndrome](#), which may be associated with cardiac anomalies including [coarctation of the aorta](#) and [bicuspid aortic valve](#). It is not associated with [QT interval prolongation](#).

D - Brachial-femoral pulse delay

Explanation Why

Brachial-femoral [pulse](#) delay, or a delay in palpable pulses between the brachial and [femoral arteries](#), is present in [coarctation of the aorta](#). This condition is often asymptomatic in children and does not typically cause [syncope](#). It is not associated with [QT interval prolongation](#).

E - Subvalvular ventricular outflow obstruction murmur

Explanation Why

Subvalvular ventricular outflow obstruction is present in [hypertrophic obstructive cardiomyopathy \(HOCM\)](#). This causes a [systolic murmur](#) that increases in intensity with maneuvers that decrease [venous return](#) such as standing up from squatting or the active phase of valsalva. While [HOCM](#) is associated with [arrhythmias](#), [syncope](#), and [sudden cardiac death](#), it is not associated with a prolonged [QT interval prolongation](#).

Question # 32

A 69-year-old man with hypertension and congestive heart failure is brought to the emergency department because of a 9-day history of worsening shortness of breath and swelling of his legs. His respirations are 25/min, and blood pressure is 160/98 mm Hg. Pulse oximetry on 5 L O₂ via nasal cannula shows an oxygen saturation of 92%. Examination shows 2+ pretibial edema bilaterally. Crackles are heard at both lung bases. The patient's symptoms are due in part to an increase in the rate of bradykinin breakdown. The substance responsible for bradykinin breakdown is primarily produced in which of the following?

	Answer	Image
A	Pulmonary endothelium	The diagram illustrates the RAAS pathway. Renin converts Angiotensinogen to Angiotensin I. ACE converts Angiotensin I to Angiotensin II. Angiotensin II causes arteriolar vasoconstriction, efferent arteriole constriction, and stimulates the release of Aldosterone and ADH. Aldosterone acts on the distal tubule and collecting duct to increase Na⁺ and H₂O reabsorption. ADH acts on the collecting duct to increase H₂O reabsorption. Both lead to increased blood pressure and cardiac output. The diagram also shows the breakdown of Bradykinin to inactive metabolites by ACE, and the inhibition of ACE by ACE inhibitors (ACEIs).
B	Liver	
C	Atria	
D	Zona glomerulosa	
E	Juxtaglomerular cells	


Hint

Bradykinin is degraded by angiotensin-converting enzyme (ACE), a component of the renin-angiotensin-aldosterone system (RAAS).

Correct Answer

A - Pulmonary endothelium

Image



Explanation Why

ACE is produced by cells of the pulmonary **endothelium**. Circulating **bradykinin** is therefore primarily metabolized in the pulmonary vasculature.

B - Liver

Explanation Why

ACE is not made in the **liver**. Angiotensinogen, however, is produced in the **liver** before being converted to **angiotensin I** by **renin** in the **kidneys**.

C - Atria

Explanation Why

[ACE](#) is not made by the cardiac [atria](#). [Atrial natriuretic peptide \(ANP\)](#), however, is produced by atrial [myocytes](#), which then acts on the [kidneys](#).

D - Zona glomerulosa

Explanation Why

[ACE](#) is not made in the [zona glomerulosa](#). [Aldosterone](#), however, is produced in the [zona glomerulosa](#) in response to [angiotensin II](#), a substance converted from [angiotensin I](#) by [ACE](#).

E - Juxtaglomerular cells

Explanation Why

[ACE](#) is not made by [juxtaglomerular cells](#). [Renin](#), however, is produced by [juxtaglomerular cells](#) and converts angiotensinogen to [angiotensin I](#).

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

Question # 33

A 72-year-old man with coronary artery disease comes to the physician because of intermittent episodes of substernal chest pain and shortness of breath. The episodes occur only when walking up stairs and resolves after resting for a few minutes. He is a delivery man and is concerned because the chest pain has impacted his ability to work. His pulse is 98/min and blood pressure is 132/77 mm Hg. Physical examination is unremarkable. An ECG shows no abnormalities. A drug that blocks which of the following receptors is most likely to prevent future episodes of chest pain from occurring?

	Answer	Image
A	Alpha-2 adrenergic receptors	
B	Angiotensin II receptors	
C	Aldosterone receptors	
D	M ₂ muscarinic receptors	
E	Beta-1 adrenergic receptors	

Hint

This patient has stable angina and will benefit from a drug that reduces myocardial oxygen demand.

Correct Answer

A - Alpha-2 adrenergic receptors

Explanation Why

Alpha-2 [adrenergic receptor](#) blockade would worsen this patient's [chest pain](#) by increasing the release of [catecholamines](#) in the presynaptic [neurons](#) of the [sympathetic nervous system](#) (PSNS). This, in turn, increases blood pressure ([afterload](#)) and workload of the [heart](#), which increases myocardial oxygen demand and consumption. Alpha-2 blockers have limited clinical application and are not used for [coronary artery disease](#). Instead, alpha-2 adrenergic agonists, such as [clonidine](#), are used to treat [hypertension](#).

B - Angiotensin II receptors

Explanation Why

[Angiotensin II](#) receptor blockade (e.g., [ARBs](#)) causes [vasodilation](#), which decreases blood pressure, [afterload](#), and cardiac oxygen demand and consumption. Although these drugs are commonly used in patients post-[MI](#) and are proven to decrease [mortality rates](#), [ARBs](#) are not used for symptomatic treatment of [stable angina](#).

C - Aldosterone receptors

Explanation Why

[Aldosterone](#) receptor blockade (e.g., [spironolactone](#), [eplerenone](#)) decreases sodium reabsorption and inhibits hydrogen and potassium excretion in the [collecting ducts](#) of the [kidneys](#). These agents are commonly used in [CHF](#) and are proven to decrease [mortality rates](#). However, [aldosterone antagonists](#) are not used for symptomatic treatment of stable [angina](#).

D - M₂ muscarinic receptors

Explanation Why

M₂ [muscarinic receptor](#) blockade would worsen this patient's [chest pain](#) by decreasing [parasympathetic](#) activity in the [AV node](#), which increases [heart rate](#), cardiac workload, and myocardial oxygen demand and consumption. [Muscarinic antagonists](#) (e.g., [atropine](#)) are first-line agents for the treatment of symptomatic [bradycardia](#) (e.g., [third-degree AV block](#)).

E - Beta-1 adrenergic receptors

Explanation Why

[Beta-1 adrenergic receptor](#) blockade decreases [heart rate](#) and cardiac contractility (negative [chronotropic](#) and [inotropic](#) effect, respectively). This, in turn, decreases blood pressure and myocardial oxygen demand, and therefore [anginal](#) symptoms. [Beta-1 receptor](#) blockade also decreases [AV node](#) conduction (increased [refractory period](#)), giving drugs targeting this receptor an antiarrhythmic effect.

Question # 34

A 52-year-old man with hypertension and hyperlipidemia comes to the emergency department 30 minutes after the sudden onset of substernal chest pain while sitting in a chair at home. He reports drinking 2 glasses of whiskey earlier that day, after a stressful day at work. Current medications include hydrochlorothiazide and atorvastatin. He has smoked one pack of cigarettes daily for the past 20 years. His pulse is 102/min, and blood pressure is 135/88 mm Hg. Cardiopulmonary examination shows normal heart sounds. An ECG obtained on arrival at the emergency department shows ST-segment elevations in the anterior leads. 15 minutes later, the patient's chest pain has resolved and a repeat ECG shows no abnormalities. Which of the following is the most likely underlying mechanism of this patient's chest pain?

	Answer	Image
A	Coronary artery vasospasm	
B	Coronary artery vasodilation	
C	Atherosclerotic coronary artery narrowing	The diagram illustrates two types of myocardial infarction. On the left, a 'Transmural infarct' is shown with a cross-section of the heart wall where the entire thickness is affected by a 'Complete occlusion of a major coronary artery'. Below this, a box titled 'Common ECG changes' lists 'ST elevation, pathological Q wave' and shows a corresponding ECG trace. On the right, a 'Subendocardial infarct' is shown with a cross-section where only the inner layer of the heart wall is affected due to a 'Transient or partial occlusion of a major coronary artery'. Below this, a box titled 'Common ECG changes' lists 'ST depression, T-wave inversion' and shows a corresponding ECG trace.
D	Pulmonary artery occlusion	
E	Atherosclerotic plaque rupture	

	Answer	Image
F	Pericardial inflammation	

Hint

Substernal chest pain with transient ST-segment elevations is consistent with Prinzmetal angina.

Correct Answer

A - Coronary artery vasospasm

Explanation Why

[Coronary artery](#) vasospasm is the underlying mechanism of [Prinzmetal angina](#). Episodes of [Prinzmetal angina](#) typically occur at rest (in contrast to exertional [angina](#)) and common triggers include stress, smoking, alcohol, drugs (e.g., [cocaine](#)), and [triptan](#) use. The transient appearance of [anterior](#) ST-segment elevations (as opposed to depressions) and spontaneous resolution of symptoms in the setting of recent alcohol use and stress support this diagnosis.


B - Coronary artery vasodilation

Explanation Why

[Coronary artery vasodilation distal](#) to an atherosclerotic stenosis is the underlying cause of [coronary steal syndrome](#). Although this condition also presents with symptoms of [angina](#), symptoms are usually triggered by [vasodilators](#) (e.g., [dipyridamole](#), adenosine, or [hydralazine](#)), which this patient does not take. Moreover, [ECG](#) changes, if present, almost always consist of [ST-segment depressions](#).

C - Atherosclerotic coronary artery narrowing

Image



Explanation Why

Atherosclerotic narrowing of a [coronary artery](#) of $> 70\%$ is the underlying mechanism of [stable angina](#), which is typically characterized by exertional [chest pain](#) that improves with rest. Physical exertion leads to increased [myocardial](#) demand and subsequent transient hypoperfusion of the [myocardium](#) (mismatch between oxygen supply and demand). Although this patient has [risk factors](#) for atherosclerotic narrowing (e.g., [hypertension](#), [hyperlipidemia](#), smoking), his symptoms are atypical ([chest pain](#) at rest) and [ST-segment depressions](#) would be expected on [ECG](#).

D - Pulmonary artery occlusion

Explanation Why

Patients with [pulmonary artery](#) occlusion due to [pulmonary embolism](#) typically present with [pleuritic](#)

[chest pain](#), rather than the substernal [pain](#) reported here. The absence of [dyspnea](#) and prominent [risk factors for PE](#) (e.g., recent surgery, immobility, [hypercoagulable state](#)) in this patient also make [pulmonary artery](#) occlusion an unlikely cause of his symptoms. Moreover, [ECG](#) findings are usually normal or suggestive of right [heart](#) strain (e.g., [right bundle branch block](#), [right axis deviation](#)).

E - Atherosclerotic plaque rupture

Explanation Why

[Atherosclerotic plaque](#) rupture with subsequent thrombosis and acute occlusion of a [coronary artery](#) is the underlying mechanism of [chest pain](#) due to [unstable angina](#) and [myocardial infarction \(MI\)](#). [Unstable angina](#) may occur at rest but classically manifests with transient [ST-segment depressions](#) on [ECG](#), rather than the [ST elevations](#) seen in this patient. [MI](#) typically results in [ST elevations](#) on [ECG](#), but would not resolve spontaneously within 15 minutes. This patient's [ECG](#) findings and resolution of symptoms are inconsistent with either diagnosis and an [atherosclerotic plaque](#) rupture is unlikely to be a cause of his symptoms.


F - Pericardial inflammation

Explanation Why

[Pericardial inflammation](#), which is characteristic of [pericarditis](#), most commonly presents with sharp, positional [chest pain](#) (e.g., worse when laying back and relieved when hunched forward), as opposed to the substernal [pain](#) reported here. Moreover, [ECG](#) typically shows diffuse [ST-segment elevations](#) throughout the [limb leads](#) and [precordial leads](#) (V2–V6) with PR segment depressions.

Question # 35

One day after a 4700-g (10-lb 6-oz) male newborn is delivered to a 28-year-old primigravid woman, he has bluish discoloration of his lips and fingernails. Oxygen saturation on room air is 81%. Examination shows central cyanosis. A continuous machine-like murmur is heard over the left upper sternal border. A single S2 heart sound is present. Supplemental oxygen does not improve cyanosis. Echocardiography shows the pulmonary artery arising from the posterior left ventricle and aorta arising from the right ventricle with active blood flow between the right and left ventricles. Further evaluation of the mother is most likely to show which of the following?

	Answer	Image
A	Prenatal lithium intake	
B	Elevated serum TSH	
C	Prenatal alcohol use	
D	Prenatal phenytoin intake	
E	Positive rapid plasma reagin test	
F	Elevated fasting blood glucose	

Hint

Echocardiography showing a pulmonary artery arising from the left ventricle and aorta arising from the right ventricle confirms the diagnosis of transposition of the great arteries, which is an important cause of neonatal cyanosis that will not improve with supplemental oxygen. The continuous machine-like murmur indicates a coexistent patent ductus arteriosus, which is critical for life, as it allows mixing of oxygenated blood and its entry into systemic circulation. Patients may also have coexistent arterial or ventricular septal defects that allow mixing of blood between the two circuits.

Correct Answer

A - Prenatal lithium intake

Explanation Why

Prenatal [lithium](#) intake is associated with [Ebstein anomaly](#), which can have a wide range of clinical presentations, including congenital [cyanosis](#) from [right-to-left shunting](#). Depending on the severity of the anomaly, the patient may have a [holosystolic murmur](#) (due to [tricuspid regurgitation](#)), as heard in this patient. However, [echocardiography](#) would show right-sided [cardiomegaly](#) with apical displacement of the [tricuspid valve](#) and [tricuspid regurgitation](#), not the [TGA](#) seen in this patient.

B - Elevated serum TSH

Explanation Why

Elevated serum [TSH](#) levels suggest [hypothyroidism](#). Subclinical maternal [hypothyroidism](#) may increase the risk for future cognitive impairment, but it is not associated with [TGA](#).

C - Prenatal alcohol use

Explanation Why

Prenatal alcohol consumption can cause [fetal alcohol spectrum disorder](#). [FASD](#) can present with a wide range of cardiac abnormalities, including [atrial septal defect](#), [VSD](#), and great [heart](#) vessel defects. However, structural [heart](#) defects would also usually present with other more characteristic [FASD](#) findings, such as facial abnormalities (short [palpebral fissures](#), thin [vermillion border](#), smooth [philtrum](#)) and growth retardation.

D - Prenatal phenytoin intake

Explanation Why

Prenatal [phenytoin](#) intake is associated with [fetal hydantoin syndrome](#). Although cases of congenital [heart](#) defects have been reported with maternal [phenytoin](#) use, [transposition of the great arteries](#) is rare. Furthermore, [fetal hydantoin syndrome](#) also presents with [nail hypoplasia](#), [hypoplasia](#) of the digits, and abnormal facial features, which are all absent in this patient.


E - Positive rapid plasma reagin test

Explanation Why

A positive rapid plasma reagin test suggests maternal [syphilis](#), which can infect the [infant](#) in utero. Most cases of [early congenital syphilis](#) are asymptomatic at [birth](#) and do not develop symptoms until 5 weeks of life. If symptomatic at [birth](#), findings include [hepatosplenomegaly](#), [jaundice](#), nasal discharge, [generalized lymphadenopathy](#), skeletal abnormalities, and/or and pemphigus syphiliticus, none of which are present in this [infant](#). [Transposition of the great arteries](#) is not associated with [congenital syphilis](#).

F - Elevated fasting blood glucose

Image



Explanation Why

[Transposition of the great arteries](#) is a [cyanotic heart](#) condition that is associated with maternal [diabetes](#), caused by abnormal rotation of the [pulmonary artery](#) and aorta during [fetal development](#). Mothers with untreated [diabetes](#) will have elevated [fasting blood glucose](#) levels. Other potential neonatal complications from maternal [diabetes](#) include fetal [macrosomia](#) (seen in this patient), [polycythemia](#), [hypoglycemia](#), [hypocalcemia](#), and [neonatal respiratory distress syndrome](#).

Question # 36

A group of investigators is studying hemodynamic regulatory mechanisms in a human volunteer subject. The volunteer is administered a stimulant drug, and a sudden increase in blood pressure is observed. It is hypothesized that the increase in blood pressure causes stretching of the atria. Increased atrial stretch would most likely lead to which of the following changes?

	Answer	Image
A	Increased excretion of potassium and hydrogen ions	
B	Vasodilation and decreased platelet aggregation	
C	Decreased reabsorption of sodium	
D	Increased reabsorption of solute-free water	
E	Vasoconstriction and stimulation of thirst	The flowchart illustrates the RAAS pathway. It starts with 'Hypotension, Hypovolemia (activates juxtaglomerular cells)' and 'Hypovolemia (activates osmotic sensors)' leading to 'Renin'. Renin converts 'Angiotensinogen' to 'Angiotensin I'. 'Angiotensin I' is converted to 'Angiotensin II' by 'ACE'. 'Angiotensin II' has several effects: it stimulates 'Aldosterone release' (which leads to 'distal tubule & collecting duct' and '↑ K⁺ urinary secretion'), it acts on 'Renal prox. tubule' (leading to '↑ Na⁺ reabsorption + H₂O retention'), it acts on 'Central' (leading to '↑ ADH release', '↑ Na⁺ appetite', and '↑ Thirst'), and it causes 'Arteriole constriction'. All these effects ultimately lead to '↑ Extracellular fluid volume + ↑ blood pressure'. There are also feedback loops: 'Angiotensin II' inhibits 'Renin' release and 'Aldosterone release'.

Hint

Atrial cardiomyocytes are stimulated by increased atrial stretch to produce atrial natriuretic peptide (ANP).

Correct Answer

A - Increased excretion of potassium and hydrogen ions

Explanation Why

Increased excretion of potassium and hydrogen ions in the [distal](#) tubules and [collecting ducts](#) of the [kidney](#), as well as reabsorption of sodium, are effects of [aldosterone](#), not of [ANP](#). [Aldosterone](#) is stimulated by [RAAS](#), increased serum potassium, [ACTH](#), the [sympathetic nervous system](#), and decreased [mean arterial pressure](#).

B - Vasodilation and decreased platelet aggregation

Explanation Why

[Nitric oxide](#) (NO) results in [vasodilation](#) and decreased [platelet aggregation](#). NO is produced by vascular [endothelial](#) cells in response to a variety of triggers, including vascular injury and [hypoxia](#). While [ANP](#) does cause [vasodilation](#), it has no effect on [platelet](#) aggregation.

C - Decreased reabsorption of sodium

Explanation Why

Decreased reabsorption of sodium from the [distal convoluted tubule](#) and cortical [collecting duct](#) is one of the effects of [ANP](#), which responds to increased atrial stretch (e.g., in [hypertension](#) or volume overload) by increasing [natriuresis](#) and diuresis in the body. In addition to decreasing sodium reabsorption, [ANP](#) increases sodium and water excretion by increasing the [GFR](#) via dilation of the [afferent arteriole](#) and constriction of the [efferent arteriole](#). Lastly, [ANP](#) inhibits [renin](#) secretion and thereby [RAAS](#). Together, these changes work to reduce blood volume, blood pressure, and [cardiac output](#).


D - Increased reabsorption of solute-free water

Explanation Why

Increased reabsorption of solute-free water is an effect of vasopressin ([ADH](#)), not of [ANP](#). [ADH](#) acts by increasing [transcription](#) and insertion of [aquaporins](#) into the renal [collecting ducts](#). [ADH](#) is secreted if the body detects increased plasma osmolality or decreased arterial blood volume.

E - Vasoconstriction and stimulation of thirst

Image



Explanation Why

Systemic vasoconstriction and stimulation of thirst, as well as sodium retention and increased water reabsorption, are stimulated by [RAAS](#), not of [ANP](#). [RAAS](#) is activated in response to decreased [glomerular](#) perfusion and by the renal [sympathetic nervous system](#) and is actually inhibited by [ANP](#).

Question # 37

A 29-year-old man is brought to the emergency department 20 minutes after being stabbed in the left thigh. His pulse is 110/min, respirations are 20/min, and blood pressure is 110/70 mm Hg. Examination shows a 2-cm wound overlying a pulsatile mass on the left anterior thigh, 4 cm below the inguinal crease. A thrill is palpated, and a bruit is heard over this area. Peripheral pulses are normal bilaterally. The patient is at greatest risk for which of the following?

	Answer	Image
A	Pudendal nerve compression	
B	High-output cardiac failure	
C	Pulmonary embolism	
D	Iliac artery aneurysm	
E	Femoral head necrosis	
F	Erectile dysfunction	

Hint

This patient has a traumatic arteriovenous fistula, which typically manifests as a pulsatile mass with a palpable thrill and audible bruit.

Correct Answer

A - Pudendal nerve compression

Explanation Why

The [aneurysmal](#) dilation of an [arteriovenous fistula](#) in the [anterior](#) thigh 4 cm below the inguinal crease can result in compression of surrounding structures. However, [pudendal nerve](#) compression is unlikely given the location of this patient's injury.

B - High-output cardiac failure

Explanation Why

[High-output cardiac failure](#) can occur following the formation of a traumatic [arteriovenous fistula](#) due to the shunting of blood from the arterial high-pressure system to the venous low-pressure system. As a result of this shunt, [systemic vascular resistance](#) decreases, which, in turn, results in an increased [heart rate](#), [stroke volume](#), and [cardiac output](#). However, despite increased [cardiac output](#), low [systemic vascular resistance](#) results in arterial underfilling, decreased renal perfusion, and retention of salt and water, which leads to volume overload and [heart failure](#).

C - Pulmonary embolism

Explanation Why

Although this patient has an increased risk of thrombus development at the site of an [arteriovenous fistula](#), a [pulmonary embolism](#) would be an unlikely complication because a dislodged [thrombus](#) would travel distally into the affected [vein](#) as a result of the reverse blood flow caused by the pressure differential from the arterial to venous systems.

D - Iliac artery aneurysm

Explanation Why

[Risk factors](#) for the development of an [iliac artery aneurysm](#) include [connective tissue disorders](#), older age, history of smoking, [hypertension](#), and [atherosclerosis](#). The mechanism of this patient's injury could compromise the vascular integrity at the level of the injury but is unlikely to affect the iliac artery given how [distal](#) the wound is.

E - Femoral head necrosis

Explanation Why

[Osteonecrosis of the femoral head](#) can result from trauma to the [medial circumflex femoral artery](#) (e.g., from [femoral head fracture](#) or [lateral](#) or [medial femoral neck fracture](#), especially with dislocation). However, a stab wound in the [anterior](#) thigh is unlikely to affect the [medial circumflex femoral artery](#).

F - Erectile dysfunction

Explanation Why

[Erectile dysfunction](#) could result from an untreated [arteriovenous fistula](#) of the [internal pudendal artery](#), which gives rise to the [deep artery of the penis](#). However, the site of this patient's injury is not near the [internal pudendal artery](#).

Question # 38

A 63-year-old woman comes to the physician because of a 6-month history of fatigue and shortness of breath on exertion in addition to a 1-month history of ankle swelling. She has a history of sleep apnea. Examination of the lower extremities shows bilateral pitting pedal and ankle edema. Cardiac examination shows a widely split S₂ that varies with respiration. An ECG shows right bundle branch block. Which of the following is the most likely etiology of the patient's lower extremity edema?

	Answer	Image
A	Localized lymphatic obstruction	
B	Dermal accumulation of glycosaminoglycans	
C	Thrombus formation in a deep vein	
D	Increased capillary hydrostatic pressure	
E	Decreased plasma oncotic pressure	

Hint

The patient's symptoms of exertional dyspnea and lower extremity edema, and physical exam findings of a widely split S2 and right bundle branch block are consistent with right heart failure caused by pulmonary hypertension (cor pulmonale). Sleep apnea can lead to pulmonary hypertension.

Correct Answer

A - Localized lymphatic obstruction

Explanation Why

Patients who have undergone [lymph node](#) removal, [radiation therapy](#), or have a history of cancer are at risk for [edema](#) caused by lymphatic obstruction (i.e., [lymphedema](#)). This type of [edema](#) is generally not pitting and would not explain the [shortness of breath](#) seen in this patient. Her [edema](#), [shortness of breath](#), and [physical examination](#) findings are instead related to [heart failure](#).

B - Dermal accumulation of glycosaminoglycans

Explanation Why

[Myxedema](#), a specific type of [edema](#) that manifests in a small portion of patients with [Graves disease](#) or severe [hypothyroidism](#), is caused by the [dermal](#) accumulation of [glycosaminoglycans](#). Although this patient's [edema](#) is in the lower extremities, which is similar to [myxedema](#), she has no signs of [thyroid](#) dysfunction (e.g., temperature intolerance, weight changes, [constipation/diarrhea](#)).

C - Thrombus formation in a deep vein

Explanation But

The formation of a [pulmonary embolism](#) from a [DVT](#) could also cause [shortness of breath](#), although the presentation is typically acuter than seen in this patient.

Explanation Why

[Deep venous thrombosis](#) can cause decreased venous drainage and [edema](#) in the affected extremity. However, this woman's bilateral lower extremity [edema](#) would have to be caused by bilateral thromboses, which is very rare.

D - Increased capillary hydrostatic pressure

Explanation But

Reduced [cardiac output](#) also activates [RAAS](#), leading to a second mechanism for [edema](#) formation in patients with [congestive heart failure](#): renal sodium and water retention.

Explanation Why

Increased [capillary hydrostatic pressure](#) occurs in [right-sided heart failure](#) in which reduced [right ventricular](#) output leads to systemic venous congestion. Venous congestion increases the [capillary hydrostatic pressure](#), altering the physiological [Starling forces](#) to cause fluid leakage into the [interstitium](#) and [edema](#).

E - Decreased plasma oncotic pressure

Explanation Why

Decreased plasma [oncotic pressure](#) is seen in patients with conditions that cause low levels of [plasma proteins](#), especially [albumin](#) (e.g., [liver](#) failure, [nephrotic syndrome](#)). Low levels of protein in the blood alter the physiological [Starling forces](#) to cause fluid leakage out of [capillaries](#), resulting in generalized peripheral [edema](#). This patient's [right heart failure](#) has no effect on total protein levels in the bloodstream and therefore has no effect on [oncotic pressure](#).

Question # 39

A 57-year-old woman is admitted to the intensive care unit for management of shock. Her pulse is feeble and blood pressure is 86/45 mm Hg. The patient undergoes pulmonary artery catheterization which shows an elevated pulmonary capillary wedge pressure and increased systemic vascular resistance. Which of the following additional findings is most likely in this patient?

	Answer	Image
A	Cold skin due to loss of intravascular fluid volume	
B	Bradycardia due to neurologic dysfunction	
C	Bronchospasm due to excessive histamine release	
D	Mottled skin due to release of endotoxins	
E	Confusion due to decreased stroke volume	

Hint

This patient might have presented with substernal chest pain radiating to her jaw.

Correct Answer

A - Cold skin due to loss of intravascular fluid volume

Explanation Why

Loss of intravascular fluid volume is consistent with [hypovolemic shock](#). In [hypovolemic shock](#), the [sympathetic nervous system](#) is activated, which causes vasoconstriction and thus an increased SVR. Although SVR is increased in this patient, her [PCWP](#) is inconsistent with [hypovolemic shock](#). [Hypovolemic shock](#) would result in a decreased [PCWP](#) due to [preload](#) reduction from intravascular depletion.

B - Bradycardia due to neurologic dysfunction

Explanation Why

[Bradycardia](#) due to neurologic dysfunction is consistent with [increased intracranial pressure](#) ([Cushing reflex](#)) and [spinal cord](#) damage above T₆ level. [Spinal cord](#) damage can cause [hypotension](#) and feeble peripheral pulses ([neurogenic shock](#)). However, [neurogenic shock](#) would be associated with decreased SVR (due to peripheral [vasodilation](#)) and decreased [PCWP](#) (due to [preload](#) reduction from intravascular depletion). Patients with [increased ICP](#) would have [hypertension](#).

C - Bronchospasm due to excessive histamine release

Explanation Why

Bronchospasm due to excessive [histamine](#) release is consistent with [anaphylactic shock](#). [Anaphylactic shock](#) is caused by a [type I hypersensitivity reaction](#) that may lead to circulatory failure ([distributive shock](#)). In [anaphylactic shock](#), the SVR and [PCWP](#) are both decreased due to [histamine](#)-mediated [vasodilation](#) and [capillary](#) leak.

D - Mottled skin due to release of endotoxins

Explanation Why

Release of [endotoxins](#) is consistent with [septic shock](#). [Septic shock](#) results from systemic [vasodilation](#) and [capillary](#) leakage. Initially, the [skin](#) is typically warm and flushed, but as [shock](#) progresses, the [skin](#) becomes mottled, cool and pale. However, [septic shock](#) typically results in decreased SVR (due to peripheral [vasodilation](#)) and decreased [PCWP](#) (due to decreased [left ventricular preload](#) caused pooling of blood in peripheral circulation).


E - Confusion due to decreased stroke volume

Explanation Why

Confusion due to decreased [stroke volume](#) is consistent with [cardiogenic shock](#), which is the most likely cause of [shock](#) in this patient. Poor [left ventricular](#) output results in elevated [pulmonary capillary wedge pressure](#) ([PCWP](#)). In addition, [hypotension](#) activates the [sympathetic nervous system](#), resulting in vasoconstriction and increased [systemic vascular resistance](#) (SVR). Peripheral vasoconstriction also causes cold, clammy extremities.

Question # 40

A 37-year-old woman comes to the physician for increasing shortness of breath with exercise over the past 2 months. She used to walk around her neighborhood every evening, but now has difficulty carrying out her daily activities. She has also started sleeping with three pillows at night. She remembers being hospitalized due to a serious illness during her childhood but has not seen a doctor since emigrating from Bangladesh 7 years ago. Physical examination shows crackles at the bilateral lung bases. There is an opening snap followed by a late diastolic rumble heard best at the 5th intercostal space at the midclavicular line. Which of the labeled points in the overview best corresponds to the physiologic opening time of the heart valve that is affected in this patient?



	Answer	Image
A	A	
B	B	

	Answer	Image
C	C	
D	D	
E	E	The image displays a series of four vertically stacked graphs sharing a common x-axis representing time. The top graph shows pressure in mmHg (0 to 120) with curves for aortic pressure (orange), left ventricular pressure (blue), and left atrial pressure (green). The middle graph shows volume in ml (0 to 150) with curves for stroke volume (red) and left ventricular volume (blue). The bottom graph shows the propagation of excitation (ECG) and heart sounds (S1, S2). The cycle is divided into phases: Atrial contraction, Ventricular contraction, Relaxation, and Diastole. Key points 1, 2, 3, and 4 are marked on the pressure and volume curves.

Hint

This patient's features of left heart failure (e.g., exertional dyspnea, orthopnea) and heart murmur indicate mitral stenosis. Her history of childhood illness suggests rheumatic fever, a known cause of mitral stenosis.

Correct Answer

A - A

Explanation Why

Point A corresponds to atrial [systole](#), which occurs in late ventricular [diastole](#) when the [atria](#) contract to eject blood into the ventricle. Atrial contraction can be associated with an [S4 heart sound](#), which is generated by increased atrial pressure due to contraction against a noncompliant ventricle (e.g., due to ventricular [hypertrophy](#)).

B - B

Explanation Why

Point B corresponds to the [mitral valve](#) closure, marking the beginning of [systole](#). The interval between the closure of the [mitral valve](#) and the opening of the [aortic valve](#) is the [isovolumetric contraction](#) phase, in which [left ventricular](#) pressure increases significantly before the [aortic valve](#) opens and [systolic ejection](#) starts. Impaired closure of the [mitral valve](#) during [systole](#) (e.g., due to [infective endocarditis](#)) causes [mitral regurgitation](#) (MR). MR is characterized by a [holosystolic murmur](#), which is best heard at the apex and radiates to the [axilla](#).

C - C

Explanation Why

Point C corresponds to the [aortic valve](#) opening, which follows [isovolumetric contraction](#) and marks the beginning of the [systolic ejection](#) phase of the [cardiac cycle](#). Impaired opening of the [aortic valve](#) is seen in [aortic stenosis](#), which is often caused by [aortic valve](#) calcification in elderly persons. [Aortic stenosis](#) is characterized by a harsh crescendo-decrescendo [systolic ejection murmur](#) and a soft S2 [heart sound](#) that is best heard at the right 2nd intercostal space.


D - D

Explanation Why

Point D corresponds to the [aortic valve](#) closure. The closure of the [aortic valve](#) corresponds to the [2nd heart sound](#) and marks the end of [systole](#) and the beginning of [diastole](#). The interval between the closure of the [aortic valve](#) and the opening of the [mitral valve](#) is the [isovolumetric relaxation](#) phase, in which the [left ventricular](#) pressure drops significantly before the [ventricular filling](#) starts during [diastole](#). Impaired closure of the [aortic valve](#) (e.g., due to a [bicuspid aortic valve](#)) causes [aortic regurgitation](#), which is characterized as a high-pitched early [diastolic](#) decrescendo [murmur](#) that is best heard at the left 3rd intercostal space ([Erb's point](#)).

E - E

Image



Explanation Why

Point E corresponds to the [mitral valve](#) opening. The opening of the [mitral valve](#) follows [isovolumetric relaxation](#) of the ventricles and marks the beginning of the rapid [filling phase](#) of [diastole](#). In [mitral stenosis](#), [mitral valve](#) opening is impaired, which is associated with a decreased [left ventricular](#) filling and a decreased [left ventricular end-diastolic volume \(EDV\)](#). Decreased [EDV](#) results in decreased [stroke volume](#) and decreased [cardiac output](#), which causes this patient's features of [left heart failure](#).

Question # 1

A 54-year-old woman comes to the physician because of a 6-month history of dull, persistent pain and swelling of her right leg. The pain is worse at the end of the day and is relieved by walking or elevating her feet. Two years ago, she developed acute deep vein thrombosis in her right calf after a long flight, which was treated with anticoagulants for 6 months. Physical examination shows 2+ pitting edema of her right leg. The skin around the right ankle shows a reddish-brown discoloration and multiple telangiectasias. She has dilated varicose veins in the right leg. Which of the following is most likely to establish the diagnosis?

	Answer	Image
A	Computerized tomography scan with contrast	
B	Nerve conduction studies	
C	D-dimer assay	
D	Ankle-brachial pressure index	
E	Duplex ultrasonography	

Hint

This patient's unilateral chronic dull pain, edema, dilated veins in the lower extremity, and history of DVT suggest chronic venous insufficiency.

Correct Answer

A - Computerized tomography scan with contrast

Explanation Why

[Computed tomography scan](#) with contrast of the lower extremities is used to evaluate patients for deep infections (e.g., [abscesses](#)), [malignancy](#), or injury in the lower extremities. It is not used to evaluate for [chronic venous insufficiency](#).

B - Nerve conduction studies

Explanation Why

[Nerve conduction studies](#) are used to evaluate the function of the [peripheral nervous system](#). They are particularly critical for diagnosing polyneuropathies (e.g., due to chronic [diabetes mellitus](#)). Although a [polyneuropathy](#) can manifest with asymmetric or unilateral [pain](#), which is seen in this patient, the [pain](#) usually takes the form of a buzzing or tingling sensation, not a dull ache. Moreover, a [polyneuropathy](#) would not cause the [edema](#) and [telangiectasias](#) seen in this patient.

C - D-dimer assay

Explanation Why

The [D-dimer](#) assay is a nonspecific test that is indicated for patients with a low [pretest probability](#) of acute [deep venous thrombosis](#) ([DVT](#)) to rule out an acute thrombotic event. The test has a high [negative predictive value](#) (> 95%); therefore, a negative test result effectively rules out the presence of a [deep venous thrombosis](#) or a [pulmonary embolism](#).

D - Ankle-brachial pressure index

Explanation Why

The [ankle-brachial pressure index \(ABPI\)](#) is used to screen for [peripheral arterial disease \(PAD\)](#) in patients who present with lower limb [pain](#) that suggests [claudication pain](#) (i.e., [pain](#) on walking that is relieved with rest). This patient's [pain](#) improves with walking, making [PAD](#) an unlikely cause of her symptoms. Her symptoms are consistent with a pathology of the venous system, which cannot be evaluated with [ABPI](#).

E - Duplex ultrasonography

Explanation Why

[Duplex ultrasonography](#) ([compression ultrasonography](#)) is the first-line investigation in patients with features suggestive of [chronic venous insufficiency \(CVI\)](#), as seen here. Evidence of venous reflux on [duplex ultrasonography](#) confirms the diagnosis of [CVI](#). [Duplex ultrasonography](#) can help determine the course of treatment by identifying which valves are incompetent and assessing the patency of the previously thrombosed deep veins.

Question # 2

A 62-year-old man is brought to the emergency department because of a 4-hour history of abdominal pain, nausea, vomiting, and confusion. His wife reports that he had blurry vision on the way to the hospital. Two weeks ago, he lost his job and since then has been extremely worried about their financial situation and future. He has congestive heart failure and atrial fibrillation well controlled with combination medical therapy. His temperature is 36.5°C (97.7°F), pulse is 57/min and irregular, respirations are 14/min, and blood pressure is 118/63 mm Hg. The patient is oriented only to person. Serum studies show:

Na ⁺	138 mEq/L
Cl ⁻	100 mEq/L
K ⁺	5.3 mEq/L
HCO ₃ ⁻	25 mEq/L
Blood urea nitrogen	14 mg/dL
Creatinine	0.9 mg/dL

An ECG shows premature ventricular beats. The drug most likely responsible for this patient's symptoms has which of the following mechanisms of action?

	Answer	Image
A	Inhibition of funny channels	
B	Inhibition of Na ⁺ -K ⁺ -2Cl ⁻ -cotransporters	
C	Blockade of aldosterone receptors	

	Answer	Image
D	Inhibition of Na^+/K^+ -ATPase	
E	Blockade of beta-adrenergic receptors	
F	Inhibition of angiotensin-converting enzyme	
G	Blockade of Na^+ channels	

Hint

This patient's symptoms of abdominal pain, nausea, vomiting, vision disturbance, confusion, bradycardia, premature ventricular beats, and hyperkalemia suggest digoxin toxicity.

Correct Answer

A - Inhibition of funny channels

Explanation Why

[Ivabradine](#) is a drug that inhibits the [funny current](#) responsible for cardiac pacemaker activity and thus reduces [heart rate](#). While this medication is prescribed as a second-line agent for symptomatic treatment of patients with [heart failure](#), [ivabradine](#) overdose would likely cause profound [bradycardia](#) rather than the symptoms seen here.

B - Inhibition of $\text{Na}^+\text{-K}^+\text{-2Cl}^-$ -cotransporters

Explanation Why

[Loop diuretics](#) (e.g., [furosemide](#)) are drugs that inhibit [\$\text{Na}^+\text{-K}^+\text{-2Cl}^-\$ cotransporters](#) in the thick [ascending loop of Henle](#) within [nephrons](#). This leads to increased production of [urine](#) and a subsequent reduction of blood volume and cardiac [preload](#). While this medication is commonly prescribed to patients with [heart failure](#), [loop diuretic](#) overdose would likely cause [hypotension](#), [hyponatremia](#), [hypokalemia](#), [hypocalcemia](#), and [metabolic alkalosis](#) rather than the symptoms seen here.

C - Blockade of aldosterone receptors

Explanation Why

[Aldosterone receptor antagonists](#) (e.g., [spironolactone](#), [eplerenone](#)) [competitively inhibit aldosterone](#) receptors in the [distal](#) tubule and [collecting duct](#) of [nephrons](#). This leads to decreased Na^+ reabsorption and K^+ excretion, subsequently causing diuresis. While [potassium-sparing diuretics](#) such as [aldosterone receptor antagonists](#) are commonly prescribed to patients with [heart failure](#) and can cause [hyperkalemia](#), an overdose would likely cause [hypotension](#), [hyponatremia](#), and [metabolic acidosis](#) rather than the symptoms seen here.

D - Inhibition of Na^+/K^+ -ATPase

Explanation Why

[Cardiac glycosides](#) are drugs that inhibit Na^+/K^+ -ATPases, and [digoxin](#) is the only drug of this class commonly used clinically as a positive [inotrope](#) in [heart failure](#). [Digoxin](#) has a [narrow therapeutic index](#), and this patient exhibits symptoms and laboratory findings consistent with [digoxin poisoning](#). Given his recent life stressors (job loss, financial worries), the possibility that this patient overdosed on his prescribed [digoxin](#) to attempt [suicide](#) should be considered.

E - Blockade of beta-adrenergic receptors

Explanation Why

[Beta blockers](#) (e.g., [metoprolol](#)) are drugs that inhibit [sympathetic](#) activation of [beta-adrenergic](#) receptors. They reduce blood pressure by decreasing [heart rate](#), reducing contractility, and causing peripheral vasoconstriction. While [beta blockers](#) are commonly prescribed to patients with [heart failure](#) and can cause [hyperkalemia](#) at high doses, overdose would likely cause [hypotension](#), [bradycardia](#), bronchospasm, and [hypoglycemia](#) rather than the symptoms seen here.

F - Inhibition of angiotensin-converting enzyme

Explanation Why

ACE inhibitors (e.g., [enalapril](#)) reduce the conversion of [angiotensin I](#) to [angiotensin II](#) and thereby reduce blood pressure as well as cardiac [preload](#) and [afterload](#). While this medication is commonly prescribed to patients with [heart failure](#), ACE inhibitor overdose would likely cause severe [hypotension](#) rather than the symptoms seen here.

G - Blockade of Na⁺ channels

Explanation Why

[Epithelial sodium channel blockers](#) (e.g., [triamterene](#) and [amiloride](#)) directly inhibit [epithelial](#) Na⁺ channels in the [distal](#) tubule and [collecting duct](#) of [nephrons](#). This decreases Na⁺ reabsorption and K⁺ excretion, which causes diuresis. Though [potassium-sparing diuretics](#) such as [epithelial sodium channel blockers](#) are commonly prescribed to patients with [heart failure](#) and can cause [hyperkalemia](#), an overdose would likely cause [hypotension](#), [hyponatremia](#), and [metabolic acidosis](#) rather than the symptoms seen here.

Question # 3

A 25-year-old woman is brought to the emergency department 12 hours after ingesting 30 tablets of an unknown drug in a suicide attempt. The tablets belonged to her father, who has a chronic heart condition. She has had nausea and vomiting. She also reports blurring and yellowing of her vision. Her temperature is 36.7°C (98°F), pulse is 51/min, and blood pressure is 108/71 mm Hg. Abdominal examination shows diffuse tenderness with no guarding or rebound. Bowel sounds are normal. An ECG shows prolonged PR-intervals and flattened T-waves. Further evaluation is most likely to show which of the following electrolyte abnormalities?

	Answer	Image
A	Increased serum Na^+	
B	Decreased serum K^+	
C	Increased serum Ca^{2+}	
D	Decreased serum Ca^{2+}	
E	Decreased serum Na^+	
F	Increased serum K^+	

Hint

Abdominal pain, nausea, vomiting, and yellow-tinged, blurry vision are hallmarks of digoxin intoxication, a cardiac glycoside. The ECG findings support this diagnosis.

Correct Answer

A - Increased serum Na^+

Explanation Why

[Digoxin](#) inhibits the [\$\text{Na}^+/\text{K}^+\$ -ATPase](#), which leads to decreased Na^+ efflux from the cells. Decreased Na^+ efflux reduces the efficacy of the [\$\text{Na}^+/\text{Ca}^{2+}\$ exchanger](#), which usually facilitates Na^+ influx and [\$\text{Ca}^{2+}\$ efflux](#). The reduced activity of both pumps increases cardiac [inotropy](#), while preserving the intracellular and extracellular concentrations of sodium. [Hypernatremia](#) would not be expected.

B - Decreased serum K^+

Explanation Why

[Digoxin](#), which is responsible for this patient's symptoms, inhibits [\$\text{Na}^+/\text{K}^+\$ -ATPase](#), but it does not decrease extracellular potassium levels. While [hypokalemia](#) can also manifest with flattened [T-waves](#) on [ECG](#), this patient's [ECG](#) findings are more likely a direct effect of [digoxin](#) use and not related to her serum potassium level.

C - Increased serum Ca^{2+}

Explanation Why

[Digoxin](#) inhibits [\$\text{Na}^+/\text{K}^+\$ -ATPase](#), which reduces efflux of Na^+ out of the cell and thereby reduces the activity of [\$\text{Na}^+/\text{Ca}^{2+}\$ exchangers](#). In [digoxin toxicity](#), which is responsible for this patient's symptoms, the concentration of [\$\text{Ca}^{2+}\$](#) therefore increases within the cells, but not in the serum.

D - Decreased serum Ca^{2+}

Explanation Why

[Digoxin](#) indirectly increases intracellular Ca^{2+} , which is the desired effect at non-toxic concentrations. However, [digoxin](#) does not significantly decrease serum Ca^{2+} levels.

E - Decreased serum Na^+

Explanation Why

[Digoxin](#), which is responsible for this patient's symptoms, inhibits [\$\text{Na}^+/\text{K}^+\$ -ATPase](#), which leads to decreased Na^+ efflux from the cells. However, decreased Na^+ efflux reduces the efficacy of the [\$\text{Na}^+/\text{Ca}^{2+}\$ exchanger](#), thus preserving rather than decreasing extracellular sodium concentrations.

F - Increased serum K^+

Explanation But

[ECG](#) findings seen in [digitalis](#) toxicity do not always correlate with other clinical findings of toxicity and while they may be seen with acute toxicity, they are more commonly seen with chronic [digoxin](#) use.

Explanation Why

[Digoxin](#), a [cardiac glycoside](#), inhibits [\$\text{Na}^+/\text{K}^+\$ -ATPase](#), which leads to decreased K^+ influx into the cells and if taken in excess, [hyperkalemia](#). Inhibition of [\$\text{Na}^+/\text{K}^+\$ -ATPase](#) also decreases Na^+ efflux from the cell, which decreases the activity of the [\$\text{Na}^+/\text{Ca}^{2+}\$ exchanger](#). The resulting increase in intracellular calcium levels results in increased cardiac [inotropy](#) and risk of [cardiac arrhythmias](#). While [hyperkalemia](#) typically causes peaked [T-waves](#), this patient's [ECG](#) findings of flattened [T-waves](#) and a prolonged [PR interval](#) are a direct manifestation of [digoxin](#) on the atrial and ventricular

refractory periods and the [AV node](#) conduction velocity. Other [ECG](#) findings seen in [digitalis](#) use include downsloping ST-depressions and [QT interval](#) shortening. [Cardiac glycoside poisoning](#) can cause severe [arrhythmias](#) (e.g., [AV block](#)).

Question # 4

A 62-year-old woman comes to the physician because of a 2-month history of a rash on her ankles with intermittent itching. After the rash developed, she started applying a new scented lotion to her legs daily. She works as a cashier at a grocery store. She has type 2 diabetes mellitus and hypertension. Current medications include metformin and enalapril. Examination shows enlarged superficial veins of the right lower extremity and red-brown discoloration with indistinct margins over the medial ankles. There is 1+ edema in the lower extremities. Which of the following is the most likely cause of this patient's skin findings?

	Answer	Image
A	Bacterial spread through the superficial dermis	
B	Contact of antigen with pre-sensitized T lymphocytes	
C	Intracellular accumulation of sorbitol	
D	Insufficient arterial tissue perfusion	

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

	Answer	Image
E	Infection with dermatophyte	
F	Dermal deposition of hemosiderin	

Hint

This woman has a pruritic, red-brown rash over the medial ankles, varicose veins, and bilateral lower extremity edema. Her occupation likely requires her to stand for prolonged periods. These findings suggest stasis dermatitis due to chronic venous disease.

Correct Answer

A - Bacterial spread through the superficial dermis

Image



Explanation Why

Bacterial spread through the [superficial dermis](#) and [lymphatics](#) is the underlying cause of [erysipelas](#), which typically results in lower extremity [erythema](#) and [edema](#) and is more common in individuals with [diabetes mellitus](#) (due to impaired [neutrophil](#) function from [hyperglycemia](#) and [diabetic neuropathy](#)). However, [skin](#) lesions of [erysipelas](#) are usually well-demarcated, unilateral, nonpruritic, and associated with systemic manifestations (e.g., [fever](#) and chills).

B - Contact of antigen with pre-sensitized T lymphocytes

Image



Explanation Why

Contact of antigen with pre-sensitized [T lymphocytes](#) is the underlying mechanism of [allergic contact dermatitis](#), which usually manifests with a [pruritic](#), [erythematous](#) rash with indistinct margins. Products that contain fragrance are common contact [allergens](#). However, this patient's rash preceded her use of a new lotion. Additionally, [allergic contact dermatitis](#) typically only causes [edema](#) in areas that have thin [skin](#) (e.g., the [lips](#) and eyelids) and does not result in [varicose veins](#).

C - Intracellular accumulation of sorbitol

Explanation Why

Intracellular accumulation of [sorbitol](#) is involved in the pathogenesis of [diabetic neuropathy](#), a condition that causes abnormal sensation in the lower extremities in patients with [diabetes](#). [Sorbitol](#)

accumulates in cells that lack [sorbitol dehydrogenase](#) (e.g., [Schwann cells](#)), which leads to osmotic damage. However, [diabetic neuropathy](#) does not cause [edema](#), [varicose veins](#), or red-brown discoloration.

D - Insufficient arterial tissue perfusion

Explanation Why

Insufficient arterial tissue perfusion can cause [ischemic](#) ulcers, which often manifest at sites of focal pressure such as the [medial](#) malleoli. These ulcers develop in patients with [peripheral artery disease \(PAD\)](#), which is associated with advanced age, [diabetes](#), and [hypertension](#). However, they are typically painful and well-demarcated. Additionally, this patient does not have other common symptoms of [PAD](#) such as [claudication](#), [hair](#) loss on the leg, or [nail](#) changes.

E - Infection with dermatophyte

Image



Explanation Why

[Tinea pedis](#), a [dermatophyte infection](#), most commonly develops on the feet and causes a [pruritic](#), scaly rash. [Erythema](#) can also occur but is less common. [Tinea pedis](#) typically affects the digital interspaces and soles of the feet (not the [medial](#) ankles) and is not associated with [edema](#) or [varicose veins](#).

F - Dermal deposition of hemosiderin

Image



Explanation Why

[Dermal](#) deposition of [hemosiderin](#) is a key feature of [stasis dermatitis](#). Elevated venous pressures from [chronic venous disease](#) result in [red blood cell](#) breakdown, which leads to [hemosiderin](#) production. [Hemosiderin](#) penetrates defective [capillaries](#) and accumulates in the [dermis](#), which causes a [pruritic](#), red-brown rash, as seen in this patient. The [medial](#) ankles are the most frequently involved site.

Question # 5

A previously healthy 24-year-old male is brought to the emergency department because of fevers, congestion, and chest pain for 3 days. The chest pain is exacerbated by deep inspiration. He takes no medications. His temperature is 37.5°C (99.5°F), blood pressure is 118/75 mm Hg, pulse is 130/min, and respirations are 12/min. He appears weak and lethargic. Cardiac examination shows a scratchy sound best heard along the left sternal border when the patient leans forward. There are crackles in both lung bases. Examination of the lower extremities shows pitting edema. Results of a rapid influenza test are negative. EKG shows diffuse ST-elevations with depressed PR interval. An echocardiogram shows left ventricular chamber enlargement with contractile dysfunction. Infection with which of the following pathogens is the most likely cause of this patient's symptoms?

	Answer	Image
A	Togavirus	
B	Paramyxovirus	
C	Flavivirus	
D	Orthomyxovirus	
E	Picornavirus	

Hint

This patient has evidence of both pericarditis and myocarditis (i.e., inflammation of the heart muscle and pericardium). The most commonly identified pathogen responsible for myocarditis belongs to the same family as poliovirus and echovirus.

Correct Answer

A - Togavirus

Explanation Why

Infection with [togaviruses](#) (e.g., [rubella](#) virus, western and [eastern equine encephalitis](#) virus, [chikungunya virus](#)) is not a common cause of viral [myocarditis](#) or [pericarditis](#). This patient lacks [risk factors](#) for Western and [Eastern equine encephalitis](#) and [chikungunya virus](#) (e.g., mosquito bites, travel to [endemic](#) areas), and has no signs of [rubella](#) infection (e.g., [lymphadenopathy](#), polyarthrititis, [skin](#) rash).

B - Paramyxovirus

Explanation Why

Infections with [paramyxoviruses](#) (e.g., [parainfluenza virus](#), [RSV](#), [measles virus](#), [mumps virus](#)) can present with [fever](#) and congestion, as seen in this patient, but they do not commonly cause [myocarditis](#) or [pericarditis](#). [Croup](#), [measles](#), and [mumps](#) are all caused by [paramyxovirus](#) infection.

C - Flavivirus

Explanation Why

Infections with [flaviviruses](#) (e.g., [dengue](#) virus, [yellow fever virus](#), [St. Louis encephalitis virus](#), [West Nile virus](#), [Zika virus](#), and [hepatitis C virus](#)) do not commonly cause [myocarditis](#) or [pericarditis](#). Furthermore, this patient lacks typical symptoms and [risk factors](#) (e.g., exposure to *Aedes aegypti* mosquito, IV drug use, etc) for [flavivirus](#) infection.

D - Orthomyxovirus

Explanation Why

Infections with [orthomyxoviruses](#) (e.g, [influenza](#) virus) can present with [fever](#), [lethargy](#), and congestion, as seen in this patient. While [influenza](#) is not a common cause of [myocarditis](#), the H1N1 [influenza A](#) strain has been implicated in cases of fulminant [myocarditis](#). This patient's negative [influenza](#) testing and lack of other typical symptoms (e.g., myalgias) make this unlikely.


E - Picornavirus

Explanation Why

[Coxsackie B](#) is one of the most commonly identified infectious cause of viral [myocarditis](#). It belongs to the [picornavirus](#) family and is transmitted via airborne droplets or the fecal-oral route. Other viral pathogens that can cause [myocarditis](#) include [parvovirus B19](#), [HHV-6](#), [HIV](#), and [adenovirus](#).

Question # 6

Two weeks after undergoing an emergency cardiac catheterization for unstable angina pectoris, a 65-year-old man has decreased urinary output. He takes naproxen for osteoarthritis and was started on aspirin, clopidogrel, and metoprolol after the coronary intervention. His temperature is 38.1°C (100.5°F), pulse is 96/min, and blood pressure is 128/88 mm Hg. Examination shows mottled, reticulated purplish discoloration of the feet and ischemic changes on the right big toe. His leukocyte count is 16,500/mm³ with 56% segmented neutrophils, 12% eosinophils, 30% lymphocytes, and 2% monocytes. His serum creatinine concentration is 4.5 mg/dL. A photomicrograph of a kidney biopsy specimen is shown. Which of the following is the most likely cause of this patient's presentation?



	Answer	Image
A	Allergic interstitial nephritis	
B	Cholesterol embolization	

	Answer	Image
C	Contrast-induced nephropathy	
D	Renal papillary necrosis	
E	Eosinophilic granulomatosis with polyangiitis	

Hint

This condition can also be seen in patients undergoing endovascular abdominal aortic aneurysm repair.

Correct Answer

A - Allergic interstitial nephritis

Explanation Why

[Allergic interstitial nephritis](#) ([AIN](#)) is caused by a [hypersensitivity reaction](#) to medications such as [NSAIDs](#) or [penicillins](#). Unlike here, patients present with [fever](#), rash, [eosinophilia](#), eosinophiluria, and [sterile pyuria](#). Although this patient's symptoms overlap with those of [AIN](#), the presence of [blue toe syndrome](#) and the biopsy results rule out this diagnosis.

B - Cholesterol embolization

Explanation Why

[Cholesterol embolization syndrome](#) is a known complication of [cardiac catheterization](#), vascular interventions, and [anticoagulant](#) therapy. [Cholesterol](#) is released into the circulation, where it can travel to many organs, especially the [kidneys](#) (as they receive a large amount of the blood supply). This patient presents with the classic symptoms of [blue toe syndrome](#), [livedo reticularis](#), and [acute kidney failure](#). [Eosinophilia](#) and eosinophiluria in the setting of [cholesterol embolism](#) are very suggestive findings. The [kidney biopsy](#) shows intravascular [cholesterol](#) deposits (spindle-shaped vacuoles) with multiple intra-arterial '[cholesterol](#) clefts' and associated giant-cells, which confirms the diagnosis.

C - Contrast-induced nephropathy

Explanation Why

In [contrast-induced nephropathy](#), [creatinine](#) is highest after 3–5 days and usually normalizes to the patient's baseline within 1 week. [Nephrotoxic drugs](#), such as [NSAIDs](#), increase the risk of [kidney](#) damage following contrast administration. Although this patient takes [naproxen](#) regularly, neither the onset of symptoms nor the [renal biopsy](#) results is consistent with this diagnosis.

D - Renal papillary necrosis

Explanation Why

[Kidney biopsy](#) of a patient with [renal papillary necrosis](#) would reveal [coagulative necrosis](#) of the [renal medulla](#) and papillae rather than spindle-shaped vacuoles. Unlike here, patients with [renal papillary necrosis](#) typically present with [hematuria](#), flank [pain](#), [fever](#), and colicky [pain](#). [Urinalysis](#) may show [pyuria/WBC casts](#) without [eosinophils](#). In addition, [skin](#) findings and [blue toe syndrome](#) are not expected in [renal papillary necrosis](#).


E - Eosinophilic granulomatosis with polyangiitis

Explanation Why

[Eosinophilic granulomatosis with polyangiitis](#) is a [vasculitis](#) characterized by necrotizing [granulomas](#) with [eosinophilia](#). It commonly involves the [lungs](#), [skin](#), and [kidneys](#). It classically presents with severe allergic asthma attacks and [allergic rhinitis](#), neither of which is present in this patient. In addition, a biopsy of [eosinophilic granulomatosis with polyangiitis](#) would show eosinophilic tissue with granulomatous reactions.


Question # 7

A 3-month-old boy is brought to the physician by his mother because of poor weight gain. She also reports a dusky blue discoloration to his skin during feedings and when crying. On examination, there is a harsh, systolic murmur heard over the left upper sternal border. An x-ray of the chest is shown below. Which of the following is the most likely cause of his symptoms?



	Answer	Image
A	Persistent connection between the aorta and pulmonary artery	1 2 3

	Answer	Image
B	Hypoplasia of the left ventricle	
C	Narrowing of the distal aortic arch	Diagram 1: Patent ductus arteriosus with stenosis. ↑BP ~SaO₂ (upper body), ~BP ↓SaO₂ (lower body). Diagram 2: Obliterated ductus arteriosus. ↑BP ~SaO₂ (upper body), ↓BP ~SaO₂ (lower body).
D	Right ventricular outflow obstruction	Diagram 1: Normal flow. Diagram 2: Obstruction at the pulmonary valve.
E	Anatomic reversal of aorta and pulmonary artery	Patent ductus arteriosus, Atrial septal defect.

	Answer	Image
F	Tricuspid valve regurgitation	 The diagram illustrates the heart with two specific congenital conditions highlighted. A 'Patent ductus arteriosus' is shown as a persistent connection between the aorta and the pulmonary artery, with a white arrow indicating the direction of blood flow from the aorta to the pulmonary artery. A 'Patent foramen ovale' is shown as an opening in the interatrial septum, with a white arrow indicating the direction of blood flow from the right atrium to the left atrium. The heart is colored in shades of red and purple, and the surrounding vessels are blue.


Hint

The boot-shaped heart on this boy's x-ray is caused by right ventricular hypertrophy (RVH). His harsh, systolic murmur heard over the left upper sternal border is caused by pulmonary stenosis. These findings, along with his acute hypoxic spells (which cause cyanosis), indicate a diagnosis of tetralogy of Fallot (TOF).

Correct Answer

A - Persistent connection between the aorta and pulmonary artery

Image



Explanation Why

A persistent connection between the aorta and [pulmonary artery](#) characterizes a [patent ductus arteriosus \(PDA\)](#), an [acyanotic heart disease](#) that can cause [cyanosis](#) if [Eisenmenger syndrome](#) develops. However, patients with a [PDA](#) and [Eisenmenger syndrome](#) typically develop [differential cyanosis](#) because [right-to-left shunting](#) through the [ductus arteriosus](#) delivers unoxygenated blood [distal](#) to the [left subclavian artery](#). Additionally, [PDA](#) causes a continuous, [machine-like murmur](#) heard throughout [systole](#) and [diastole](#) (unlike this patient's [systolic murmur](#)), and chest x-ray shows a prominent [mediastinum](#) caused by an enlarged [pulmonary artery](#) and [aortic knob](#) (not a boot-shaped heart).


B - Hypoplasia of the left ventricle

Explanation Why

Hypoplasia of the left ventricle is caused by hypoplastic left heart syndrome (HLHS), a heart condition that is typically acyanotic in the early postnatal period because a patent ductus arteriosus (PDA) ensures systemic perfusion. Cyanosis develops when the PDA closes, usually within a few days of birth (not as late as 3 months as in this case). Neonates with HLHS also have weak peripheral pulses, hypotension, and metabolic acidosis as the hypoplastic left ventricle cannot maintain adequate cardiac output. Additionally, HLHS does not cause a murmur, and findings on chest x-ray are variable (e.g., small, normal, or enlarged heart and increased pulmonary venous markings) but do not include a boot-shaped heart.

C - Narrowing of the distal aortic arch

Image




Explanation Why

Narrowing of the [distal aortic arch](#), or [coarctation of the aorta](#), manifests variably depending on the degree of [aortic stenosis](#) (AS) and the presence of a [patent ductus arteriosus](#) (PDA). Patients with significant AS and PDA manifest with [differential cyanosis](#) because [right-to-left shunting](#) through the [ductus arteriosus](#) delivers unoxygenated blood [distal](#) to the [left subclavian artery](#). Once the PDA closes (usually within a few days after [birth](#)), [heart failure](#) develops and manifests as [dyspnea](#), sweating during feeds, and poor weight gain. Examination shows pallor (rather than [cyanosis](#)), features of inadequate [left ventricular](#) output (absent or weak femoral pulses, brachio-femoral delay), and signs of [right heart failure](#) (e.g., raised [jugular venous pressure](#), [edema](#), and [hepatomegaly](#)). Chest x-ray shows [cardiomegaly](#) and increased pulmonary venous markings, not a boot-shaped [heart](#).

D - Right ventricular outflow obstruction

Image



Explanation But

Older children with TOF tend to squat during a [tet spell](#). Squatting increases [systemic vascular resistance](#), which increases [left ventricular](#) pressure and decreases the [right-to-left shunting](#) of blood through the [VSD](#). [Infants](#) who are having a [tet spell](#) should be placed with their [knees](#) toward their


chest to achieve the same effect.

Explanation Why

Right ventricular outflow obstruction (RVOO) in this patient is caused by [pulmonary stenosis](#), which is one of the components of TOF, along with [RVH](#), an overriding aorta, and a [VSD](#). Exertion, such as crying or feeding, increases the RVOO. This results in [cyanosis](#) ([tet spells](#)) due to [right-to-left shunting](#) of deoxygenated blood through the [VSD](#) instead of the [pulmonary valve](#), which in turn causes the [murmur](#) to become softer. The [right ventricle hypertrophies](#) because it chronically pumps against high resistance caused by the RVOO.

E - Anatomic reversal of aorta and pulmonary artery

Image




Explanation Why

Anatomic reversal of the aorta and [pulmonary artery](#) is known as [transposition of the great vessels](#) ([TGV](#)). This condition causes [cyanosis](#) because the aorta arises from the [right ventricle](#) and thus results in [systemic circulation](#) of deoxygenated blood. However, unlike this patient, [cyanosis](#) in [TGV](#) manifests immediately after [birth](#). Additionally, chest x-ray in [TGV](#) shows an egg-on-a-string appearance due to an enlarged [heart](#) with a narrowed [mediastinum](#), which resembles an egg lying on

its side.

F - Tricuspid valve regurgitation

Image



Explanation Why

[Tricuspid regurgitation](#) (TR) in [infants](#) is usually due to a malformed and displaced [tricuspid valve](#) in [Ebstein anomaly](#). These [infants](#) may develop [cyanosis](#) only if there's an associated ASD or [PFO](#) that allows for right-to-left shunting of blood. Auscultation would reveal a [holosystolic murmur](#) at the left sternal border, like in this patient, with a widely [split S2](#). However, a chest [x-ray](#) in [Ebstein anomaly](#) will show [right atrial enlargement](#) (due to chronic reflux of blood from the [right ventricle](#) into the [right atrium](#)) - not a boot-shaped [heart](#) as seen in this [infant](#).

Question # 8

A 57-year-old man comes to the emergency department because of shortness of breath and palpitations for 3 hours. He has had similar episodes intermittently for 4 months. His pulse is 140/min and blood pressure is 90/60 mm Hg. An ECG shows irregular narrow-complex tachycardia with no discernable P waves. Emergent electrical cardioversion is performed and the patient reverts to normal sinus rhythm. Pharmacotherapy with sotalol is begun. Which of the following is the most likely physiologic effect of this drug?

	Answer	Image
A	Decreased AV nodal conduction	The diagram illustrates the effects of various antiarrhythmic drug classes on cardiac action potentials. It shows six panels: Class IA, Class IB, Class IC, Class II, Class III, and Class IV. Each panel plots Membrane potential (mV) against Time. A legend at the bottom identifies the blue curve as the Ventricular action potential and the pink curve as the Pacemaker cell action potential. Class IA: Prolonged AP duration, Slow conduction velocity. Class IB: Shortened AP duration, Slow conduction velocity. Class IC: Slow conduction velocity. Class II: Decreased slope of phase 4 repolarization, Prolonged repolarization. Class III: Prolonged AP duration. Class IV: Decreased slope of phase 4 and phase 0 repolarization, Prolonged repolarization.
B	Increased ventricular repolarization rate	This diagram shows the effect of Class IB antiarrhythmic drugs on the ventricular action potential. It plots Membrane potential (mV) against Time. The action potential is shown in blue. The diagram indicates a shortened AP duration and slow conduction velocity.
C	Decreased Purkinje fiber conduction	
D	Increased myocyte inotropy	

	Answer	Image
E	Increased K^+ efflux from myocytes	


Hint

Irregular narrow-complex tachycardia with no discernable P waves indicates atrial fibrillation. After multiple episodes of atrial fibrillation, this man was cardioverted and administered sotalol for continued rhythm control. Sotalol is a unique antiarrhythmic drug with both class II (beta blocker) and class III (potassium-channel blocker) properties.

Correct Answer

A - Decreased AV nodal conduction

Image



Explanation But


Decreased [AV nodal](#) conduction is also caused by non-dihydropyridine [calcium channel blockers](#) ([class IV antiarrhythmics](#)), [class IC antiarrhythmics](#) (e.g., [flecainide](#), [propafenone](#)), [digoxin](#), and [adenosine](#).

Explanation Why

Decreased [AV nodal](#) conduction is a potent effect of all [beta blockers](#) ([class II antiarrhythmics](#)), including [sotalol](#). As a consequence of [sympathetic](#) blockade (primarily [β1 receptors](#)), the conduction velocity through the [AV node](#) is reduced and the [heart rate](#) slows. Decreased [AV nodal](#) conduction would be evident on [ECG](#) as an increased [PR interval](#). Class III effects of [sotalol](#) include increased [myocyte action potential](#) duration (due to delayed [repolarization](#) caused by decreased K^+ efflux), increased [effective refractory period](#), and prolonged [QT interval](#).

B - Increased ventricular repolarization rate

Image



Explanation Why

Increased ventricular [repolarization](#) rate is seen with [class IB antiarrhythmics](#) (e.g., [lidocaine](#), [mexiletine](#)), which are typically used to treat [ventricular arrhythmias](#) in patients with [myocardial infarction](#), and [digitalis](#)-induced [arrhythmias](#). [Class III antiarrhythmic drugs](#), which block potassium channels in cardiac [myocytes](#), would decrease rather than increase the ventricular [repolarization](#) rate.

C - Decreased Purkinje fiber conduction

Explanation Why

Decreased conduction in cardiac [myocytes](#) and [Purkinje fibers](#) is the mechanism by which sodium channel blockers (class I antiarrhythmics) function. By inhibiting the flow of sodium ions, the rapid [depolarization](#) phase (phase 0) of cardiac [action potentials](#) is slowed. [Sotalol](#) is not a class I

antiarrhythmic and does not have a clinically apparent effect on conduction velocity in the [Purkinje fibers](#).

D - Increased myocyte inotropy

Explanation Why

Increased [myocyte inotropy](#) is caused by [digoxin](#), which can be used to treat [atrial fibrillation](#) in patients with [heart failure](#). [Sotalol](#) is an [antiarrhythmic drug](#) with the properties of a [beta blocker](#) ([class II antiarrhythmic drug](#)), and thus causes a decrease in [inotropy](#).

E - Increased K^+ efflux from myocytes

Explanation Why

[Hyperpolarization](#) due to increased K^+ efflux from [myocytes](#) is the mechanism of action of adenosine. Adenosine is used in the acute setting to terminate stable [paroxysmal supraventricular tachycardias](#) ([PSVT](#)) such as [AVNRT](#) and AVRT, which are typically regular narrow complex tachycardias. Adenosine should not be used to treat [atrial fibrillation](#) (AF) because it worsens AF. This patient received [sotalol](#), a [class III antiarrhythmic drug](#), which would slow the rate of [repolarization](#) by decreasing K^+ efflux from the [myocyte](#).

Question # 9

A 72-year-old man with congestive heart failure is brought to the emergency department because of chest pain, shortness of breath, dizziness, and palpitations for 30 minutes. An ECG shows a wide complex tachycardia with a P-wave rate of 105/min, an R-wave rate of 130/min, and no apparent relation between the two. Intravenous pharmacotherapy is initiated with a drug that prolongs the QRS and QT intervals. The patient was most likely treated with which of the following drugs?

	Answer	Image
A	Verapamil	
B	Carvedilol	
C	Flecainide	
D	Lidocaine	
E	Sotalol	
F	Quinidine	

Hint

The drug administered to this patient acts to block cardiac sodium channels and prolongs the action potential.

Correct Answer

A - Verapamil

Explanation Why

[Verapamil](#) is a [class IV antiarrhythmic](#). These drugs are used for controlling ventricular rate in [supraventricular tachycardia](#) and act on calcium-dependent cardiac cells, such as those in the [AV node](#). Class IV drugs reduce inward calcium current during the AP and phase 4. As a result, conduction velocity through the [AV node](#) is decreased and the [refractory period](#) is prolonged, which increases the [PR interval](#). [Verapamil](#) does not have an effect on QRS or QT duration, though.

B - Carvedilol

Explanation Why

[Carvedilol](#) is a nonselective [beta blocker](#)/alpha 1 blocker and a [class II antiarrhythmic](#). It blocks β_1 , β_2 , and [\$\alpha_1\$ receptors](#) on cardiac [myocytes](#). The alpha blockade decreases [peripheral vascular resistance](#), resulting in an antihypertensive effect. Beta inhibition leads to decreased [heart rate](#) (increased [PR interval](#)) and contractility. [Carvedilol](#) does not impact the [QT interval](#) or QRS duration.

C - Flecainide

Explanation Why

[Flecainide](#) is a [class IC](#) antiarrhythmic. While these drugs are powerful depressants of the sodium current, they have no effect on ventricular AP duration or the [QT interval](#).

D - Lidocaine

Explanation Why

[Lidocaine](#) is a [class IB antiarrhythmic](#). While all class I drugs slow conduction through the fast-inward sodium current (I_{Na}), class IB drugs selectively affect [ischemic](#) or depolarized tissue. They have little effect on atrial tissue and do not reduce the Phase 3 potassium current (I_K). This results in a shortened [action potential](#) duration but prolonged [effective refractory period](#). These drugs would not be expected to increase the QRS duration or the [QT interval](#).

E - Sotalol

Explanation Why

[Sotalol](#) is generally classified as a [class III antiarrhythmic drug](#) but also has [class II antiarrhythmic](#) properties (β -blocking effect). The hallmark of class III is prolongation of the [action potential](#) duration by blockade of potassium channels (I_K) that are responsible for [repolarization](#). This results in PR and [QT prolongation](#) but not an increase in QRS duration. Other class III agents, such as [amiodarone](#), do prolong both QRS and QT as a result of sodium, calcium, and potassium channel blockade as well as β [adrenoceptors](#) antagonism.

F - Quinidine

Explanation Why

[Quinidine](#) is a [class IA antiarrhythmic agent](#) used to treat or prevent [ventricular arrhythmias](#). All class I drugs slow conduction through the fast-inward sodium current (I_{Na}) in phase 0 and phase 4. This causes the phase 0 [depolarization](#) rate of the [cardiac action potential](#) to decrease. This occurs in a “use-dependent” fashion: at higher [heart rates](#), the sodium channel block intensifies, while the opposite is true at lower [heart rates](#). Class IA drugs also reduce phase 3 potassium current (I_K) and prolong the [effective refractory period](#). The overall effect of [quinidine](#), as seen on this patient's [ECG](#) a prolongation of the QT and QRS durations.

Question # 10

A 32-year-old man is brought to the emergency department 10 minutes after he sustained a stab wound to the left chest just below the clavicle. On arrival, he is hypotensive with rapid and shallow breathing and appears anxious and agitated. He is intubated and mechanically ventilated. Infusion of 0.9% saline is begun. Five minutes later, his pulse is 137/min and blood pressure is 84/47 mm Hg. Examination shows a 3-cm single stab wound to the left chest at the 4th intercostal space at the midclavicular line without active external bleeding. Cardiovascular examination shows muffled heart sounds and jugular venous distention. Breath sounds are normal bilaterally. Further evaluation of this patient is most likely to show which of the following findings?

	Answer	Image
A	A 15 mm Hg decrease in systolic blood pressure during inspiration	
B	Cough productive of frank blood	
C	Lateral shift of the trachea toward the right side	
D	Subcutaneous crepitus on palpation of the chest wall	
E	Inward collapse of part of the chest with inspiration	

Hint

Muffled heart sounds, hypotension, and jugular venous distention (also known as Beck's triad) indicate that this patient likely has cardiac tamponade.

Correct Answer

A - A 15 mm Hg decrease in systolic blood pressure during inspiration

Explanation Why

A > 10 mm Hg drop in [systolic](#) blood pressure during inspiration is known as [pulsus paradoxus](#), and is a sign observed in pathologies such as severe [asthma](#), [tension pneumothorax](#), [constrictive pericarditis](#), and moderate and severe [cardiac tamponade](#). In [cardiac tamponade](#), the outward expansion of the [right ventricle](#) is limited, especially as blood flows during inspiration. This effect causes the interventricular septum to bulge into the [left ventricle](#) and together with a reduction in [left ventricular](#) filling causes a decrease in [stroke volume](#) as well as a subsequent drop in the blood pressure.

B - Cough productive of frank blood

Explanation Why

[Hemoptysis](#) can occur in conditions such as [tuberculosis](#) and [pulmonary contusion](#). Although this patient exhibits some features consistent with [pulmonary contusion](#) ([tachycardia](#), rapid breathing), this condition is not associated with muffled [heart sounds](#) or [jugular venous distention](#). Also, the nature of his trauma makes this diagnosis unlikely.

C - Lateral shift of the trachea toward the right side

Explanation Why

[Tracheal deviation](#) toward the contralateral side of a lesion indicates a [tension pneumothorax](#). Although [tension pneumothorax](#) can also be caused by [penetrating chest trauma](#), and the patient has some features of this condition, such as [tachycardia](#), [hypotension](#), and [jugular venous distention](#), he does not have the key finding of reduced/absent breath sounds on the side of the lesion.

D - Subcutaneous crepitus on palpation of the chest wall

Explanation Why

[Subcutaneous emphysema](#) or [crepitus](#) on [palpation of the chest](#) wall suggests either [bronchial rupture](#), [esophageal rupture](#), or [pneumothorax](#), which also may be caused by [penetrating trauma](#). None of these conditions cause muffled [heart sounds](#).

E - Inward collapse of part of the chest with inspiration

Explanation Why

Paradoxical motion of part of the chest with breathing such as inward collapse with inspiration, suggests [flail chest](#), which is usually caused by severe blunt thoracic trauma rather than [penetrating chest trauma](#). Although this patient has shallow breathing, which can be a feature of [flail chest](#), the condition is not associated with muffled [heart sounds](#) or [jugular venous distention](#). Also, it is unlikely that a single stab wound would cause several [rib fractures](#).

Question # 11

A 14-year-old boy comes to the physician for a follow-up after a blood test showed a serum triglyceride level of 821 mg/dL. Several of his family members have familial hypertriglyceridemia. The patient is prescribed a drug that increases his risk of gallstone disease. The expected beneficial effect of this drug is most likely due to which of the following actions?

	Answer	Image
A	Increased lipoprotein lipase activity	
B	Decreased LDL-receptor degradation	
C	Decreased lipolysis in adipose tissue	
D	Increased PPAR-gamma activity	
E	Decreased HMG-CoA reductase activity	
F	Increased bile acid sequestration	

Hint

This patient with severe hypertriglyceridemia is at increased risk for acute pancreatitis. He requires a drug that causes a significant decrease in triglyceride levels.

Correct Answer

A - Increased lipoprotein lipase activity

Explanation Why

[Fibrates](#) (e.g., [bezafibrate](#), [fenofibrate](#), [gemfibrozil](#)) significantly lower [triglycerides](#) levels. [Fibrates](#) activate peroxisome proliferator-activated receptor alpha (PPAR- α), which increases [lipoprotein lipase](#) activity and thereby increases [triglyceride](#) clearance. Increased PPAR- α activity also increases hepatic [HDL](#) production. [Fibrates](#) are associated with an increased risk of [cholesterol gallstones](#) because they inhibit the cytochrome p450 enzyme [cholesterol 7- \$\alpha\$ hydroxylase](#) ($\uparrow$ [cholesterol](#) and $\downarrow$ [bile acid](#) concentration in [bile](#)). Since [gallstone](#) disease is itself a [risk factor](#) for [pancreatitis](#), patients are often switched from [fibrates](#) to [statins](#) after an adequate reduction in [triglyceride](#) levels has been achieved. Other side effects of [fibrates](#) include hepatotoxicity and [myopathy](#).

B - Decreased LDL-receptor degradation

Explanation Why

Decreased [LDL-receptor](#) degradation is caused by [PCSK9 inhibitors](#) (e.g., [alirocumab](#), [evolocumab](#)). These drugs can dramatically reduce serum [LDL](#) levels but they only minimally decrease [triglyceride](#) levels. Moreover, the typical side effects of this class of drugs include myalgias and neurocognitive defects (e.g., [delirium](#), [dementia](#)), but not [gallstone](#) formation.

C - Decreased lipolysis in adipose tissue

Explanation Why

Decreased [lipolysis](#) in [adipose tissue](#) is caused by [niacin](#), which inhibits [hormone-sensitive lipase](#). [Niacin](#) can decrease serum [LDL](#) and increase serum [HDL](#) levels but they only minimally decrease [triglyceride](#) levels. Moreover, typical side effects of [niacin](#) include flushing, [hyperuricemia](#), and [hyperglycemia](#), not [gallstone](#) formation.

D - Increased PPAR-gamma activity

Explanation Why

[Glitazones](#), which increase PPAR- γ activity, can cause a reduction in [triglyceride](#) levels. However, [glitazones](#) are primarily used as antidiabetic medications. Moreover, the typical side effects of [glitazones](#) are weight gain, [heart failure](#), and increased risk of [fractures](#), not [gallstone](#) formation. The drug that is most likely prescribed to this patient increases the activity of PPAR- α , not PPAR- γ .

E - Decreased HMG-CoA reductase activity

Explanation Why

HMG-CoA reductase is inhibited by [statins](#), which are often the first-line drugs for [dyslipidemia](#). However, [statins](#) primarily decrease [LDL](#) levels and only modestly decrease [triglyceride](#) levels. Moreover, the typical side effects of [statins](#) include hepatotoxicity and [myopathy](#). [Statin](#) therapy actually decreases the risk of [gallstone](#) disease.

F - Increased bile acid sequestration

Explanation Why

Increased [bile acid](#) sequestration is caused by [bile acid resin](#) (e.g., [cholestyramine](#)), which is used as an [LDL](#) lowering drug and can cause [gallstones](#) as a side effect. However, these drugs can slightly increase [triglyceride](#) levels, which is undesirable in this patient.

Question # 12

An 84-year old woman is brought to the emergency department from a nursing facility because of confusion and aggressive behavior. She has had multiple episodes of diarrhea the last 3 days. Her temperature is 37.5°C (99.5°F), blood pressure is 82/51 mm Hg, and pulse is 110/min. She is disoriented to place and time. Her extremities are warm and dry; her pulses are weak. Placement of a foley catheter yields cloudy urine with a foul odor. This patient is most likely to have which of the following hemodynamic profiles?

	Cardiac output	Pulmonary capillary wedge pressure	Systemic vascular resistance	Central venous pressure
A	Decreased	decreased	decreased	decreased
B	Increased	decreased	decreased	decreased
C	Decreased	increased	increased	increased
D	Increased	increased	increased	increased
E	Decreased	decreased	increased	decreased

	Answer	Image
A	A	
B	B	
C	C	
D	D	

	Answer	Image
E	E	

Hint

Infection should always be considered as a cause of altered mental status in an elderly patient.

Correct Answer

A - A

Explanation Why

This profile is suggestive of [neurogenic shock](#), which is an uncommon consequence of [CNS](#) trauma and is due to loss of [sympathetic](#) tone. It results in a form of [distributive shock](#) with [vasodilation](#) and reduced SVR, as well as relative [bradycardia](#) and decreased [cardiac output](#). Without history or evidence of significant head, neck, or [spine](#) trauma, this etiology is very unlikely to explain the patient's presentation.

B - B

Explanation But

The most important initial steps in management of [septic shock](#) are aggressive [fluid resuscitation](#) with [crystalloids](#) and the early administration of [empiric antibiotic therapy](#). Vasopressors should be added in patients who are refractory to intravenous fluid management.

Explanation Why

This profile is suggestive of [distributive shock](#), which is a collective term for [shock](#) that results in pathologic [vasodilation](#), leading to decreased SVR. The most common etiology is [septic shock](#) (in this case, a patient with likely [urinary tract infection](#)), but it can also be caused by [neurogenic shock](#) or [anaphylactic shock](#). Circulating [cytokines](#) and [bacterial toxins](#) cause pathologic [vasodilation](#) and lead to decreased SVR ([afterload](#)), as well as decreased CVP and [PCWP](#) ([preload](#)). Consequently, there is an increase in [heart rate](#) and [cardiac output](#). Patients with [septic shock](#) will often have warm, flushed [skin](#) as a result of this pathophysiology.

C - C

Explanation Why

This profile is suggestive of [cardiogenic shock](#), which results from any process that significantly decreases [cardiac output](#). It is generally characterized either as [obstructive shock](#) (due to [cardiac tamponade](#), massive [pulmonary embolism](#), [tension pneumothorax](#), etc) versus non-[obstructive shock](#) (due to acute [myocardial infarction](#), acute [heart failure](#), [arrhythmia](#), [valvular heart disease](#), etc). In response to the decrease in [cardiac output](#), there is increase in SVR. Poor forward flow leads to increases in [PCWP](#) and CVP. In [cardiogenic shock](#), patients will have cold, clammy [skin](#) on exam. This patient has no historical or examination elements to suggest a cardiogenic etiology of her presentation.

D - D

Explanation Why

This hemodynamic profile would be most consistent with an increase in overall [sympathetic](#) tone, such as secondary to [sympathomimetic drug](#) ingestion or illicit drug use ([cocaine](#), [amphetamines](#), etc). This patient's presentation and vital signs are not consistent with this answer, as she does not have evidence of [hypertension](#) but is hypotensive on exam.

E - E

Explanation Why

This profile is suggestive of [hypovolemic shock](#), which is [shock](#) related to loss of intravascular fluid. It is most often due to hemorrhage (often referred to as hemorrhagic shock). To maintain tissue perfusion, there is a compensatory [tachycardia](#) and increase in SVR. Despite these responses, there is an overall decrease in CVP, [PCWP](#), as well as reduced [stroke volume](#) and [cardiac output](#). On exam these patients will have cold and clammy [skin](#). The patient described here has no history suggestive of significant blood loss.

Question # 13

A 65-year-old woman comes to the physician because of a 3-month history of progressive shortness of breath and a dry cough. She has also noticed gradual development of facial discoloration. She has coronary artery disease, hypertension, and atrial fibrillation. She does not remember which medications she takes. Her temperature is 37°C (98.6°F), pulse is 90/min, respirations are 18/min, and blood pressure is 150/85 mm Hg. Pulse oximetry on room air shows an oxygen saturation of 95%. Examination shows blue-gray discoloration of the face and both hands. Diffuse inspiratory crackles are heard. An x-ray of the chest shows reticular opacities around the lung periphery and particularly around the lung bases. The most likely cause of this patient's findings is an adverse effect to which of the following medications?

	Answer	Image
A	Digoxin	
B	Procainamide	
C	Lisinopril	
D	Metoprolol	
E	Diltiazem	
F	Amiodarone	
G	Warfarin	

Hint

This woman's chest x-ray findings, along with her progressive shortness of breath, dry cough, and inspiratory crackles, are consistent with pulmonary fibrosis, an adverse effect of a medication commonly used for one of her pre-existing conditions.

Correct Answer

A - Digoxin

Explanation Why

[Digoxin](#) is a [cardiac glycoside](#) used in the treatment of A-fib and often has adverse effects due to its narrow therapeutic window. However, typical symptoms of [cardiac glycoside](#) toxicity are nausea, vomiting, blurry vision, changes in color vision (objects appear yellow), and [cardiac arrhythmias](#), not pulmonary [fibrosis](#) or blue-gray discoloration of the face.

B - Procainamide

Explanation Why

[Procainamide](#) is an antiarrhythmic that has many adverse effects, including gastrointestinal and [central nervous system](#) dysfunction, [thrombocytopenia](#), and [torsades de pointes](#) (due to a [prolonged QT interval](#)). It can also cause [drug-induced lupus erythematosus](#), which presents with [fever](#), polyarthritis, and phototoxicity. [Procainamide](#) would not cause this patient's pulmonary [fibrosis](#) or blue-gray discoloration of the face.

C - Lisinopril

Explanation Why

[Lisinopril](#) is an ACE inhibitor indicated for [hypertension](#). Although ACEIs can induce dry [cough](#), they would not cause this patient's pulmonary [fibrosis](#). Other common adverse effects include [angioedema](#) and [hyperkalemia](#).

D - Metoprolol

Explanation Why

[Metoprolol](#) has cardioprotective, antihypertensive, and antiarrhythmic properties and can cause pulmonary and cutaneous adverse effects. However, [beta blockers](#) typically cause bronchoconstriction, particularly in patients with [asthma](#) and reactive [airway](#) disease, not pulmonary [fibrosis](#). It can also cause flares of [psoriasis](#), which would not present with blue-gray discoloration of the face. Other adverse effects include [bradycardia](#), [hypotension](#), fatigue, and [erectile dysfunction](#).

E - Diltiazem

Explanation Why

[Diltiazem](#) is a [nondihydropyridine calcium channel blocker](#) that can be used in the treatment of [angina](#), [hypertension](#), or [atrial fibrillation](#). However, typical adverse effects include flushing, [constipation](#), [bradycardia](#), and worsening [cardiac output](#), not pulmonary [fibrosis](#) or blue-gray discoloration of the face.

F - Amiodarone

Explanation Why

[Amiodarone](#) is an [antiarrhythmic drug](#) that treats [atrial fibrillation](#), though it also has several adverse effects in many organ systems. Pulmonary toxicity, which can manifest as pulmonary [fibrosis](#), is responsible for most deaths associated with [amiodarone](#) therapy and correlates with total cumulative dose. Patients who develop pulmonary toxicity should stop [amiodarone](#) therapy and begin [glucocorticoid](#) therapy. A more than a two-fold elevation in [AST](#) and [ALT](#) is also a contraindication to continued therapy, although patients are usually asymptomatic. Blue-gray [skin](#) discoloration typically affects the face but is not dangerous.

G - Warfarin

Explanation Why

[Warfarin](#) is an [oral anticoagulant](#) that prevents [thromboembolic events](#), a common complication of A-fib. Its typical adverse effect is an increased risk of bleeding, which this patient has no signs of. Less commonly, it can cause [warfarin-induced skin necrosis](#), which presents with painful [purpura](#), hemorrhagic [blisters](#), and large areas of [necrosis](#), not with blue-gray discoloration of the face. [Warfarin](#) would also not cause this patient's pulmonary [fibrosis](#).

Question # 14

A 44-year-old woman comes to the office because of 4 episodes of loss of consciousness over the past 2 weeks. She recovered immediately and was not confused following the episodes. For the past 6 months, she has also had increased shortness of breath, palpitations, and chest tightness that resolves with rest. She immigrated with her family from India 10 years ago. Pulse is 115/min and irregular and blood pressure is 108/70 mm Hg. Cardiac examination shows an accentuated and split S₂. There is an opening snap followed by a low-pitched diastolic murmur in the fifth left intercostal space at the midclavicular line. An ECG shows atrial fibrillation and right axis deviation. Which of the following is the most likely underlying mechanism of these findings?

	Answer	Image
A	Increased pulmonary capillary wedge pressure	
B	Increased central venous pressure	
C	Increased oxygenated blood in the right ventricle	
D	Increased mean arterial pressure	
E	Increased left ventricular end diastolic pressure	


Hint

This patient has features of mitral stenosis (opening snap and diastolic murmur along with atrial fibrillation) that has progressed and caused further hemodynamic changes. MS commonly occurs as a late complication of rheumatic fever, which is more common in developing countries where patients may not have been adequately treated.

Correct Answer

A - Increased pulmonary capillary wedge pressure

Image



Explanation Why

[Mitral stenosis](#) leads to elevated left atrial pressure, which can trigger [atrial fibrillation](#) and cause increased [pulmonary capillary wedge pressure \(PCWP\)](#). Over time, the increased [PCWP](#) leads to remodeling of the [pulmonary arteries](#), increased [pulmonary vascular resistance](#), an S2 split, and [right ventricular hypertrophy](#) (seen as [right axis deviation](#)). In MS, shorter interval between the [opening snap](#) and S2 indicates more severe disease. This is because left atrial pressure is greater than left ventricular end-diastolic pressure.

B - Increased central venous pressure

Explanation Why

Central venous pressure ([preload](#)) is a marker of [venous return](#) to the [heart](#) and may be increased in states that expand blood volume (e.g., [congestive heart failure](#), intravenous fluid administration). While this patient shows signs of [right-sided heart failure](#), the presence of [mitral stenosis](#) with [right axis deviation](#) suggests a different underlying mechanism of the patient's disease.

C - Increased oxygenated blood in the right ventricle

Explanation Why

Increased oxygenated blood in the [right ventricle](#) may be present in patients with a [ventricular septal defect \(VSD\)](#) or an [atrial septal defect \(ASD\)](#) due to [left-to-right shunting](#). However, this patient's [cardiovascular examination](#) is consistent with [mitral stenosis](#) ([opening snap](#) with low-pitched [diastolic murmur](#) at the apex). She does not have a [murmur](#) that suggests a [VSD](#) or ASD ([holosystolic murmur](#)).

D - Increased mean arterial pressure

Explanation Why

Increased [mean arterial pressure](#) can be seen in patients with [essential hypertension](#) or secondary causes of [hypertension](#). This may eventually lead to [left ventricular hypertrophy](#) that can evolve into [diastolic heart failure](#). As this patient shows [right axis deviation](#) on [ECG](#), she is unlikely to have LV [hypertrophy](#). In addition, the patient has normal blood pressure on examination, which suggests a different diagnosis.


E - Increased left ventricular end diastolic pressure

Explanation Why

Increased left ventricular end diastolic pressure (LVEDP) is seen in [left heart failure](#), but this patient demonstrates signs of [mitral stenosis](#) (MS). MS causes a decrease in LVEDP due to impaired ability to fill the LV in [diastole](#) through the stenotic valve.

Question # 15

A previously healthy 2-month-old girl is brought to the emergency department because her lips turned blue while passing stools 30 minutes ago. She is at the 40th percentile for length and below the 35th percentile for weight. Pulse oximetry on room air shows an oxygen saturation of 65%, which increases to 76% on administration of 100% oxygen. Physical examination shows perioral cyanosis and retractions of the lower ribs with respiration. Cardiac examination shows a harsh grade 2/6 systolic crescendo-decrescendo murmur heard best at the left upper sternal border. Which of the following is most likely to improve this patient's symptoms?

	Answer	Image
A	Cooling of the face	
B	Elevation of the lower extremities	
C	Administration of indomethacin	
D	Hyperextension of the neck	
E	Knee to chest positioning	
F	Prone positioning	

Hint

Right ventricular outflow obstruction is a significant feature of this common congenital cardiac defect. Certain activities that slightly decrease oxygen saturation (e.g., defecating, crying) can acutely worsen the right ventricular outflow obstruction, triggering a cyanotic spell.

Correct Answer

A - Cooling of the face

Explanation Why

Cooling of the [infant](#)'s face with ice or cold water for 15–30 seconds can provoke the [diving reflex](#) to terminate [paroxysmal supraventricular tachycardias](#). [Supraventricular tachycardias](#) in [infants](#) typically present with nonspecific signs of [congestive heart failure](#) (e.g., poor feeding, sweating) and would not account for the patient's central [cyanosis](#), [hypoxemia](#), or ejection [murmur](#).

B - Elevation of the lower extremities

Explanation Why

Elevation of the lower extremities increases [venous return](#) to the [heart](#), which results in increased [preload](#) and [cardiac output](#). Lower limb elevation is a useful maneuver for patients suffering from [vasovagal syncope](#) and [hypovolemic shock](#). Although defecation can trigger [vasovagal syncope](#), these episodes would unlikely produce central (perioral) [cyanosis](#), as seen in this [infant](#). Increased [venous return](#) and the subsequent increase in right-to-left ventricular shunting of deoxygenated blood would worsen this patient's condition.

C - Administration of indomethacin

Explanation Why

Administration of [indomethacin](#) is used to treat [patent ductus arteriosus](#). [PDA](#) presents with a continuous machine-like [murmur](#), not a [systolic](#) crescendo-decrescendo [murmur](#), as heard in this patient. [Infants](#) with [PDA](#) are typically acyanotic at presentation, but eventually [Eisenmenger reaction](#) may result in shunt reversal and [differential cyanosis](#). [Indomethacin](#) will not improve this patient's condition.


D - Hyperextension of the neck

Explanation Why

Hyperextension of the neck can improve [tracheal](#) compression due to vascular rings. This anatomical anomaly can sometimes present as respiratory distress ([hypoxemia](#), lower [rib](#) retractions) and may be accompanied by [heart murmurs](#); up to 50% of affected [infants](#) also have an associated [heart](#) defect, as seen here. However, this [infant](#) lacks classic features of [tracheal](#) and/or esophageal compression from a [vascular ring anomaly](#), such as [stridor](#), [cough](#), difficulty feeding, or regurgitation, and hyperacute onset of symptoms. Hyperextension of the neck will not improve this patient's condition.

E - Knee to chest positioning

Image



Explanation But

More severe [tet spells](#) can be managed with IV [morphine](#) for sedation and fluids for volume

expansion. If those measures fail, IV beta blockers can be administered to reduce [sympathetic nervous system](#) activity and infundibular musculature spasms.

Explanation Why

[Infants](#) with [tetralogy of Fallot](#) frequently suffer from [tet spells](#), which are most likely due to [spastic](#) contractions of the [hypertrophic](#) infundibular musculature at the transition from the [right ventricle](#) to the [pulmonary artery](#). This contraction results in severe right outflow tract obstruction, causing increased right-to-left shunting of deoxygenated blood through the [VSD](#). Affected [infants](#) present with severe [hypoxemia](#) and profound [cyanosis](#). First-line treatments for [tet spells](#) include [knee](#)-chest positioning (older children can assume a squatting position) and administration of supplemental oxygen (acts as a pulmonary [vasodilator](#)). [Knee](#)-chest positioning increases arterial oxygenation via increased [systemic vascular resistance](#) and thus left-sided [heart](#) pressures. This, in turn, decreases right-to-left shunting across the [VSD](#) and promotes blood flow into the [pulmonary circulation](#) rather than the aorta.


F - Prone positioning

Explanation Why

[Prone positioning](#) can improve [airway](#) obstruction secondary to [laryngomalacia](#), which can cause lower [rib](#) retractions and [cyanosis](#) in rare cases. Most [infants](#) with [laryngomalacia](#) will instead have an [inspiratory stridor](#) and, sometimes, reflux and feeding difficulties, all of which this patient lacks. This [infant's](#) harsh [systolic murmur](#), central [cyanosis](#), and profound [hypoxemia](#) are not consistent with this diagnosis.

Question # 16

A 6-month-old girl is brought to the emergency department because her lips turned blue for several minutes earlier during feeding. Her father reports that the patient had similar episodes while playing that resolved quickly. During the examination, she sits calmly in her father's lap. When her ears are examined, the patient begins to cry and her lips and fingers begin to turn blue. Further evaluation of this patient is most likely to show which of the following?

	Answer	Image
A	Anomalous pulmonary venous return on MR angiography	
B	Machine-like hum on auscultation	
C	Diminutive left ventricle on echocardiogram	
D	Decreased blood pressure in both lower extremities	
E	Pulmonary vascular congestion on chest x-ray	
F	Right axis deviation on ECG	

Hint

The cyanotic episode described here is characteristic of the tet spells seen in tetralogy of Fallot (TOF).

Correct Answer

A - Anomalous pulmonary venous return on MR angiography

Explanation Why

Anomalous pulmonary [venous return](#) on MR angiography is diagnostic of [total anomalous pulmonary venous return](#) (TAPVR). Although [TAPVR](#) is survivable and can cause central [cyanosis](#) in patients with a [right-to-left shunt](#), symptoms are continuous rather than episodic, as seen here.

B - Machine-like hum on auscultation

Explanation Why

A machine-like [murmur](#) over the [heart](#) is pathognomonic for [patent ductus arteriosus](#) ([PDA](#)). [PDA](#) is usually an acyanotic congenital condition because most blood passes through normal circulation and thus sufficient [oxygen saturation](#) is maintained; only a small amount of blood is pumped through the [PDA](#). Furthermore, [PDA](#) causes continuous symptoms, as opposed to the episodic symptoms described for this patient.

C - Diminutive left ventricle on echocardiogram

Explanation Why

A diminutive [left ventricle](#) on [echocardiogram](#) is seen in [hypoplastic left heart syndrome](#), which can also present with central [cyanosis](#). Because the left [heart](#) is unable to adequately perfuse the peripheral tissues, the associated [cyanosis](#) would be constant, rather than episodic.

D - Decreased blood pressure in both lower extremities

Explanation Why

Decreased blood pressure in the lower extremities is characteristic of [aortic coarctation](#), which presents with [differential cyanosis](#) (i.e., of the lower extremities) instead of the central [cyanosis](#) (i.e., blue [lips](#) and fingers) seen in this patient. Moreover, [aortic coarctation](#) rarely causes [hypoxemia](#); in the rare case of [hypoxemia](#), it is usually mild. Finally, [aortic coarctation](#) results in chronic rather than episodic symptoms.


E - Pulmonary vascular congestion on chest x-ray

Explanation Why

Pulmonary vascular congestion seen on chest [x-ray](#) is a sign of [left heart failure](#), with subsequent “backing up” of blood into the pulmonary vasculature, leading to enlarged pulmonary vessels. Although pulmonary vascular congestion is classically associated with [congestive heart failure](#) in adults (especially elderly patients), some congenital cardiac defects can also cause this finding (e.g., [truncus arteriosus](#), [total anomalous pulmonary venous return](#)). In TOF, there is a right ventricular outflow obstruction, leading to decreased blood flow to the pulmonary vasculature and, consequently, decreased pulmonary vascular markings on chest [x-ray](#).

F - Right axis deviation on ECG

Image




Explanation Why

[Right axis deviation](#) is typically seen on [ECG](#) in cases of TOF due to [right ventricular hypertrophy](#). The thickened [myocardium](#) in the [right ventricle](#) produces increased electrical activity, which shifts the axis of cardiac [depolarization](#). TOF also involves an overriding aorta, [ventricular septal defect](#), and [pulmonary stenosis](#). It is the most common congenital [cyanotic heart](#) lesion. In addition to these [ECG](#) findings, a chest [x-ray](#) showing a boot-shaped [heart](#) also supports the diagnosis. [Echocardiography](#) is used to confirm TOF.

Question # 17

A 44-year-old woman comes to the physician because of progressively worsening shortness of breath with exertion and intermittent palpitations over the last 2 months. She has had neither chest pain nor a cough. Her pulse is 124/min and irregular. Physical examination shows a grade 4/6 high-pitched holosystolic murmur that is best heard at the apex and radiates to the back. The murmur increases in intensity when she clenches her hands into fists. The lungs are clear to auscultation. Further evaluation of this patient is most likely to show which of the following findings?

	Answer	Image
A	Pulmonary artery thrombus on computed tomography scan	
B	Diffuse ST elevations on electrocardiogram	
C	Obstruction of the right marginal artery on coronary angiogram	
D	Reversible area of myocardial ischemia on nuclear stress test	
E	Dilation of left atrium on echocardiogram	

Hint

A high-pitched, holosystolic murmur at the apex with radiation to the back and higher intensity with increased systemic vascular resistance (hand grip, squatting) in a patient with exertional dyspnea is characteristic of mitral regurgitation.

Correct Answer

A - Pulmonary artery thrombus on computed tomography scan

Explanation Why

A [pulmonary artery thrombus](#) on CT is seen in [pulmonary embolism \(PE\)](#). While this condition can cause [dyspnea](#), it typically manifests with an acute onset of [cough](#), [chest pain](#), [hypotension](#), and [jugular venous distention](#). This patient's 2-month history of progressive [shortness of breath](#) and absence of [chest pain](#) and [cough](#) make this diagnosis unlikely.

B - Diffuse ST elevations on electrocardiogram

Explanation Why

Diffuse [ST elevations](#) on [ECG](#) suggest [acute pericarditis](#), which can cause [dyspnea](#), as seen in this patient. However, this condition manifests with an acute onset of severe [chest pain](#) that is relieved by sitting up and leaning forward, and a [pericardial friction rub](#) on [cardiac auscultation](#), neither of which are present.

C - Obstruction of the right marginal artery on coronary angiogram

Explanation Why

Obstruction of the [right marginal artery](#) on a coronary angiogram is consistent with [acute coronary syndrome \(ACS\)](#), which typically presents with acute retrosternal [chest pain](#)/pressure, radiation to the [jaw](#) and/or left arm, [shortness of breath](#), and nausea/vomiting. It also produces symptoms of [right-sided heart failure](#) due to reduced blood supply of the [right ventricle](#), such as [jugular venous distention](#). However, these symptoms are not present here, and this patient's progressive 2-month history of increasing [dyspnea](#) makes this diagnosis unlikely. Also, a significant [systolic murmur](#) characteristic of [mitral regurgitation](#) requires a different diagnostic method than an angiogram.


D - Reversible area of myocardial ischemia on nuclear stress test

Explanation Why

Reversible [myocardial ischemia](#) due to CAD can be diagnosed by a nuclear stress test that includes myocardial perfusion imaging (MPI). While this condition also causes [shortness of breath](#), it typically presents with an acute onset of retrosternal [pain](#)/pressure that may radiate to the [jaw](#) and/or left arm, and autonomic symptoms such as sweating, pallor, and nausea/vomiting, which are not present here.

E - Dilation of left atrium on echocardiogram

Image



Explanation Why

Dilatation of the [left atrium](#) (LA) results from volume and pressure overload due to chronic [mitral regurgitation](#) (MR) in this patient. LA enlargement can cause [atrial fibrillation](#), which explains the

[tachycardia](#) and [irregular heartbeat](#) seen here. [Echocardiography](#) would be the best diagnostic method for visualizing the suspected structural and flow abnormalities of [MR](#).

Question # 18

A 55-year-old man is brought to the emergency department 3 hours after ingesting approximately 30 tablets of an unknown drug in an apparent suicide attempt. His temperature is 36.5°C (97.7°F), pulse is 40/min, respiratory rate is 19/min, and blood pressure is 85/50 mm Hg. Examination shows cold, clammy extremities. Scattered expiratory wheezing is heard throughout both lung fields. His fingerstick blood glucose concentration is 62 mg/dL. ECG shows prolonged PR intervals and narrow QRS complexes. Intravenous fluid resuscitation and atropine do not improve his symptoms. Administration of which of the following is the most appropriate next step in treatment of this patient?

	Answer	Image
A	Sodium bicarbonate	
B	Pralidoxime	
C	Glucagon	
D	Activated charcoal	
E	Digoxin-specific antibody	
F	Naloxone	

Hint

Bradycardia, hypotension, wheezing, hypoglycemia, and prolonged PR intervals suggest severe beta-blocker intoxication.

Correct Answer

A - Sodium bicarbonate

Explanation But

Sodium bicarbonate is also an [antidote](#) to [barbiturates](#), [salicylates](#), [tricyclic antidepressants](#), and a number of other drugs. It works by stabilizing the cardiac membrane (mostly in widened QRS) and also by alkalizing the [urine](#) and enhancing the excretion of these substances.

Explanation Why

Although sodium bicarbonate is used to treat [propranolol](#) overdose, it is not effective in the treatment of other types of [beta-blocker](#) toxicity. In addition to the typical manifestations of [beta-blocker](#) toxicity, [propranolol](#) causes QRS widening due to sodium channel blockade. This patient's narrow [QRS complexes](#) are unlikely caused by [propranolol](#).

B - Pralidoxime

Explanation Why

[Pralidoxime](#) is used in the treatment of [cholinergic syndrome](#). Although this syndrome can cause [bradycardia](#) and wheezing (due to bronchoconstriction), other typical manifestations such as sweating, lacrimation, or urination ([SLUDGE-M](#)) would also be expected in this patient. Furthermore, [hypoglycemia](#) is not a common manifestation of [cholinergic syndrome](#).

C - Glucagon

Explanation Why

In patients with [beta-blocker](#) overdose, intravenous [glucagon](#) should be administered when blood pressure and [heart rate](#) do not respond sufficiently to [IV fluids](#) and [atropine](#) alone. [Glucagon](#) increases intracellular [cAMP](#) by activating [adenylate cyclase](#), which improves cardiac contractility

and [heart rate](#) while bypassing the [beta receptors](#). It also improves [hypoglycemia](#).

D - Activated charcoal

Explanation Why

[Activated charcoal](#) may be used in acute [poisoning](#) to bind medications in the [gastrointestinal tract](#) and prevent their absorption. However, it is only effective within 1–2 hours (maximum) following ingestion of the poisonous substance. [Activated charcoal](#) would not be useful for this patient who presented 3 hours after ingestion of the tablets.

E - Digoxin-specific antibody

Explanation Why

[Digoxin](#)-specific [antibody](#) fragments are the treatment of choice for [digitalis poisoning](#), which typically presents with nausea, vomiting, abdominal [pain](#), [diarrhea](#), blurry vision with a yellow tint, and [cardiac arrhythmias](#). While [digitalis](#) toxicity can also present with [bradycardia](#) and [hypotension](#), wheezing (indicating bronchoconstriction) and [hypoglycemia](#) are not consistent with this condition.







F - Naloxone







Explanation Why

[Naloxone](#) is the first-line treatment for [opioid overdose](#), which can present with mild [bradycardia](#) and [hypotension](#). However, intoxication with [opioids](#) usually also presents with decreased [respiratory rate](#) and [miosis](#), which are not seen in this patient. Wheezing and prolongation of the [PR interval](#) are not consistent with [opioid overdose](#).

Question # 19

A 2300-g (5.07-lb) male newborn is delivered at term to a 39-year-old woman. Examination shows a sloping forehead, a flat nasal bridge, increased interocular distance, low-set ears, a protruding tongue, a single palmar crease and an increased gap between the first and second toe. There are small white and brown spots in the periphery of both irises. The abdomen is distended. An x-ray of the abdomen shows two large air-filled spaces in the upper quadrant. This patient's condition is most likely associated with which of the following cardiac anomalies?

	Answer	Image
A	Patent ductus arteriosus	1  2  3 
B	Atrial septal defects	1  2  3 

	Answer	Image
C	Atrioventricular septal defect	Down syndrome Epidemiology Incidence: ~ 1:700 live births Etiology Three complete copies of chromosome 21; due to meiotic nondisjunction in 95% of cases Karyotype ♀ : 47,XX,+21 ♂ : 47,XY,+21 Complications Due to organ malformations and immunodeficiency. Increased risk of AML/ALL. Early onset Alzheimer's disease Important Risk increases with maternal age Life expectancy ~50 years Karyotype   Short stature Intellectual disability Hearing loss, recurrent otitis media Brushfield spots Hypothyroidism Cardiac malformations (e.g., ASD, VSD) Gastrointestinal malformations (e.g., duodenal atresia/stenosis, annular pancreas) Urogenital malformations (e.g., hypospadias; low fertility in men) Obesity Single transverse palmar crease (I), clinodactyly (II), brachydactyly (III), camptodactyly (IV) Umbilical and inguinal hernias Delayed motor development, hypotonia Sandal gap
D	Tetralogy of Fallot	 1  2
E	Ventricular septal defect	 1  2  3


Hint

The characteristic morphological features, Brushfield spots in the eye, and the “double bubble” sign on abdominal x-ray (indicating duodenal atresia) are all suggestive of trisomy 21 (Down syndrome).

Correct Answer

A - Patent ductus arteriosus

Image




Explanation Why

A [patent ductus arteriosus](#) is most commonly associated with prematurity, [fetal alcohol syndrome](#), [birth](#) at high altitude, or [congenital rubella infection](#). Although it occurs in approx. 5% of patients with [Down syndrome](#), it is not the most common congenital cardiac anomaly seen in this condition.


B - Atrial septal defects

Image


1



2



3




Explanation Why

[Atrial septal defects](#) (ASDs) are among the most common congenital [heart](#) defects and are commonly associated with [fetal alcohol syndrome](#) and [trisomy 21](#). Although ASD occurs in approx. 15% of patients with [Down syndrome](#), it is not the most common congenital cardiac anomaly seen in this condition.

C - Atrioventricular septal defect

Image




Explanation Why

[Atrioventricular septal defects](#) (AVSD; also referred to as [endocardial cushion defects](#)) are the most common congenital [heart](#) defects in patients with [Down syndrome](#), occurring in about 40% of [newborns](#) with [trisomy 21](#). AVSDs are usually diagnosed in the prenatal period, and discovery of an [AVSD](#) should warrant prompt genetic testing, as there is a strong association with [trisomy 21](#). [Down syndrome](#) is diagnosed in approx. 40–50% of children with [AVSD](#).

D - Tetralogy of Fallot

Image




Explanation Why

[Tetralogy of Fallot](#) (TOF) is the most common cause of [cyanotic heart disease](#) in children. Most cases arise sporadically, but approx. 15% of patients with TOF have an underlying genetic syndrome (such as [Down syndrome](#) or [DiGeorge syndrome](#)). Although it occurs in approx. 5% of patients with [Down syndrome](#), it is not the most common congenital cardiac anomaly seen in this condition.

E - Ventricular septal defect

Image




Explanation Why

[Ventricular septal defects \(VSDs\)](#) account for approx. 50% of all congenital [heart](#) defects, making them the most common cause of congenital [heart](#) disease overall. They are commonly associated with [fetal alcohol syndrome](#), intrauterine infections, and many genetic syndromes (including [trisomy 13](#), [trisomy 18](#), and [trisomy 21](#)). Although [VSD](#) occurs in approx. 30% of patients with [Down syndrome](#), it is not the most common congenital cardiac anomaly seen in this condition.

Question # 20

A 72-year-old woman with hypertension comes to the physician because of swelling and pain in both legs for the past year. The symptoms are worse at night and improve in the morning. Current medications include losartan and metoprolol. Her temperature is 36°C (96.8°F), pulse is 67/min, and blood pressure is 142/88 mm Hg. Examination shows normal heart sounds; there is no jugular venous distention. Her abdomen is soft and the liver edge is not palpable. Examination of the lower extremities shows bilateral pitting edema and prominent superficial veins. The skin is warm and there is reddish-brown discoloration of both ankles. Laboratory studies show a normal serum creatinine and normal urinalysis. Which of the following is the most likely underlying cause of this patient's symptoms?

	Answer	Image
A	Increased platelet aggregation	
B	Decreased lymphatic flow	
C	Decreased intravascular oncotic pressure	
D	Decreased arteriolar resistance	
E	Increased venous valve reflux	

	Answer	Image
F	Increased capillary permeability	

Hint

This patient most likely has varicose veins, a condition that can be problematic for patients with occupations that require prolonged standing.

Correct Answer

A - Increased platelet aggregation

Explanation Why

Increased [platelet aggregation](#) is involved in the pathophysiology of [deep venous thrombosis](#), which also classically manifests with lower extremity swelling and [pain](#). However, symptoms of [DVT](#) are typically acute in onset and unilateral; the bilateral, chronic nature of this patient's symptoms makes [DVT](#) unlikely.

B - Decreased lymphatic flow

Explanation Why

Decreased lymphatic flow can result in [lymphedema](#), a type of [nonpitting edema](#) that results due to the accumulation of filtered [lymph](#) fluid in the [interstitium](#). This patient's [pitting edema](#), chronic [skin](#) changes, and lack of [risk factors](#) for [lymphedema](#) ([radiation therapy](#), surgery, [tumor](#) obstruction) suggest a different underlying mechanism.

C - Decreased intravascular oncotic pressure

Explanation Why

Decreased plasma [oncotic pressure](#) is typically seen in diseases that cause hypoproteinemia, such as [nephrotic syndrome](#), [malnutrition](#), and/or [liver cirrhosis](#). While low [oncotic pressure](#) can lead to bilateral peripheral [edema](#), this patient does not have any [risk factors](#) or exam findings (e.g., [hepatomegaly](#), [proteinuria](#)) consistent with renal or [liver](#) disease.


D - Decreased arteriolar resistance

Explanation Why

Decreased arteriolar resistance can be caused by medications that induce vascular [smooth muscle](#) relaxation, such as [dihydropyridine calcium channel blockers](#) (e.g., [amlodipine](#)). Patients on [dihydropyridines](#) can also present with [pitting edema](#). However, this patient is not taking [dihydropyridines](#), and neither [losartan](#) nor [metoprolol](#) cause decreased arteriolar resistance.

E - Increased venous valve reflux

Image



Explanation Why

A female patient of advanced age with bilateral leg swelling, [skin](#) discoloration, and [pain](#) most likely has [chronic venous insufficiency](#) and [varicose veins](#). In [CVI](#), defective venous valve leaflets result in reflux of venous blood back into the periphery, leading to increased venous pressure ([hydrostatic](#)

[pressure](#)) and subsequent extravasation of venous blood into the [interstitium](#). [Plasma proteins](#) in the [interstitium](#) in turn cause [edema](#) and tissue hypoperfusion, which results in [inflammation](#), [atrophy](#), and possibly ulcer formation.

F - Increased capillary permeability

Explanation Why

Increased [capillary](#) permeability is seen in patients with systemic [inflammation](#) and/or [distributive shock](#) (e.g., from [anaphylaxis](#) or [sepsis](#)). Extravasation of [plasma proteins](#) through the permeable [capillary](#) walls causes [edema](#). This patient is afebrile, not [tachycardic](#) or hypotensive, and has no symptoms to suggest infection or [inflammation](#). Her chronic [skin](#) changes suggest a different underlying cause of her symptoms.

Question # 21

A 75-year-old man is brought to the emergency room because of dizziness and palpitations for the past hour. His pulse is 185/min and blood pressure is 100/52 mm Hg. An ECG shows a narrow-complex, regular tachycardia without P waves. A drug is administered that activates cardiac potassium channels and causes transient atrioventricular block. Which of the following adverse effects is most likely to occur?

	Answer	Image
A	Severe headache	
B	Sedation	
C	Chest pain	
D	Dry cough	
E	Perioral paresthesia	
F	Yellow vision	

Hint

The patient has been given the antiarrhythmic drug adenosine.

Correct Answer

A - Severe headache

Explanation Why

Severe [headache](#) is a common side effect of [nitrates](#) (e.g., [nitroglycerin](#), [isosorbide dinitrate](#)), which release [nitric oxide](#) in [smooth muscle cells](#), resulting in [vasodilation](#) and [increased intracranial pressure](#). Other side effects of [nitrates](#) include [hypotension](#), [reflex tachycardia](#), and [syncope](#). Severe [headaches](#) are not a common side effect of adenosine, which this patient received.

B - Sedation

Explanation Why

Sedation is a known side effect of [beta blockers](#) (e.g., [metoprolol](#), [labetalol](#)) and [class IB antiarrhythmics](#) (e.g., [lidocaine](#), [mexiletine](#)). These agents are [lipophilic](#) and can penetrate the [blood-brain barrier](#), resulting in [CNS](#) side effects. Adenosine does not cause sedation.

C - Chest pain

Explanation But

[Theophylline](#) and [caffeine](#) are adenosine receptor antagonists that can inhibit the effect of adenosine.

Explanation Why

[Chest pain](#) is a common adverse effect in patients treated with adenosine. Adenosine causes a transient atrioventricular (AV) block by increasing potassium efflux from cells, which causes [hyperpolarization](#), decreased calcium influx, and decreased levels of intracellular [cAMP](#). The brief [AV block](#) (usually < 15 secs) can cause [chest pain](#), and a feeling of impending doom. Other side effects of adenosine include [hypotension](#), flushing, bronchospasm, [bradycardia](#), and nausea. Prolonged [asystole](#) is rare but also a potential complication.

D - Dry cough

Explanation Why

Dry [cough](#) is a side effect of ACE-inhibitors that results from decreased breakdown of [bradykinin](#). This increases production of [arachidonic acid](#) metabolites (e.g., [prostaglandins](#)), which accumulate in the upper respiratory tract and the [lungs](#), resulting in a dry [cough](#). Adenosine can cause [shortness of breath](#) but not dry [cough](#).

E - Perioral paresthesia

Explanation Why

Perioral [paresthesia](#) is a side effect of [lidocaine](#), a [class IB antiarrhythmic](#) agent used to treat [ventricular tachycardia](#). When administered intravenously, [lidocaine](#) works by blocking the sodium channels in cardiac cells. Adenosine, which this patient received for his [supraventricular arrhythmia](#), does not cause perioral [paresthesia](#).

F - Yellow vision

Explanation Why

Blurry vision with a yellow tint and halos is a side effect of [digoxin](#). [Digoxin](#) inhibits [Na⁺/K⁺-ATPase](#) and indirectly increases calcium in the cell, leading to increased cardiac contractility. Other side effects of [digoxin](#) include [arrhythmias](#), [kidney failure](#), [hyperkalemia](#), and [hypercalcemia](#). Yellow vision is not a known side effect of adenosine.

Question # 22

A 55-year-old man comes to the physician because of a 2-day history of severe perianal pain and bright red blood in his stool. Examination shows a bulging, red nodule at the rim of the anal opening. Which of the following arteries is the most likely source of blood to the mass found during examination?

	Answer	Image
A	Deep circumflex iliac	Deep circumflex iliac artery Superficial circumflex iliac artery Ascending branch of lateral circumflex femoral artery Lateral circumflex femoral artery Descending branch of lateral circumflex femoral artery Inferior epigastric artery Superficial epigastric artery External pudendal arteries Femoral artery Profunda femoris artery Medial circumflex femoral artery
B	Internal pudendal	Lymphatic channel to inferior mesenteric lymph nodes Inferior hypogastric plexus Lymphatic channel to internal iliac lymph nodes Pectinate line Lymphatic channel to internal iliac lymph nodes Inferior rectal nerve Lymphatic channel to superficial inguinal lymph nodes Superior rectal vein Superior rectal artery Middle rectal vein Middle rectal artery Corpus cavernosum recti Inferior rectal vein Inferior rectal artery
C	Median sacral	

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

	Answer	Image
D	Inferior gluteal	
E	Superior rectal	


Hint

Bright red blood in stool with a painful nodule at the rim of the anal opening suggests an external hemorrhoid.

Correct Answer

A - Deep circumflex iliac

Image




Explanation Why

The deep circumflex iliac artery is a branch of the external iliac artery that supplies the [iliacus](#), iliac fossa, and deep abdominal wall of the inguinal region. This [artery](#) does not supply the [anal canal](#).

B - Internal pudendal

Image



Explanation Why

The [internal pudendal artery](#) gives rise to the [inferior rectal artery](#), which supplies the part of the [anal canal](#) below the [pectinate line](#), where [external hemorrhoids](#) develop. An [external hemorrhoid](#) is often painful because the region [below the pectinate line](#) arises from [ectoderm](#) and is innervated by the inferior rectal nerve, which transmits somatic sensations such as [pain](#). [Internal hemorrhoids](#), which arise [above the pectinate line](#), are supplied by the [superior rectal artery](#), a branch of the [inferior mesenteric artery](#)). Unlike [external hemorrhoids](#), [internal hemorrhoids](#) are typically painless because the region of the [anal canal above the pectinate line](#) arises from [endoderm](#) and is innervated by the [inferior hypogastric plexus](#), which carries visceral sensation only.


C - Median sacral

Explanation Why

The [median sacral artery](#) is a [branch of the abdominal aorta](#) that supplies part of the lumbar [vertebral column](#), the [coccyx](#), and the [sacrum](#). This [artery](#) does not supply the [anal canal](#).

D - Inferior gluteal

Image




Explanation Why

The [inferior gluteal artery](#) is a branch of the internal iliac artery that supplies the [gluteus maximus](#), [quadratus femoris](#), and [piriformis](#) muscles. Although [external hemorrhoids](#) are also supplied by a branch of the [internal iliac artery](#), the [inferior gluteal artery](#) does not supply the [anal canal](#).

E - Superior rectal

Image




Explanation Why

The [superior rectal artery](#) supplies the [anal canal](#) above the [pectinate line](#) (where [internal hemorrhoids](#) develop). In contrast to [external hemorrhoids](#), [internal hemorrhoids](#) are typically painless.

Question # 23

A 71-year-old man with hypertension is taken to the emergency department after the sudden onset of stabbing abdominal pain that radiates to the back. He has smoked 1 pack of cigarettes daily for 20 years. His pulse is 120/min and thready, respirations are 18/min, and blood pressure is 82/54 mm Hg. Physical examination shows a periumbilical, pulsatile mass and abdominal bruit. There is epigastric tenderness. Which of the following is the most likely underlying mechanism of this patient's current condition?

	Answer	Image
A	Coronary artery narrowing	
B	Aortic wall stress	
C	Mesenteric atherosclerosis	
D	Gastric mucosal ulceration	
E	Abdominal wall defect	
F	Portal vein stasis	

Hint

Tearing back pain, a pulsatile abdominal mass, and sudden hypotension in a male patient with a history of smoking is concerning for pathology located in the retroperitoneal space.

Correct Answer


A - Coronary artery narrowing

Explanation Why

[Coronary atherosclerosis](#) resulting narrowing of the vessel lumen is involved in the pathophysiology of [angina pectoris](#) and [myocardial infarction](#). [Coronary atherosclerosis](#) leads to narrowing of the arterial lumen, which decreases the blood flow to the [myocardium](#). During exercise, [myocardial](#) flow is unable to meet increased [myocardial](#) demand, which causes [myocardial ischemia](#) and symptoms of [angina](#). While smoking increases the risk for [MI](#), this man's history of acute onset, stabbing abdominal [pain](#) with an umbilical mass and [hypotension](#) suggest a different disease process.

B - Aortic wall stress

Image



Explanation Why

This patient most likely has a ruptured [abdominal aortic aneurysm](#). The pathophysiology of abdominal [aortic aneurysms](#) is not fully understood, but numerous genetic and environmental factors, such as smoking, are believed to cause [inflammation](#) and proteolysis of [connective tissue](#) within the aortic wall. This leads to loss of structural integrity and weakening of the aorta. When high aortic wall stresses are applied, as in [hypertension](#), further weakening, widening, and rupture of the aortic wall can occur. Abdominal [aortic aneurysms](#) are most commonly located below the [renal arteries](#) (infrarenal).

C - Mesenteric atherosclerosis

Explanation Why

[Mesenteric atherosclerosis](#) is involved in the pathophysiology of [chronic mesenteric ischemia](#), in which intestinal blood flow is reduced. In the postprandial period, there is a mismatch between low splanchnic blood flow and high intestinal metabolic demand, which leads to postprandial [ischemia](#) and [pain](#) within an hour of eating. Bloating, nausea, and weight-loss due to food aversion are common symptoms. This man has acute-onset, stabbing abdominal [pain](#) with an umbilical mass and [hypotension](#), features that suggest a different, more emergent disease process.

D - Gastric mucosal ulceration

Explanation Why

Gastric mucosal ulceration is involved in the pathophysiology of [peptic ulcer disease](#). One of the complications of this disease is gastric or duodenal perforation, which presents with sudden onset, stabbing abdominal [pain](#), and distention that can progress to peritonitis. This man's periumbilical mass, abdominal [bruit](#), and [hypotension](#) are inconsistent with a perforated [duodenal](#) or [gastric ulcer](#).

E - Abdominal wall defect

Explanation Why

Abdominal wall defects can predispose patients to hernia formation. Uncomplicated hernias are typically asymptomatic and occur when intra-abdominal contents protrude through congenital defects or areas of weakness in the abdominal wall. Complicated hernias include [incarcerated hernias](#) (irreducible), [strangulated hernias](#) (disrupted blood supply), or obstructed hernias (kinked intestine), and typically present with acute [pain](#) at the site of the hernia and possible [bowel obstruction](#). Although this patient has a periumbilical mass, he also has an abdominal [bruit](#) and [hypotension](#), suggesting a different etiology for his symptoms.

F - Portal vein stasis

Explanation Why

[Portal vein](#) stasis predisposes patients to the formation of acute [portal vein thrombosis](#), which presents with [right upper quadrant](#) or generalized abdominal [pain](#). [Portal vein thrombosis](#) often arises as a complication of [cirrhosis](#), but this patient lacks symptoms of chronic [liver](#) disease ([jaundice](#), [pruritus](#), [hepatosplenomegaly](#), [telangiectasias](#), [caput medusae](#), [gynecomastia](#)). Furthermore, this man's periumbilical mass and abdominal [bruit](#) suggest a different cause for his symptoms.

Question # 24

A 43-year-old man is brought to the emergency department by his wife after a near-syncopal episode. He was doing yard work when he began feeling dizzy and had to lie down. Earlier in the day, he was started on lisinopril. On arrival, he is fully oriented. His pulse is 100/min and blood pressure is 92/60 mm Hg. Serum electrolytes are within normal limits. An ECG shows no evidence of ischemia. Concurrent treatment with which of the following agents most likely predisposed the patient to this episode?

	Answer	Image
A	Atorvastatin	
B	Ibuprofen	
C	Hydrochlorothiazide	
D	Lithium	
E	Trimethoprim/sulfamethoxazole	
F	Metformin	

Hint

The patient presents with a symptomatic hypotensive episode after initiating lisinopril (first-dose hypotension), an angiotensin-converting enzyme inhibitor (ACE inhibitor). ACE inhibitors work synergistically with other antihypertensive agents.

Correct Answer

A - Atorvastatin

Explanation Why

[Atorvastatin](#) is a type of [statin](#), or [HMG-CoA reductase inhibitor](#), used to treat [hypercholesterolemia](#). It inhibits [cholesterol synthesis](#) in the [liver](#). Although [atorvastatin](#) would have interactions with other lipid-lowering agents and [CYP3A4](#) inhibitors, it has no significant interactions with [antihypertensives](#) like [lisinopril](#).

B - Ibuprofen

Explanation Why

[Ibuprofen](#) is an [NSAID](#), which has anti-inflammatory effects by reversibly inhibiting the enzymes [COX-1](#) and [COX-2](#). Because it would actually decrease an ACE inhibitor's kinin-mediated synthesis of [prostaglandins](#), [ibuprofen](#) would decrease [lisinopril](#)'s hypotensive effect rather than enhance it as in this patient.

C - Hydrochlorothiazide

Explanation Why

[Hydrochlorothiazide](#) is a [thiazide diuretic](#) which inhibits the NCC transporter of the [distal convoluted tubule](#). It is a first-line agent for treating [hypertension](#). Patients who are taking [diuretics](#) or are volume depleted are at higher risk for first-dose [hypotension](#) when additional [antihypertensives](#) such as ACE inhibitors are initiated, as in this patient.

D - Lithium

Explanation Why

[Lithium](#) is used to treat [bipolar disorders](#) and has numerous adverse effects and interactions with other drugs. ACE inhibitors can lead to increased [lithium](#) reabsorption in the [proximal](#) tubules and increased risk of [lithium toxicity](#), which would present with [tremor](#), [ataxia](#), vomiting, [diarrhea](#), and other neuropsychiatric impairments but not [hypotension](#) as this patient is experiencing.

E - Trimethoprim/sulfamethoxazole

Explanation Why

[TMP/SMX](#) is a combination [antibiotic](#) that has a broad spectrum of activity against many gram-positive and gram-negative bacteria. It may interact with ACE inhibitors like [lisinopril](#) by synergistically increasing the risk of [hyperkalemia](#). However, it has no direct effect on blood pressure and would not lead to [hypotension](#) as in this patient.

F - Metformin

Explanation Why

[Metformin](#) is a [biguanide](#) used as a first-line treatment for [type 2 diabetes](#) to increase [insulin](#) sensitivity and decrease [gluconeogenesis](#). It has interactions with [sulfonylureas](#) and is contraindicated before the use of contrast medium because of the increased risk of [lactic acidosis](#). However, [metformin](#) has no known interaction with ACE inhibitors.

Question # 25

A 56-year-old man comes to the emergency department because of pain and swelling in his left leg. He has a history of pancreatic cancer and is currently receiving chemotherapy. Three weeks ago, he had a similar episode in his right arm that resolved without treatment. His temperature is 38.2°C (100.8°F). Palpation of the left leg shows a tender, cord-shaped structure medial to the medial condyle of the femur. The overlying skin is erythematous. Which of the following vessels is most likely affected?

	Answer	Image
A	Posterior tibial artery	
B	Posterior tibial vein	
C	Popliteal artery	
D	Anterior tibial artery	
E	Popliteal vein	
F	Superficial femoral artery	
G	External iliac artery	

	Answer	Image
H	Great saphenous vein	
I	External iliac vein	
J	Deep femoral vein	
K	Anterior tibial vein	

Hint

Recurrent episodes of pain, swelling, and erythema in various locations in a patient with a history of pancreatic cancer are suggestive of malignant migratory thrombophlebitis (Trousseau syndrome).

Correct Answer

A - Posterior tibial artery

Explanation Why

The [posterior tibial artery](#) resides in the deep [posterior compartment of the leg](#). Occlusion of this [artery](#) can be asymptomatic owing to collateral circulation to the foot or show areas of [watershed ischemia](#) marked by pale, cool tissue. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

B - Posterior tibial vein

Explanation Why

The [posterior tibial vein](#) is a deep vein that courses through the deep [posterior compartment of the leg](#). This [vein](#) is susceptible to clotting but classically presents with calf [pain](#), especially on ankle [dorsiflexion](#) ([Homan's sign](#)). It would not be appreciated [medial](#) to the [medial](#) femoral condyles. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

C - Popliteal artery

Explanation Why

The [popliteal artery](#) is the continuation of the [superficial femoral artery](#) as it courses through the [popliteal fossa](#). A clot in this vessel would lead to acute arterial insufficiency in the tissues [distal](#) to the [knee](#) manifesting as [pain](#), pallor, [poikilothermia](#), pulselessness, [paresthesias](#), and paralysis. The patient described here has [pain](#) alone, making limb [ischemia](#) exceptionally unlikely. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

D - Anterior tibial artery

Explanation Why

The [anterior tibial artery](#) runs through the [anterior compartment of the leg](#) and is one of two major arterial inflows to the foot. Occlusion of this [artery](#) can be asymptomatic owing to collateral circulation to the foot or show areas of [watershed ischemia](#) marked by pale, cool tissue. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

E - Popliteal vein

Explanation Why

The [popliteal vein](#) gives rise to the [femoral vein](#) as it passes through the [popliteal fossa](#). This vessel is commonly affected in [migratory thrombophlebitis](#) and would present as a palpable cord. However, it is appreciated in the [popliteal fossa](#) between the tendons of the [hamstring muscles](#).

F - Superficial femoral artery

Explanation Why

The [superficial femoral artery](#) is the principal arterial supply to the leg. A clot in this vessel would lead to signs of acute arterial insufficiency: [pain](#), pallor, [poikilothermia](#), pulselessness, [paresthesias](#), and paralysis. The patient described here has [pain](#) alone, making limb [ischemia](#) exceptionally unlikely. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.


G - External iliac artery

Explanation Why

The external iliac artery courses through the deep [pelvis](#). A clot in that vessel would lead to [critical limb ischemia](#) of the entire ipsilateral leg manifesting as [pain](#), pallor, [poikilothermia](#), pulselessness, [paresthesias](#), and paralysis. The patient described here has [pain](#) alone, making limb [ischemia](#) exceptionally unlikely. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

H - Great saphenous vein

Image



Explanation Why

The [great saphenous vein](#) (GSV) originates at the base of the hallux, continues proximally by passing [anterior](#) to the [medial malleolus](#) and then courses along the [posterior](#) aspect of the [medial](#) femoral

condyle before running deep into the thigh to join the [femoral vein](#). The GSV and the [small saphenous vein](#) make up the main [veins](#) of superficial venous system of the leg, which is commonly involved in cases of malignant [migratory thrombophlebitis](#). [Migratory thrombophlebitis](#) in [pancreatic cancer](#) is believed to be a direct result of [procoagulants](#) by the [tumor](#) itself. This patient will require acute treatment with intravenous [heparin](#) followed by outpatient anticoagulation.

I - External iliac vein

Explanation But

Acute, massive clot in the iliac venous system can lead to [phlegmasia cerulea dolens](#).

Explanation Why

The external iliac vein is a deep vein of the [pelvis](#) and [proximal](#) thigh. Clots in this vessel are exceptionally dangerous as they can easily migrate into the [inferior vena cava](#) and the pulmonary vasculature. However, this [vein](#) cannot be palpated on clinical exam and requires imaging to evaluate for clots. In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

J - Deep femoral vein

Explanation Why

The deep [femoral vein](#) travels alongside the [deep femoral artery](#) deep in the thigh. This [vein](#) is a common location for [deep venous thrombosis](#) and [thrombophlebitis](#) alike, but it would not be palpated [medial](#) to the [medial](#) condyle of the [femur](#). In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

K - Anterior tibial vein

Explanation Why


The [anterior](#) tibial [vein](#) is a deep vein that runs parallel to the [anterior tibial artery](#) in the [anterior compartment of the leg](#). In cases of [thrombophlebitis](#), [skin](#) changes and [pain](#) are present along the

[anterior](#) aspect of the leg but the [vein](#) can be difficult to palpate due to the overlying muscle bellies and the curved surface of the underlying [tibia](#). In addition, malignant [migratory thrombophlebitis](#) generally affects superficial veins.

Question # 26

A 47-year-old woman comes to the emergency department after coughing up 2 cups of bright red blood. A CT angiogram of the chest shows active extravasation from the right bronchial artery. A coil embolization is planned to stop the bleeding. During this procedure, a catheter is first inserted into the right femoral artery. Which of the following represents the most likely subsequent route of the catheter to the source of the bleeding?

	Answer	Image
A	Thoracic aorta, right subclavian artery, right internal thoracic artery, right bronchial artery	
B	Thoracic aorta, pulmonary trunk, right pulmonary artery, right bronchial artery	
C	Thoracic aorta, right posterior intercostal artery, right bronchial artery	
D	Thoracic aorta, right superior epigastric artery, right bronchial artery	

	Answer	Image
E	Thoracic aorta, right pulmonary artery, right bronchial artery	 Brachiocephalic artery Aorta Superior vena cava Right pulmonary artery Pulmonary trunk Right pulmonary veins Pulmonary valve Right atrium Opening of coronary sinus Tricuspid valve Right ventricle Inferior vena cava Left common carotid artery Left subclavian artery Ligamentum arteriosum Left pulmonary artery Left pulmonary veins Left atrium Aortic valve Mitral valve Tendinous cords Left ventricle Papillary muscle Interventricular septum

Hint

The bronchial arteries supply the lung tissue with oxygenated blood via the systemic circulatory system. Bleeding from the bronchial arteries is responsible for approx. 90% of cases of hemoptysis.

Correct Answer

A -

Thoracic aorta, right subclavian artery, right internal thoracic artery, right bronchial artery


Explanation Why

The [internal thoracic artery](#) originates from the [subclavian artery](#) and supplies oxygenated blood to the [anterior chest wall](#) and [breasts](#). However, unlike the [left subclavian artery](#), the [right subclavian artery](#) is not a direct branch of the [thoracic aorta](#) but arises from the brachiocephalic [artery](#). In addition, the right bronchial [artery](#) does not commonly originate from the [internal thoracic artery](#).

B -

Thoracic aorta, pulmonary trunk, right pulmonary artery, right bronchial artery

Image



Explanation Why

The [pulmonary arteries](#) are branches of the [pulmonary trunk](#), which originates from the [right ventricle](#) (not from the [thoracic aorta](#)). They carry deoxygenated blood to the [lungs](#) for [gas exchange](#) and oxygenation. The [bronchial arteries](#), on the other hand, carry oxygenated blood to supply to [lung](#) parenchyma. They do not originate from the [pulmonary arteries](#).

C - Thoracic aorta, right posterior intercostal artery, right bronchial artery

Explanation But

While the [pulmonary arteries](#) supply almost 99% of the blood flow to the [lungs](#), the blood they carry

is deoxygenated and participates in [gas exchange](#) rather than supplying nourishment. The blood of the [bronchial arteries](#), the 2nd of the 2 vascular systems of the [lungs](#), supplies only 1–2% of the blood flow to the [lungs](#) but provides oxygen to the [lungs'](#) supporting structures.

Explanation Why

The right bronchial [artery](#) has several possible origins, including directly from the [thoracic aorta](#), from the left superior bronchial [artery](#), and, most commonly, from one of the right [posterior intercostal arteries](#) (sometimes referred to as the intercostobronchial trunk). The [posterior intercostal arteries](#) branch directly from the [thoracic aorta](#). Therefore, in order to embolize the right bronchial [artery](#) from the right [femoral artery](#), the catheter would need to travel to the [thoracic aorta](#) and then to one of the [intercostal arteries](#) before arriving in the right bronchial [artery](#). On the left side, there are typically two [bronchial arteries](#) that originate directly from the [thoracic aorta](#).


D - Thoracic aorta, right superior epigastric artery, right bronchial artery

Explanation Why

The superior epigastric artery originates from the [internal thoracic artery](#) and supplies oxygenated blood to the [anterior abdominal wall](#) and diaphragm. However, it does not supply the [chest wall](#) or [lung](#) tissue, and the right bronchial [artery](#) does not originate from it.

E - Thoracic aorta, right pulmonary artery, right bronchial artery

Image




Explanation Why

The [pulmonary arteries](#) originate from the [right ventricle](#), not the [left atrium](#), and carry deoxygenated blood to the [lungs](#) for [gas exchange](#) and oxygenation. The [bronchial arteries](#) do not originate from the [pulmonary arteries](#).

Question # 27

Three weeks after undergoing transurethral prostate resection for benign prostatic hyperplasia, a 70-year-old man has fever, malaise, and pain in his extremities. Physical examination shows subungual petechiae and tender red papules on his fingers and toes. A new holosystolic murmur is heard on chest auscultation. A photomicrograph of a Gram stain of an isolate from blood culture is shown. The organism does not cause hemolysis on blood agar. Addition of pyrrolidonyl- β -naphthylamide gives the bacterial colonies a cherry red color. Which of the following is the most likely causal organism?



	Answer	Image
A	<i>Staphylococcus aureus</i>	
B	<i>Enterococcus faecalis</i>	
C	<i>Pseudomonas aeruginosa</i>	

	Answer	Image
D	<i>Streptococcus mutans</i>	
E	<i>Streptococcus gallolyticus</i>	
F	<i>Cardiobacterium hominis</i>	
G	<i>Staphylococcus epidermidis</i>	

Hint

Presence of a new holosystolic murmur, Osler nodes, and splinter hemorrhages is suggestive of infective endocarditis. Photomicrograph and blood agar culture reveal gram-positive cocci that are γ -hemolytic and PYR-positive.

Correct Answer

A - *Staphylococcus aureus*

Explanation Why

[*Staphylococcus aureus*](#) is a gram-positive coccus and the most common causative agent for acute [infective endocarditis](#). However, [*S. aureus*](#) is typically [β-hemolytic](#) and PYR-negative unlike the [γ-hemolytic](#), PYR-positive organisms seen here.

B - *Enterococcus faecalis*

Explanation Why

[*Enterococcus faecalis*](#) is a gram-positive, PYR positive, catalase-negative coccus that can grow in 6.5% NaCl and produces variable [hemolysis](#) ([α-hemolytic](#) or [γ-hemolytic](#)) on blood agar. It can cause [subacute infective endocarditis](#) in patients who undergo gastrointestinal or genitourinary procedures (e.g., transurethral [prostate](#) resection). [Enterococcus](#) spp. are resistant to drugs typically used to treat [infective endocarditis](#) such as [penicillin G](#), 1st to 4th generation [cephalosporins](#), and [vancomycin](#) (in the case of [VRE](#)). [Enterococcus](#) is sensitive to [aminopenicillins](#) and 5th generation [cephalosporins](#) (e.g., [ceftaroline](#)).

C - *Pseudomonas aeruginosa*

Explanation Why

[*Pseudomonas aeruginosa*](#) is a gram-negative [β-hemolytic](#) PYR-negative [bacillus](#) that has been associated with [infective endocarditis](#) in patients with IV drug use (IVDU). This patient has no history of IVDU, and the presence of gram-positive, [γ-hemolytic](#), PYR-positive bacteria makes pseudomonal [endocarditis](#) unlikely.

D - *Streptococcus mutans*

Explanation Why

Streptococcus mutans (a type of [viridans streptococci](#)) is a gram-positive coccus that can cause [infective endocarditis](#). However, *Streptococcus mutans* is [α-hemolytic](#), unlike the [γ-hemolytic](#) organisms seen here, and it typically causes [endocarditis](#) in patients with previously damaged [heart valves](#) who undergo a dental procedure, unlike this patient who develops [endocarditis](#) after undergoing a genitourinary procedure.

E - *Streptococcus gallolyticus*

Explanation Why

Streptococcus gallolyticus is a [γ-hemolytic](#), gram-positive coccus that can cause [endocarditis](#). However, *S. gallolyticus* would show a negative PYR test, and it typically causes [endocarditis](#) in patients with [colon cancer](#) or in patients who undergo GI procedures (e.g., [colon](#) surgery), unlike this patient who develops [endocarditis](#) after undergoing a genitourinary procedure.

F - *Cardiobacterium hominis*

Explanation Why

Cardiobacterium hominis belongs to the [HACEK group](#) of organisms, which can cause [infective endocarditis](#). However, [HACEK](#) organisms are fastidious (i.e., cannot be cultured), and *Cardiobacterium hominis* is a gram-negative coccobacillus, unlike the gram-positive organisms here.






G - *Staphylococcus epidermidis*


Explanation Why

Staphylococcus epidermidis is a [γ-hemolytic](#), gram-positive coccus that can cause [endocarditis](#). However, *S. epidermidis* would show a negative PYR test, and it typically causes [endocarditis](#) in patients with [prosthetic heart valves](#), unlike this patient who develops [endocarditis](#) after undergoing a genitourinary procedure.

Question # 28

A 5-day-old boy is brought to the emergency department by his mother because of a 2-day history of difficulty feeding and multiple episodes of his lips turning blue. He was born at home via spontaneous vaginal delivery and Apgar scores were 7 and 8 at 1 and 5 minutes, respectively. Physical examination shows grunting and moderate intercostal and subcostal retractions. Echocardiography shows a single vessel exiting from the heart. Which of the following is the most likely underlying cause of this patient's condition?

	Answer	Image
A	Failure of neural crest cell migration	
B	Insufficient growth of endocardial cushions	1  2  3 
C	Abnormal placement of the infundibular septum	1  2 

	Answer	Image
D	Absent fusion of septum primum and septum secundum	
E	Abnormal cardiac looping	

Hint

This patient's symptoms (poor feeding, cyanosis, respiratory distress) in early infancy and echocardiography results are consistent with persistent truncus arteriosus.

Correct Answer


A - Failure of neural crest cell migration

Explanation Why

During cardiac development, failure of [neural crest cell](#) migration leads to a lack of [aorticopulmonary septum](#) formation, resulting in a single outflow vessel that receives output from both the right and left ventricles; this condition is known as persistent [truncus arteriosus](#). The mixing of oxygen-rich blood from the [left ventricle](#) with the oxygen-poor blood from the [right ventricle](#) leads to a decrease in overall oxygenation of the blood going into the [systemic circulation](#), causing [cyanosis](#), as seen in this patient with episodic blue [lips](#). This patient also has poor feeding, grunting, and intercostal and subcostal retractions, which are concerning for respiratory distress caused by pulmonary congestion secondary to an increase in overall blood flow to the [lungs](#).

B - Insufficient growth of endocardial cushions

Image




Explanation Why

Insufficient growth of [endocardial cushions](#) leads to defects in the atrial and ventricular septa, resulting in [atrial septal defects](#) (ASD) and [ventricular septal defects](#) (VSD), respectively. Large defects in the cardiac septa can lead to [pulmonary hypertension](#) as a result of [left-to-right shunting](#), which would explain this patient's respiratory distress. While there is a strong association between a [persistent truncus arteriosus](#) and a large [VSD](#), patients with only an ASD or [VSD](#) are usually acyanotic, and the [echocardiography](#) should only show the septal defect with a normal cardiac outflow tract.

C - Abnormal placement of the infundibular septum

Image



Explanation Why

Abnormal placement of the infundibular septum leads to the characteristic findings in [Tetralogy of Fallot](#) (TOF). While TOF can also lead to the blue [lips](#) and respiratory distress seen in this patient, an [echocardiography](#) would instead show [right ventricular outflow tract obstruction](#), [right ventricular hypertrophy](#), [VSD](#), and an overriding aorta.


D - Absent fusion of septum primum and septum secundum

Explanation Why

Absent fusion of the [septum primum](#) and [septum secundum](#) leads to an opening in the atrial septum known as a [patent foramen ovale \(PFO\)](#). A [PFO](#) causes a mild [left-to-right shunt](#) that is usually asymptomatic and does not cause [cyanosis](#) or respiratory distress, unlike in this patient. Also, an [echocardiography](#) would instead show a shunt caused by the patent opening and a normal cardiac outflow tract.

E - Abnormal cardiac looping

Image



Explanation Why

Abnormal [cardiac looping](#) results in an abnormal arrangement of the [heart](#) (e.g., dextrocardia), typically in a condition known as [Kartagener syndrome](#). [Kartagener syndrome](#) may cause respiratory

distress (as in this patient) due to dysfunctional cilia of the [respiratory epithelium](#), as well as [recurrent sinusitis](#) and otitis, [hearing loss](#), and impaired [fertility](#). However, [echocardiography](#) would instead show a [heart](#) in the right side of the thorax.

Question # 29

Six hours after birth, a newborn boy is evaluated for tachypnea. He was delivered at 41 weeks' gestation via Caesarian section and the amniotic fluid was meconium-stained. His respiratory rate is 75/min. Physical examination shows increased work of breathing. X-rays of the abdomen and chest show no abnormalities. Echocardiography shows elevated pulmonary artery pressure. He is started on an inhaled medication that increases smooth muscle cGMP, and there is immediate improvement in his tachypnea and oxygenation status. Three hours later, the newborn is tachypneic and there is blue-grey discoloration of the lips, fingers, and toes. Which of the following is the most likely cause of this infant's cyanosis?

	Answer	Image
A	Increase in concentration of serum myoglobin	
B	Closure of the ductus arteriosus	
C	Oxidization of Fe^{2+} to Fe^{3+}	
D	Inhibition of mitochondrial cytochrome c oxidase	

	Answer	Image																																				
E	Allosteric alteration of heme groups	The graph illustrates the oxygen saturation of myoglobin and hemoglobin as a function of partial pressure of oxygen (pO₂). The y-axis represents O₂ saturation in percent, ranging from 0 to 100. The x-axis represents pO₂ in mmHg, ranging from 0 to 100. Myoglobin is represented by a hyperbolic curve that rises steeply at low pO₂ and reaches near 100% saturation by pO₂ = 40 mmHg. Hemoglobin is represented by a sigmoidal curve that shows cooperative binding, with a P₅₀ (the pO₂ at which hemoglobin is 50% saturated) indicated at approximately 37 mmHg. Two vertical shaded regions are marked: 'Venous blood from peripheral tissue' (pO₂ ~20-40 mmHg) and 'Arterial blood from the lungs' (pO₂ ~80-100 mmHg). A horizontal dashed line at 50% saturation intersects the hemoglobin curve at its P₅₀. <table border="1"><caption>Approximate data points from the O₂ saturation graph</caption><thead><tr><th>pO₂ (mmHg)</th><th>Myoglobin Saturation (%)</th><th>Hemoglobin Saturation (%)</th></tr></thead><tbody><tr><td>0</td><td>0</td><td>0</td></tr><tr><td>10</td><td>~80</td><td>~10</td></tr><tr><td>20</td><td>~95</td><td>~30</td></tr><tr><td>30</td><td>~98</td><td>~55</td></tr><tr><td>40</td><td>~99</td><td>~75</td></tr><tr><td>50</td><td>~100</td><td>~85</td></tr><tr><td>60</td><td>~100</td><td>~90</td></tr><tr><td>70</td><td>~100</td><td>~93</td></tr><tr><td>80</td><td>~100</td><td>~95</td></tr><tr><td>90</td><td>~100</td><td>~97</td></tr><tr><td>100</td><td>~100</td><td>~98</td></tr></tbody></table>	pO ₂ (mmHg)	Myoglobin Saturation (%)	Hemoglobin Saturation (%)	0	0	0	10	~80	~10	20	~95	~30	30	~98	~55	40	~99	~75	50	~100	~85	60	~100	~90	70	~100	~93	80	~100	~95	90	~100	~97	100	~100	~98
pO ₂ (mmHg)	Myoglobin Saturation (%)	Hemoglobin Saturation (%)																																				
0	0	0																																				
10	~80	~10																																				
20	~95	~30																																				
30	~98	~55																																				
40	~99	~75																																				
50	~100	~85																																				
60	~100	~90																																				
70	~100	~93																																				
80	~100	~95																																				
90	~100	~97																																				
100	~100	~98																																				

Hint

Elevated pulmonary artery pressure raises suspicion for persistent pulmonary hypertension of the newborn (PPHN), a condition in which the pulmonary vascular resistance remains abnormally high after birth. This results in right-to-left shunting of blood via the fetal circulatory pathways, leading to insufficient oxygenation (cyanosis) and tachypnea. Since the pathophysiology of PPHN is aberrantly constricted pulmonary vasculature, a vasodilator in the form of inhaled nitric oxide is given to reduce shunting across the ductus and foramen ovale.

Correct Answer

A - Increase in concentration of serum myoglobin

Explanation Why

Increased serum [myoglobin](#) can occur as a result of [rhabdomyolysis](#) and often manifests with myalgias, weakness, and dark [urine](#) ([myoglobinuria](#)). [Rhabdomyolysis](#) can be of traumatic (e.g., [crush injury](#) or direct injury) or nontraumatic cause (e.g., [seizures](#), substance intoxication, or adverse drug reactions of [statins](#) and [neuroleptics](#)). However, this [newborn](#) was delivered by [cesarean section](#) and has no [risk factors](#) or medications that would suggest [rhabdomyolysis](#) or myoglobinemia.

B - Closure of the ductus arteriosus

Explanation Why

Closure of the [ductus arteriosus](#) can cause [cyanosis](#) in patients with ductus-dependent anomalies including [Tetralogy of Fallot](#) (ToF), pulmonary atresia, critical [pulmonary stenosis](#), and [transposition of the great arteries](#). Since the [newborn](#)'s chest [x-ray](#) was normal and the [echocardiography](#) showed only elevated [pulmonary artery](#) pressure, a ductus-dependent anomaly is an unlikely cause for [cyanosis](#) in this patient. In a patient with [persistent pulmonary hypertension of the newborn](#) (PPHN), which is the most likely diagnosis here, closure of the ductus would likely improve [cyanosis](#), since it is being caused by right-to-left shunting.

C - Oxidization of Fe^{2+} to Fe^{3+}


Explanation Why

Inhaled nitric oxide (NO) is the mainstay treatment for [PPHN](#) because of its ability to potently [vasodilate](#) the pulmonary vasculature (reducing [V/Q mismatch](#)) without decreasing [systemic vascular resistance](#). NO induces [smooth muscle](#) relaxation (and subsequent [vasodilation](#)) by activating [guanylyl cyclase](#), which increases [cGMP](#) synthesis. A side effect of inhaled NO is the oxidation of ferrous [iron](#) (Fe^{2+}) to ferric [iron](#) (Fe^{3+}) at high levels, resulting in the formation of [methemoglobin](#).

Because Fe^{3+} has a reduced affinity for oxygen binding, [methemoglobin](#) is a poor oxygen transporter. Therefore, [methemoglobinemia](#) results in poor tissue oxygenation and causes clinical signs of [cyanosis](#). Other side effects of inhaled NO in [neonates](#) include [platelet dysfunction](#), [pulmonary edema](#), and the formation of peroxynitrites.

D - Inhibition of mitochondrial cytochrome c oxidase

Image




Explanation Why

[Cyanide poisoning](#) inhibits [mitochondrial cytochrome c oxidase](#), which disrupts the [electron transport chain](#) and results in [anaerobic metabolism](#) and [lactic acidosis](#). It is typically caused by enclosed-space explosions or fires, or by high-dose [sodium nitroprusside](#) treatment. Patients typically present with confusion, [headache](#), nausea, vomiting, [dyspnea](#), flushing, and breath that smells of [bitter](#) almonds. Despite its name, [cyanide poisoning](#) does not cause [cyanosis](#) and is not caused by inhaled [nitric oxide](#).

E - Allosteric alteration of heme groups

Image




Explanation Why

[Carboxyhemoglobin](#) is formed as a result of carbon monoxide binding to [hemoglobin](#), which results in an allosteric change that shifts the [oxygen-hemoglobin dissociation curve](#) to the left. This increases the affinity of [hemoglobin](#) for oxygen and reduces oxygen unloading in the peripheral tissues. [Carbon monoxide poisoning](#) is typically caused by exposure to fires, exhaust, or heaters, and manifests with cherry-red [skin](#) rather than [cyanosis](#). [Carboxyhemoglobinemia](#) does not occur secondary to [persistent pulmonary hypertension of the newborn](#) or administration of [nitric oxide](#).

Question # 30

A 60-year-old woman with ovarian cancer comes to the physician with a 5-day history of fever, chills, and dyspnea. She has a right subclavian chemoport in which she last received chemotherapy 2 weeks ago. Her temperature is 39.5°C (103.1°F), blood pressure is 110/80 mm Hg, and pulse is 115/min. Cardiopulmonary examination shows jugular venous distention and a new, soft holosystolic murmur heard best in the left parasternal region. Crackles are heard at both lung bases. Echocardiography shows a vegetation on the tricuspid valve. Peripheral blood cultures taken from this patient is most likely to show which of the following findings?

	Answer	Image
A	Gram-positive, catalase-negative, α -hemolytic, optochin-resistant cocci in chains	
B	Gram-positive, catalase-positive, coagulase-negative, novobiocin-resistant cocci in clusters	
C	Gram-positive, catalase-positive, coagulase-positive cocci in clusters	
D	Gram-positive, catalase-negative, nonhemolytic, salt-sensitive cocci in chains	
E	Gram-positive, catalase-negative, β -hemolytic, bacitracin-resistant cocci in chains	

Hint

This patient presents with features of infective endocarditis (2 major Duke criteria: a new valvular murmur, valvular vegetations on echocardiography). The chemoport is the most likely route of infection.

Correct Answer

A -

Gram-positive, catalase-negative, α -hemolytic, optochin-resistant cocci in chains

Explanation Why

The above microbiologic findings would indicate [Viridans streptococci](#) infection. However, these bacteria usually cause infection following dental procedures and afflict previously damaged native valves or [prosthetic valves](#), neither of which are present in this patient. Moreover, [Viridans streptococci](#) typically cause subacute infection, whereas this patient presents with a 5-day history.

B -

Gram-positive, catalase-positive, coagulase-negative, novobiocin-resistant cocci in clusters

Explanation But


[Staphylococcus epidermidis](#), a bacterium commonly causing line-associated [endocarditis](#), has similar microbiologic features but is novobiocin sensitive.

Explanation Why

The above microbiologic findings would indicate [Staphylococcus saprophyticus](#) infection. This pathogen typically causes [urinary tract infections](#) rather than [endocarditis](#).

C - Gram-positive, catalase-positive, coagulase-positive cocci in clusters

Image



Explanation But

In this clinical scenario, infection with [*Staphylococcus epidermidis*](#) would be a reasonable working diagnosis as well, especially since this bacterium is commonly responsible for infections via peripheral venous catheters. However, *S. epidermidis* is coagulase-negative.

Explanation Why

The new [holosystolic murmur](#) in the left parasternal region and the [echocardiographic](#) findings indicate affliction of the [tricuspid valve](#). Blood culture findings of gram-positive, catalase-positive, [coagulase](#)-positive [cocci](#) in clusters are characteristic of [*Staphylococcus aureus*](#), which is the most common pathogen that causes acute [infective endocarditis](#) (including in patients with previously healthy valves).

D -

Gram-positive, catalase-negative, nonhemolytic, salt-sensitive cocci in chains

Explanation Why

The above microbiologic findings would indicate [*Streptococcus gallolyticus*](#) infection. Infection with this bacterium is associated with [colorectal cancer](#) but not with [ovarian cancer](#). Moreover, *S. gallolyticus* is only responsible for a small share of [infective endocarditis](#) cases and the gastrointestinal surface is a far more likely portal of entry than a [central venous access](#).

E -


Gram-positive, catalase-negative, β -hemolytic, bacitracin-resistant cocci in chains

Explanation Why

The above microbiologic findings would indicate [*Streptococcus agalactiae*](#) infection. While this pathogen can cause respiratory symptoms in adults, it is usually responsible for neonatal infections (particularly [neonatal meningitis](#)). It is not associated with [infective endocarditis](#).

Question # 31

A 27-year-old man comes to the physician because of a 2-month history of palpitations and shortness of breath on exertion. He has no history of serious illness. He does not smoke or use illicit drugs. His pulse is 90/min, respirations are 18/min, and blood pressure is 140/40 mm Hg. Cardiac examination shows a murmur along the left sternal border. A phonocardiogram of the murmur is shown. Which of the following additional findings is most likely in this patient?



	Answer	Image
A	Increased left ventricular end-diastolic volume	

	Answer	Image
B	Decreased left ventricular wall compliance	
C	Decreased left ventricular wall stress	
D	Decreased pulmonary capillary wedge pressure	
E	Decreased left ventricular afterload	
F	Increased right ventricular oxygen saturation	


Hint

This patient's symptoms and decrescendo diastolic murmur along the left sternal border are consistent with aortic regurgitation, which is likely due to a congenital bicuspid aortic valve (given his age and the absence of other risk factors, e.g., intravenous drug use, Marfan syndrome, history of rheumatic fever).

Correct Answer

A - Increased left ventricular end-diastolic volume

Image



Explanation Why

[Aortic regurgitation](#) causes an increase in [left ventricular end-diastolic volume](#), as the [left ventricle](#) receives blood from both the [left atrium](#) and the regurgitant aorta. This results in hyperdynamic circulation and clinically manifests with [palpitations](#) and a widened [pulse pressure](#), as seen in this patient. Patients with chronic AR can be asymptomatic for decades before developing [symptoms of left heart failure](#). The recent exertional dyspnea in this patient indicates the onset of [left heart failure](#), as the [left ventricle](#) is unable to maintain adequate [cardiac output](#) during exertion.

B - Decreased left ventricular wall compliance

Explanation Why

[Left ventricular](#) wall compliance is increased in chronic [aortic regurgitation](#) because the [left ventricle](#) gradually enlarges and remodels due to the intermittent reversal of blood flow. [Eccentric hypertrophy](#) causes the [left ventricle](#) to become larger and develop a comparatively thin, more compliant wall.

C - Decreased left ventricular wall stress

Explanation Why

Decreased [left ventricular](#) wall stress is seen in conditions with decreased ventricular pressure, e.g., [hypovolemic shock](#). This patient, however, presents with [aortic regurgitation](#), which causes increased [left ventricular](#) wall stress due to intermittent blood flow reversal.

D - Decreased pulmonary capillary wedge pressure

Explanation Why

Decreased [pulmonary capillary wedge pressure](#) is seen in settings of [distributive shock](#), e.g., [septic shock](#). This patient, however, presents with symptoms of [systolic heart failure](#) secondary to [aortic regurgitation](#), which causes increased left atrial pressure and consequently raised [pulmonary capillary wedge pressure](#). [PCWP](#) can also be normal in compensated [aortic regurgitation](#) but would not be decreased.

E - Decreased left ventricular afterload

Explanation Why

Decreased [left ventricular afterload](#) is seen in conditions involving peripheral [vasodilation](#) (e.g., due to medication use or [distributive shock](#)). This patient, however, presents with [aortic regurgitation](#), which usually causes increased [afterload](#) because intermittent blood flow reversal increases [systolic stress](#).

F - Increased right ventricular oxygen saturation

Explanation Why

[Right ventricular oxygen saturation](#) would be increased in an [atrial septal defect](#). While an [atrial septal defect](#) can also manifest with a [murmur](#) and exertional dyspnea, it is typically characterized by a widely fixed S2, which is not seen here. In addition, an [atrial septal defect](#) would not manifest with increased [pulse pressure](#).

Question # 32

A 28-year-old man comes to the emergency department for an injury sustained while doing construction. Physical examination shows a long, deep, irregular laceration on the lateral aspect of the left forearm with exposed fascia. Prior to surgical repair of the injury, a brachial plexus block is performed using a local anesthetic. Shortly after the nerve block is performed, he complains of dizziness and then loses consciousness. His radial pulse is faint and a continuous cardiac monitor shows a heart rate of 24/min. Which of the following is the most likely mechanism of action of the anesthetic that was administered?

	Answer	Image
A	Inactivation of potassium channels	
B	Inactivation of sodium channels	
C	Activation of acetylcholine receptors	
D	Inactivation of ryanodine receptors	
E	Activation of GABA receptors	

Hint

This patient received a local anesthetic, probably bupivacaine, which can have severe cardiotoxicity when absorbed systemically.

Correct Answer

A - Inactivation of potassium channels

Explanation Why

Inactivation of potassium channels is the mechanism of action of [class III antiarrhythmic drugs](#) such as [amiodarone](#). Side effects of this drug class include cardiovascular depression, including [bradycardia](#), [heart block](#), and [hypotension](#). However, inactivation of potassium channels does not achieve [local anesthesia](#) and so this cannot be used for [brachial plexus](#) block.

B - Inactivation of sodium channels

Explanation Why

Local anesthetics (e.g., [bupivacaine](#)) bind to [voltage-gated sodium channels](#) of the nerve fibers and block them, which leads to an inhibition of nerve excitation and impulse conduction ([pain](#) signals). When absorbed systemically, [bupivacaine](#) also acts as a [class IB antiarrhythmic drug](#) and can cause [hypotension](#) and [bradycardia](#), as seen in this patient. [Ventricular arrhythmias](#) and [asystole](#) may also occur. Of the commonly used local anesthetics, [bupivacaine](#) is most likely to result in cardiovascular toxicity, because of its high [potency](#) and long duration of action.

C - Activation of acetylcholine receptors

Explanation Why

Activation of acetylcholine receptors is the mechanism of action of depolarizing agents such as [succinylcholine](#). Side effects include [hypercalcemia](#), [hyperkalemia](#), and [malignant hyperthermia](#), which may lead to [hypotension](#) and [bradycardia](#). However, [succinylcholine](#) is used for the induction of anesthesia, not for a [brachial plexus](#) block.

D - Inactivation of ryanodine receptors

Explanation Why

Inactivation of [ryanodine receptors](#) is the mechanism of action of [muscle relaxants](#) like [dantrolene](#). Side effects of [dantrolene](#) include [CNS](#) depression, and infrequently, [AV block](#) or [tachycardia](#). However, [dantrolene](#) is not an anesthetic drug used for [brachial plexus](#) block but rather can be used to treat toxicity associated with inhaled anesthetics (e.g., [malignant hyperthermia](#)).



E - Activation of GABA receptors



Explanation Why

Activation of [GABA](#) receptors is the mechanism of action of [benzodiazepines](#) (e.g., [midazolam](#)) and [barbiturates](#) (e.g., [phenobarbital](#)), some of which may be used for general anesthesia or sedation. Side effects may include [CNS](#) and cardiovascular depression, including [bradycardia](#) and [hypotension](#). However, [GABA](#) receptor activation cannot be used to achieve [brachial plexus](#) block.

Question # 33

Shortly after delivery, a female newborn develops bluish discoloration of the lips, fingers, and toes. She was born at term to a 38-year-old primigravid woman. Pregnancy was complicated by maternal diabetes mellitus. Pulse oximetry on room air shows an oxygen saturation of 81%. Echocardiography shows immediate bifurcation of the vessel arising from the left ventricle; the vessel emerging from the right ventricle gives out coronary, head, and neck vessels. An abnormality in which of the following developmental processes most likely accounts for this patient's condition?

	Answer	Image
A	Separation of tricuspid valve tissue from myocardium	 The diagram shows a frontal view of a heart. A blue vessel (ductus arteriosus) is shown connecting the descending aorta to the pulmonary artery. A label 'Patent ductus arteriosus' points to this vessel. Another label 'Patent foramen ovale' points to an opening in the interatrial septum. Arrows indicate the direction of blood flow.
B	Fusion of endocardial cushion	 Three diagrams labeled 1, 2, and 3 show the progression of endocardial cushion fusion. Diagram 1 shows the initial stage with the cushion as a separate structure. Diagram 2 shows the cushion beginning to fuse with the underlying myocardium. Diagram 3 shows the final stage where the cushion is fully fused, forming the atrial septum and the AV valves. Arrows indicate the direction of blood flow.

	Answer	Image
C	Alignment of infundibular septum	
D	Division of aorta and pulmonary artery	
E	Spiraling of aorticopulmonary septum	


Hint

Echocardiography shows the pulmonary trunk arising (posteriorly) from the left ventricle, and the aorta (which gives out coronary, head, and neck arteries) arising (anteriorly) from the right ventricle, which is called transposition of the great vessels. Early cyanosis and poor oxygen saturation are in line with this diagnosis

Correct Answer

A - Separation of tricuspid valve tissue from myocardium

Image




Explanation Why

Failed separation of [tricuspid valve](#) tissue from [myocardium](#) causes [Ebstein anomaly](#). Mild apical displacement is often asymptomatic, whereas individuals with marked [tricuspid valve](#) leaflet displacement and incompetence can present with [cyanosis](#) due to [right-to-left shunting](#) (interatrial) and poor [oxygen saturation](#), as observed in this patient. However, [echocardiography](#) would show apical displacement of the septal [tricuspid valve](#) leaflet, not [transposition of the great vessels](#).

B - Fusion of endocardial cushion

Image




Explanation Why

Impaired fusion of the [endocardial cushion](#) can cause [atrial septal defect](#), [ventricular septal defect](#), or [atrioventricular septal defects](#), all of which are usually asymptomatic at [birth](#). [Infants](#) with complete [atrioventricular septal defects](#) can present with symptoms of [heart failure](#). Untreated, these defects can cause significant [pulmonary hypertension](#) and reversal of the [left-to-right shunt](#) ([Eisenmenger syndrome](#)). However, early [cyanosis](#) is not typical and [echocardiography](#) will show the respective septal defect, not [transposition of the great vessels](#).

C - Alignment of infundibular septum

Image



Explanation Why

Anterosuperior displacement of the infundibular septum is one of the features in [tetralogy of Fallot](#) (TOF) that causes infundibular [pulmonary stenosis](#), TOF's key prognostic determinant. [Infants](#) with TOF can present with early [cyanosis](#) and poor [oxygen saturation](#), as observed in this [newborn](#). However, [echocardiography](#) would show the presence of [right ventricular outflow tract obstruction](#) and [hypertrophy](#), overriding aorta, and [ventricular septal defect](#), not [transposition of the great vessels](#).

D - Division of aorta and pulmonary artery


Explanation Why

Failed development of a septum dividing the aorta and [pulmonary artery](#) causes persistent [truncus](#)

[arteriosus](#). In patients with [truncus arteriosus](#), a common trunk arises from both the right and [left ventricle](#), mixing arterial and venous blood. [Infants](#) will present with early [cyanosis](#) and poor [oxygen saturation](#), as observed in this patient. However, [echocardiography](#) will show a single overriding vessel connected to both ventricles, not [transposition of the great vessels](#).

E - Spiraling of aorticopulmonary septum

Image



Explanation But

[TGV](#) is incompatible with life unless mixing occurs via an intracardiac shunt across a [patent foramen ovale](#), ventricle septal defect, or [atrial septal defect](#), or via an extracardiac connection (e.g., [patent ductus arteriosus](#)).

Explanation Why

In the development of the cardiac outflow tract, [neural crest](#) and endocardial cells migrate to form [truncal ridges](#) and [bulbar ridges](#) from the [truncus arteriosus](#) and [bulbus cordis](#), respectively. Normally, the truncal and [bulbar ridges](#) spiral and fuse to form the [aorticopulmonary septum](#) ([AP septum](#)), which further develops into the [ascending aorta](#) and [pulmonary trunk](#). Failure of the spiraling of the [aorticopulmonary septum](#) can lead to [transposition of the great vessels](#) ([TGV](#)).

[Infants](#) born to diabetic mothers are at increased risk of developing [TGV](#).

Question # 34

A 55-year-old woman is brought to the emergency department because of epigastric pain, sweating, and breathlessness for 45 minutes. She has hypertension treated with hydrochlorothiazide. She has smoked 1 pack of cigarettes daily for the past 30 years and drinks 1 glass of wine daily. Her pulse is 105/min and blood pressure is 100/70 mm Hg. Arterial blood gas analysis on room air shows:

pH	7.49
pCO ₂	32 mm Hg
pO ₂	57 mm Hg

An ECG shows ST elevations in leads V₁–V₆. Which of the following is the most likely cause of hypoxemia in this patient?

	Answer	Image
A	Decreased transpulmonary pressure	
B	Decreased minute ventilation	
C	Increased pulmonary capillary pressure	
D	Decreased total body hemoglobin	
E	Increased pulmonary capillary permeability	

Hint

This patient's multiple cardiovascular risk factors (e.g., hypertension and smoking history) in conjunction with epigastric pain, sweating, dyspnea, and tachycardia should raise concern for myocardial infarction. The ST elevations in the anterior leads are diagnostic of an extensive anterior myocardial infarction.

Correct Answer

A - Decreased transpulmonary pressure

Explanation Why

Decreased transpulmonary pressure can result from a [pneumothorax](#), which also manifests with similar [ABG](#) findings, [dyspnea](#), and [tachycardia](#), as observed in this patient. In addition, this woman's long-term smoking history increases her risk of [spontaneous pneumothorax](#). However, epigastric [pain](#) is not typically associated with [pneumothorax](#). In addition, this patient's [ECG](#) results are not consistent with a [pneumothorax](#).

B - Decreased minute ventilation

Explanation Why

Decreased [minute ventilation](#) can be seen in patients with depression in the [respiratory center](#) of the brain (e.g., [opioid overdose](#)) and in those with impairment of the peripheral nerves or respiratory muscles. A slow breathing rate and/or low [tidal volume](#) results in [hypoxemia](#), as seen on this patient's [ABG](#). However, CO₂ would also accumulate, causing [hypoxic](#) hypercapnic [respiratory acidosis](#), as opposed to the [respiratory alkalosis](#) present here.

C - Increased pulmonary capillary pressure

Explanation But

This patient shows an atypical presentation of [myocardial infarction](#), which is more likely to occur in elderly, diabetic individuals, and women. These patients can present with minimal or no [chest pain](#) while autonomic symptoms are often the [chief complaint](#). When evaluating patients with symptoms like epigastric [pain](#), sweating, and breathlessness, as seen here, [acute coronary syndrome](#) is an important differential diagnosis that must be promptly ruled out.

Explanation Why

Increased [pulmonary capillary wedge pressure](#) can occur in the setting of [myocardial infarction](#) (MI) due to [cardiogenic shock](#). An extensive [myocardial infarction](#), as seen in this patient, may result in decreased cardiac contractility and reduced [cardiac output](#), leading to increased left-sided [heart](#) pressures and ultimately backup of blood to the [lungs](#). Cardiogenic pulmonary edema thickens the alveolar-[capillary](#) membrane and impairs oxygen [diffusion capacity](#). As a compensatory mechanism, [minute ventilation](#) increases, resulting in increased CO₂ elimination. This patient's inability to oxygenate with preserved ventilation, as reflected by the [hypoxemia](#) without concurrent hypercapnia in her [ABG](#), is consistent with [type 1 respiratory failure](#).

D - Decreased total body hemoglobin

Explanation Why

Decreased total body [hemoglobin](#) (Hb) can result from acute blood loss in the setting of a ruptured [aortic aneurysm](#) (AA). A ruptured AA can manifest with [tachycardia](#), [hypotension](#), [dyspnea](#), sweating, and abdominal [pain](#), as seen in this patient. [Risk factors](#) include a history of [hypertension](#) and smoking, both of which are present here. However, patients with AA classically present with severe, tearing [pain](#) in the chest, back, and/or abdomen. Additionally, though decreased [Hb](#) results in a decreased O₂ delivery to tissue ([hypoxia](#)), it would not explain this patient's decreased pO₂ ([hypoxemia](#)) on [ABG](#) or the [ST elevations](#) on [ECG](#).

E - Increased pulmonary capillary permeability

Explanation Why

Increased pulmonary [capillary](#) permeability is seen in [acute respiratory distress syndrome](#) (ARDS). [ARDS](#) manifests with similar [ABG](#) findings, sweating, and [dyspnea](#), which are seen in this patient. A known cause of [ARDS](#) is [acute pancreatitis](#), a condition that manifests with epigastric [pain](#) and is associated with chronic alcohol use. However, this patient only drinks one glass of wine per day, making [acute pancreatitis](#), and therefore [ARDS](#), unlikely. Finally, this patient's [ECG](#) results are not consistent with a diagnosis of [ARDS](#).

Question # 35

A 39-year-old woman comes to the physician because of an 8-month history of progressive fatigue, shortness of breath, and palpitations. She has a history of recurrent episodes of joint pain and fever during childhood. She emigrated from India with her parents when she was 10 years old. Cardiac examination shows an opening snap followed by a late diastolic rumble, which is best heard at the fifth intercostal space in the left midclavicular line. This patient is at greatest risk for compression of which of the following structures?

	Answer	Image
A	Trachea	
B	Thoracic duct	
C	Vagus nerve	
D	Hemiazygos vein	
E	Esophagus	

Hint

This patient's murmur is characteristic of mitral stenosis, which, given her immigration history and recurrent fever and joint pain, is most likely secondary to rheumatic fever. Chronic untreated mitral stenosis results in dilation of the left atrium, which can compress adjacent structures.

Correct Answer

A - Trachea

Explanation Why

The [trachea](#) terminates by bifurcating into the right and left [bronchi](#) superior to the [left atrium](#). Therefore, cardiac compression of the [trachea](#) is unlikely.

B - Thoracic duct

Explanation Why

The [thoracic duct](#) lies in the [posterior mediastinum](#) and is separated from the [heart](#) by the descending [thoracic aorta](#) and the [esophagus](#). It is therefore unlikely to be compressed by [cardiac enlargement](#).

C - Vagus nerve

Explanation Why

The [vagus nerves](#) lie [posterior](#) to the hila of the [lungs](#) bilaterally. They are therefore unlikely to be compressed by [cardiac enlargement](#). In rare cases, compression of the left [recurrent laryngeal nerve](#) by a significantly enlarged [left atrium](#) may occur in patients with chronic [mitral stenosis](#). Patients may then present with hoarseness of voice.

D - Hemiazygos vein

Explanation Why

The [hemiazygos vein](#) lies in the [posterior mediastinum](#), which is separated from the [heart](#) by the descending [thoracic aorta](#) and the [esophagus](#). It is therefore unlikely to be compressed by [cardiac enlargement](#).

E - Esophagus

Explanation Why

Because the [posterior mediastinal](#) portion of the [esophagus](#) lies just behind the [left atrium](#), it is at risk of being compressed and/or displaced by an enlarged [left atrium](#). [Left atrial enlargement](#) due to increased left atrial pressure can develop in patients with chronic [mitral stenosis](#), like this patient, although this occurs very rarely.

Question # 36

An investigator is studying cardiomyocytes in both normal and genetically modified mice. The genetic modification is found to result in less coordinated muscle contractions. Both the normal and genetically modified mice are observed after aerobic exercise and their heart rates are recorded and compared. After a 10-minute session on a treadmill, the average pulse measured in the normal mice is 680/min, whereas in the genetically modified mice it is only 160/min. Which of the following is most likely to account for the increased heart rate seen in the normal mice?

	Answer	Image
A	Greater cardiomyocyte size	
B	Lower opening rate of funny current channels	
C	Greater ratio of heart to body weight	
D	Lower number of gap junctions	
E	Greater T-tubule density	

Hint

An increased heart rate requires both rapid cycling of Ca^{2+} and rapid, coordinated excitation-contraction coupling of myocytes.

Correct Answer

A - Greater cardiomyocyte size

Explanation Why

Increased [cardiomyocyte](#) size is not related to contraction rate. The contractile machinery of [cardiomyocytes](#) consists of [myofilaments](#) that comprise 45–60% of the cell's volume. Although [hypertrophy](#) of [myofilaments](#) may increase the force of contraction, it would not influence the rate of excitation-contraction coupling and, therefore, would not affect [heart rate](#) or coordination of muscle contractions.

B - Lower opening rate of funny current channels

Explanation Why

[Funny current](#) channels facilitate the influx of Na^+ and K^+ at the start of phase 4 of the pacemaker cell [action potential](#). The [funny current](#) is responsible for the [automaticity](#) of pacemaker cells. Increased influx through these channels results in an earlier onset of [depolarization](#). Therefore, a lower opening rate of the channels that facilitate this current would cause a decrease in [heart rate](#), not an increase. Additionally, a lower opening rate of these channels would not explain the less coordinated muscle contractions.

C - Greater ratio of heart to body weight

Explanation Why

A greater [ratio](#) of [heart](#) to body weight would not explain the increased [heart rate](#) or less coordinated muscle contractions seen here. Although studies have shown higher [heart rates](#) in smaller species (e.g., mice are capable of achieving higher [heart rates](#) than humans), a direct association between increased [heart](#) to body weight [ratio](#) has not been found. A different mechanism is more likely responsible for the rapid excitation-contraction coupling occurring in the [cardiomyocytes](#) of the normal mice.

D - Lower number of gap junctions

Explanation Why

[Gap junctions](#) enable the electrical coupling and synchronized conduction of impulses between [cardiomyocytes](#). A lower number of [gap junctions](#) would, therefore, result in slower conduction and thus a lower [heart rate](#).

E - Greater T-tubule density

Explanation Why

Greater [T-tubule](#) density is most likely to account for the increased [heart rate](#) seen in the normal mice. In both mice and humans, [T-tubules](#) ensure coordinated contractions in [striated muscle](#) cells (i.e., both skeletal and [cardiac muscle cells](#)). The dense organization of [T-tubules](#) within the [sarcoplasmic reticulum](#) (SR) allows for optimal cardiac excitation-contraction coupling to occur and facilitates the synchronous activation of rapid calcium release among the [cardiomyocytes](#), thereby ensuring a coordinated contraction. Mice have a much higher cardiac [T-tubule](#) density than humans, which allows them to achieve a much higher [heart rate](#).

Question # 37

A previously healthy 32-year-old woman comes to the emergency department with fatigue and bilateral leg swelling. Her pulse is 92/min, and respirations are 24/min. Physical examination shows jugular venous distention and pitting edema of the lower extremities. Her abdomen is distended with shifting dullness and tender hepatomegaly is present. Cardiovascular examination shows a holosystolic murmur heard best at the left lower sternal border that increases in intensity with inspiration. Which of the following is the most likely predisposing factor for this patient's condition?

	Answer	Image
A	45,XO genotype	
B	Intravenous drug use	
C	Vitamin B ₁ deficiency	
D	Fibrillin gene defect	
E	Streptococcal pharyngitis	

Hint

This patient has signs of right heart failure (JVD, hepatomegaly, peripheral edema) and a murmur that is consistent with tricuspid regurgitation. TR is best auscultated at the left parasternal region of the 4th ICS and manifests as a holosystolic murmur that gets louder on inspiration (Carvallo phenomenon).

Correct Answer

A - 45,XO genotype

Explanation Why

The [45,XO genotype](#) causes [Turner syndrome](#). The most common cardiovascular abnormalities in [Turner syndrome](#) are [bicuspid aortic valve](#), [coarctation of the aorta](#), and [aortic dissection](#). Chronic and/or severe [aortic valve](#) disease could lead to [left heart failure](#), which typically manifests with [dyspnea](#), orthopnea, [paroxysmal nocturnal dyspnea](#), and bilateral basilar [rales](#). However, this patient presents with features of [right heart failure](#). Left [heart murmurs](#) do not increase on inspiration, unlike the auscultatory findings in this patient. Moreover, other typical manifestations of [Turner syndrome](#) (e.g., [short stature](#), [webbed neck](#), widely spaced nipples, [amenorrhea](#)) would also be expected.

B - Intravenous drug use

Explanation Why

Acquired [tricuspid valve regurgitation](#) and signs of right-sided [heart failure](#) in a previously healthy patient suggest a diagnosis of [tricuspid valve endocarditis](#), for which intravenous drug use (IVDU) is the most likely predisposing factor. The most commonly isolated pathogens in [infective endocarditis](#) due to IVDU are [Staphylococcus aureus](#) (most common), [streptococci](#), [enterococci](#), and [Pseudomonas aeruginosa](#). Additional causes of [tricuspid valve regurgitation](#) in a young patient include [rheumatic heart disease](#), [right-sided heart failure](#), and [connective tissue disease](#) (e.g., [Marfan syndrome](#)).

C - Vitamin B₁ deficiency

Explanation Why

Severe [vitamin B₁ deficiency](#) (usually secondary to [alcohol use disorder](#) in developed countries) results in [wet beriberi](#). [Wet beriberi](#) typically manifests as high-output cardiac failure. Although examination would reveal [JVD](#), [ascites](#), and [edema](#) (as seen in this patient), this patient lacks other

typical features of high-output cardiac failure such as [dyspnea](#), [tachypnea](#), orthopnea, [paroxysmal nocturnal dyspnea](#), bounding peripheral pulses, and an [S3 gallop](#). In addition, a [holosystolic murmur](#) at the left lower sternal border would not be expected in such patients.

D - Fibrillin gene defect

Explanation Why


Defects in the [fibrillin gene](#) are responsible for [Marfan syndrome](#). The most common cardiovascular structures involved in [Marfan syndrome](#) are the aorta and valves of the left [heart](#) (e.g., [aortic root dilation](#), [aortic regurgitation](#), [mitral valve prolapse](#)). Chronic and/or severe left [valvular heart disease](#) could lead to [left heart failure](#), which typically manifests with [dyspnea](#), orthopnea, [paroxysmal nocturnal dyspnea](#), and bilateral basilar [rales](#). However, this patient presents with features of [right heart failure](#). Left [heart murmurs](#) (except for [mitral valve prolapse](#)) do not increase on inspiration, unlike the [murmur](#) in this patient. Moreover, other typical features of [Marfan syndrome](#) (e.g., [joint hypermobility](#), [lens subluxation](#)) would also be expected.

E - Streptococcal pharyngitis

Explanation Why

Untreated [streptococcal pharyngitis](#) (caused by group A [streptococcus](#)) can lead to [rheumatic heart disease](#) (RHD) that can manifest as [heart failure](#) in young or middle-aged individuals. However, as RHD mainly affects the valves of the left [heart](#) (i.e., the mitral and/or aortic valves), features of [left heart failure](#) (e.g., [dyspnea](#), orthopnea, [paroxysmal nocturnal dyspnea](#), bilateral basilar [rales](#)) would be expected and the intensity of the [murmur](#) would not increase on inspiration. This patient presents with features of right [heart failure](#) and a [murmur](#) that increases on inspiration.

A 54-year-old woman is brought to the emergency department because of a 2-hour history of nausea and retrosternal chest pain. She has a 15-year history of type 2 diabetes mellitus. Her current medications include atorvastatin, metformin, and lisinopril. She is diaphoretic. Her serum troponin level is 3.0 ng/mL (N: < 0.04). She undergoes cardiac catheterization. A photograph of coronary angiography performed prior to percutaneous coronary intervention is shown. An acute infarct associated with the finding on angiography is most likely to manifest with ST elevations in which of the following leads on ECG?



	Answer	Image
C	V ₁ –V ₆	The diagram illustrates the coronary artery system from two perspectives: the anterior (ventral) view and the diaphragmatic (dorsocaudal) view. The anterior view shows the Right coronary artery (RCA), Left coronary artery (LCA), Left circumflex artery (LCX), Obtuse marginal artery (OMA), Acute marginal artery (AMA), and Left anterior descending artery (LAD). The diaphragmatic view shows the Aorta, Right coronary artery (RCA), and Posterior descending artery (PDA). A legend indicates the regions supplied by each artery: RCA (blue), AMA (purple), LAD (green), LCX (orange), and PDA (pink).
D	II, III, and aVF	
E	V ₇ –V ₉	

Hint

This patient's coronary angiography study shows proximal narrowing of the left anterior descending artery. This artery perfuses the anterior two-thirds of the interventricular septum, the anterolateral papillary muscle, and the anterior surface of the left ventricle.

Correct Answer

A - I, aVR

Explanation Why

[ST elevations](#) in leads I and aVR are an atypical finding because these leads do not correspond with any specific territory of the [heart](#) that is supplied by a single [artery](#). This patient's [coronary angiography](#) study shows severe occlusion of the [left anterior descending artery](#).


B - V_{3R}–V_{6R}

Explanation Why

[ST elevations](#) in leads V_{3R}–V_{6R} on [ECG](#) typically develop as a result of [infarcts](#) that involve the [right coronary artery](#) and its branches, which supply the inferior aspect of the [heart](#) (e.g., [right ventricle](#)). This patient's [coronary angiography](#) study shows severe occlusion of the [proximal left anterior descending artery](#), which does not typically affect these leads.

C - V₁-V₆

Image



Explanation But

[Myocardial infarction](#) can also result in the development of pathologic Q waves on [ECG](#).

Explanation Why

[ST elevations](#) typically develop in leads V₁-V₆ on [ECG](#) as a result of [infarcts](#) that involve the [proximal left anterior descending artery](#) (LAD). Leads V₁-V₂ correspond to the anteroseptal territory; leads V₃-V₄ correspond to the anteroapical territory; leads V₅-V₆ correspond to the anterolateral territory. In some cases, [ST elevations](#) in the lateral leads (leads I and aVL) can also develop.

D - II, III, and aVF

Explanation Why

[ST elevations](#) typically develop in leads II, III, and aVF on [ECG](#) as a result of [infarcts](#) that involve the [right coronary artery](#), which supplies the inferior aspect of the [heart](#). This patient's [coronary angiography](#) study shows severe occlusion of the [left anterior descending artery](#), which does not typically affect these leads.

E - V7–V9

Explanation Why

[ST elevations](#) typically develop in leads V7–V9 on [ECG](#) as a result of [infarcts](#) that involve the [posterior descending artery](#), which supplies the [posterior](#) aspect of the [heart](#). This patient's [coronary angiography](#) study shows severe occlusion of the [left anterior descending artery](#), which does not typically affect these leads.

Question # 39

A 3-month-old boy is brought to the physician for a routine follow-up examination. He was delivered at 32 weeks' gestation to a 35-year-old woman. Cardiac examination is shown. Which of the following is the most likely cause for this patient's findings?

[Video](#)

	Answer	Image
A	Patent ductus arteriosus	
B	Ventricular septal defect	
C	Atrial septal defect	
D	Mitral valve prolapse	
E	Truncus arteriosus	
F	Tetralogy of Fallot	

Hint

Auscultation shows a continuous murmur. This finding is associated with a condition that affects infants who are born prematurely and/or have congenital rubella infection.

Correct Answer

A - Patent ductus arteriosus

Explanation Why

A [patent ductus arteriosus \(PDA\)](#) manifests with a continuous [machinery murmur](#) best heard in the left infraclavicular area. The [machinery murmur](#) is generated as blood continuously shunts from the high-pressure aorta through the [ductus arteriosus](#) into the low-pressure [pulmonary artery](#). It may envelop the S2 and is not usually affected by positional changes. Patients with moderate-sized PDAs may present with [bounding pulses](#), widened [pulse pressure](#), and symptoms of [heart failure](#).

B - Ventricular septal defect

Explanation Why

[Ventricular septal defects \(VSD\)](#) manifest with [holosystolic](#), harsh sounding [murmurs](#), best heard in the left lower sternal border. These [murmurs](#) often obliterate the S2 and are associated with a palpable thrill. [Neonates](#) present with an asymptomatic [heart murmur](#) or, in cases of moderate-to-large sized defects, with signs of heart failure. The examination findings in this patient reveal a continuous [machinery murmur](#) in the left upper sternal border, which is not suggestive of a [VSD](#).

C - Atrial septal defect

Explanation Why

[Atrial septal defects \(ASD\)](#) manifest with pulmonary outflow tract [murmurs](#) heard best at the left upper sternal border and are often associated with fixed [splitting of S2](#). ASDs are often low velocity shunts that produce insufficient flow to be audible. Larger shunts may cause [diastolic flow murmurs](#) across the [tricuspid valve](#) owing to a large left-to-right shunt. However, the continuous [machinery murmur](#) would not be seen in patients with an ASD.

D - Mitral valve prolapse

Explanation Why

[Mitral valve prolapse \(MVP\)](#) is associated with a late [systolic](#) crescendo [murmur](#) with a midsystolic click heard best at the apex. The examination findings of a continuous [machinery murmur](#) in the left upper sternal border, such as that observed in this patient, would not be seen in patients with [MVP](#).

E - Truncus arteriosus

Explanation Why

[Truncus arteriosus \(TA\)](#) results in a single conduit providing mixed venous and arterial blood to the coronary, pulmonary, and systemic circulations. Cardiac exam findings of TA would include a single loud S2 and the presence of an [ejection click](#) at the cardiac apex or left sternal border. Accentuated and bounding peripheral pulses might also be present. However, patients with an unrepaired defect present with harsh [systolic murmurs](#) at the left sternal border, unlike the continuous [machinery murmur](#) heard in this patient.

F - Tetralogy of Fallot

Explanation Why

[Tetralogy of Fallot \(TOF\)](#) is associated with a harsh crescendo-decrescendo [murmur](#) best heard along the left mid- to upper-sternal border. The [murmur](#) can decrease in intensity as the degree of [right ventricular outflow tract obstruction](#) worsens because of increased right-to-left shunting of blood. The examination findings of a continuous [machinery murmur](#) in the left upper sternal border, such as that observed in this patient, would not be seen in patients with TOF.

Question # 40

A 52-year-old man is brought to the emergency department for recurrent vomiting that began 5 hours ago. He describes the vomitus as bloody and containing black specks. He drinks 10–12 beers daily and has not seen a physician in 15 years. He appears pale. His temperature is 36.7°C (98.1°F), pulse is 122/min, and blood pressure is 85/59 mm Hg. Physical examination shows decreased radial pulses, delayed capillary refill, multiple spider angiomas, and a distended abdomen. He is oriented to person but not to place or time. Which of the following is most likely decreased in this patient in response to his hematemesis?

	Answer	Image
A	Systemic vascular resistance	
B	Capillary fluid absorption	
C	Fractional tissue oxygen extraction	
D	Carotid baroreceptor firing activity	The diagram illustrates the physiological response to decreased blood pressure and hypoxia. On the left, the baroreceptor reflex is shown: a decrease in blood pressure (↓ Blood pressure) leads to decreased parasympathetic activity (↓ Parasympathetic activity) and increased sympathetic activity (↑ Sympathetic activity). This results in vaso-dilation (Vaso-dilation) and increased heart rate (HR) and blood pressure (BP). On the right, the chemoreceptor reflex is shown: decreased pH, decreased pO₂, and increased PCO₂ lead to increased sympathetic activity (↑ Sympathetic activity). A legend identifies the components: 1. Vagus nerve, 2. Glossopharyngeal nerve, 3. Sympathetic nervous system, 4. Parasympathetic nervous system, 5. Solitary nucleus, 6. Carotid sinus, 7. Aortic arch, 8. Carotid body, 9. Aortic body, 10. SA node, 11. AV node.
E	Pulmonary vascular resistance	
F	Cardiac inotropy	

Hint

This patient's hematemesis, tachycardia, hypotension, weak pulses, and delayed capillary refill are concerning for hemorrhagic shock due to variceal bleeding.

Correct Answer

A - Systemic vascular resistance

Explanation Why

The physiologic responses to [hypovolemic shock](#) help to maintain tissue perfusion and oxygenation. Accordingly, [sympathetic](#) tone increases and results in increased, not decreased, [systemic vascular resistance](#). The peripheral vasculature constricts to help maintain blood pressure; this process is mediated by the [baroreceptor reflex](#) and chemoreceptor reflex, [catecholamine](#) release, and [renin-angiotensin-aldosterone system \(RAAS\)](#) activation.

B - Capillary fluid absorption

Explanation Why

[Capillary](#) fluid absorption is increased, not decreased, in response to [hypovolemic shock](#). [Starling forces](#) govern [capillary](#) fluid movement, which is mainly a balance between the [hydrostatic pressure](#) of blood in the [capillaries](#) (which pushes intravascular fluid out) and the [oncotic pressure](#) gradient across the [capillary](#) wall (which keeps intravascular fluid within blood vessels). [Interstitial](#) hydrostatic and oncotic pressures are typically much lower than the pressures in the [capillaries](#), and, therefore, do not have a significant influence on [capillary](#) fluid movement. In this patient with hemorrhagic shock, decreased blood pressure results in decreased intravascular [hydrostatic pressure](#) and, therefore, increased [capillary](#) fluid absorption.

C - Fractional tissue oxygen extraction


Explanation Why

The physiologic responses to [hypovolemic shock](#) help to maintain tissue perfusion and oxygenation. Accordingly, fractional tissue oxygenation extraction is increased, not decreased, to compensate for decreased [oxygen delivery](#) to peripheral tissues from a loss of [red blood cells](#) and intravascular volume. Oxygen that is bound to [hemoglobin](#) is more readily released to peripheral tissues (i.e., [right shift of the oxygen-hemoglobin dissociation curve](#)). As a result, [mixed venous oxygen saturation](#)

decreases.

D - Carotid baroreceptor firing activity

Image



Explanation But

[Baroreceptors](#) are only able to respond to short-term changes in blood pressure because their firing frequency adapts to a new blood pressure level within a few days.

Explanation Why

[Baroreceptors](#) are mechanoreceptors that fire in response to blood vessel stretching to regulate autonomic activity. Carotid [baroreceptors](#) send afferent signals via the [glossopharyngeal nerve](#) to the [solitary nucleus](#) in the medulla. In response, the [brainstem](#) modulates autonomic outflow to alter the [heart rate](#) and [peripheral vascular resistance](#) to normalize blood pressure (i.e., the [baroreceptor reflex](#)). In hypotensive states, decreased arterial pressure results in decreased stretching of the arterial wall and decreased carotid (and aortic) [baroreceptor](#) firing activity. The resulting [sympathetic](#) stimulation and [parasympathetic](#) inhibition lead to compensatory increases in [heart rate](#), cardiac contractility, vasoconstriction, and blood pressure to maintain tissue perfusion and oxygenation.

E - Pulmonary vascular resistance

Explanation Why

The physiologic responses to [hypovolemic shock](#) help to maintain tissue perfusion and oxygenation. Accordingly, [sympathetic](#) tone increases and results in increased, not decreased, [pulmonary vascular resistance](#). Additional pulmonary responses to [hypovolemic shock](#) include an increased [respiratory rate](#) (as a compensatory response to [lactic acidosis](#)) and increased alveolar-[capillary](#) permeability.

F - Cardiac inotropy

Explanation Why

The physiologic responses to [hypovolemic shock](#) help to maintain tissue perfusion and oxygenation. Accordingly, [sympathetic](#) tone increases and results in increased, not decreased, cardiac [inotropy](#). Blood loss results in [hypovolemia](#), reduced cardiac filling, and decreased [stroke volume](#). Cardiac contractility increases in response to increased [sympathetic](#) tone mediated by the [baroreceptor reflex](#) and chemoreceptor reflex, [catecholamine](#) release, and [renin-angiotensin-aldosterone system](#) (RAAS) activation.

Question # 1

A 25-year old man comes to the physician because of fatigue over the past 6 months. He has been to the emergency room several times over the past 3 years for recurrent shoulder and patella dislocations. Physical examination shows abnormal joint hypermobility and skin hyperextensibility. A high-frequency mid-systolic click is heard on auscultation. Which of the following is most likely to result in an earlier onset of this patient’s auscultation finding?

	Answer	Image
A	Hand grip	
B	Abrupt standing	
C	Rapid squatting	
D	Valsalva release phase	
E	Leaning forward	

Hint

This patient most likely has Ehlers-Danlos syndrome, which can manifest with joint hypermobility (predisposing to joint dislocations) and hyperextensible skin. Ehlers-Danlos syndrome is associated with mitral valve prolapse, a structural defect of the mitral valves due to myxomatous degeneration that leads to bulging of the leaflets into the left atrium during systole.

Correct Answer


A - Hand grip

Explanation Why

The characteristic click heard on auscultation in patients with [mitral valve prolapse \(MVP\)](#) results from the bulging of the valve leaflets into the [atrium](#), which begins when the contracting [left ventricle](#) reaches a critical volume during [systole](#). An increased [left ventricular end-diastolic volume](#), e.g., due to increased [afterload](#) and subsequent increased [left ventricular end-systolic volume](#) during hand grip (hand clenching) delays the point when that volume is reached and results in a later onset of the auscultatory findings.

B - Abrupt standing

Image



Explanation Why

The characteristic auscultatory finding of [mitral valve prolapse \(MVP\)](#) is a high-frequency mid-[systolic](#) click, which is caused by sudden tensing of the [chordae tendineae](#) as the valve leaflets prolapse. The click may be followed by a mid-to-late [systolic murmur](#) indicating [mitral regurgitation](#). Since the prolapse occurs when the contracting ventricle reaches a critical volume, the timing of the click varies according to the [left ventricular end-diastolic volume \(EDV\)](#). Decreased [EDV](#), e.g., due to decreased [venous return](#) to the [heart](#) during abrupt standing (i.e., decreased [preload](#)) causes the leaflets to prolapse sooner, resulting in an earlier onset (i.e., closer to S1) and an increased intensity of auscultatory findings. Conversely, maneuvers that increase the volume (e.g., squatting) result in a later onset and decreased intensity.

C - Rapid squatting

Explanation Why

The characteristic click heard on auscultation in patients with [mitral valve prolapse \(MVP\)](#) results from the bulging of the valve leaflets into the [atrium](#), which begins when the contracting [left ventricle](#) reaches a critical volume during [systole](#). An increased [left ventricular end-diastolic volume](#), e.g., due to increased [venous return](#) to the [heart](#) during rapid squatting (i.e., increased [preload](#)) delays the point when that volume is reached and results in a later onset of the auscultatory findings.

D - Valsalva release phase

Explanation Why

The characteristic click heard on auscultation in patients with [mitral valve prolapse \(MVP\)](#) results from the bulging of the valve leaflets into the [atrium](#), which begins when the contracting [left ventricle](#) reaches a critical volume during [systole](#). An increased [left ventricular end-diastolic volume](#), e.g., due to increased [venous return](#) to the [heart](#) during the release phase of the [Valsalva maneuver](#) (i.e., increased [preload](#)) delays the point when that volume is reached and results in a later onset of the auscultatory findings.

E - Leaning forward

Explanation Why

Leaning forward brings the [heart](#) closer to the chest, but it has no significant impact on [preload](#) or [afterload](#), so it does not affect the timing of the [murmur](#) of [mitral valve prolapse](#). However, a [pericardial friction rub](#) and the [murmur](#) of [aortic regurgitation](#) are best heard with the patient leaning forward.

Question # 2

A 3-year-old girl is brought to the emergency department because of chest pain for 2 hours. Eight days ago, she was admitted to the hospital for treatment of low-grade fever, malaise, and sore throat. The hospitalization was complicated by pharyngitis with pseudomembrane formation and severe cervical lymphadenopathy briefly requiring intubation. She has not received any routine childhood vaccinations. Serum studies show elevated cardiac troponins. An ECG shows diffuse T wave inversions and prolonged PR interval. Administration of which of the following at the time of her previous admission is most likely to have prevented this patient's cardiac symptoms?

	Answer	Image
A	Denatured bacterial toxin that contains an intact receptor binding site	
B	Electrolyte that reduces cardiomyocyte excitability	
C	Antibiotic that binds to penicillin-binding protein 3	
D	Salicylate that inhibits prostaglandin synthesis	
E	Immunoglobulin that targets circulating proteins	

Hint

Pharyngitis with pseudomembrane formation and cervical lymphadenopathy (“bull neck”) is suggestive of diphtheria. The patient's signs and symptoms of cardiac dysfunction (chest pain, elevated cardiac troponins, ECG abnormalities) are indicative of myocarditis, a concerning complication of diphtheria.

Correct Answer

A - Denatured bacterial toxin that contains an intact receptor binding site

Explanation Why

Administration of a denatured [bacterial toxin](#) that contains an intact receptor binding site (e.g., [diphtheria](#) toxoid) provides [active immunity](#). However, [active immunity](#) takes several weeks to develop; therefore, administration of the [diphtheria vaccine](#) would not have helped at the time of admission. Previous [vaccination](#) certainly would have dramatically reduced the [odds](#) of contracting [diphtheria](#).

B - Electrolyte that reduces cardiomyocyte excitability

Explanation Why

Calcium is an electrolyte that reduces [cardiomyocyte](#) excitability and is typically administered to protect the [myocardium](#) from the deleterious effects of [hyperkalemia](#). Although [hyperkalemia](#) can present with malaise and prolonged [PR interval](#), as seen in this patient, elevated cardiac [troponins](#), diffuse [T wave](#) inversions, and [chest pain](#) are not common. Furthermore, this patient's presentation is strongly suggestive of [diphtheria](#), for which calcium is not indicated.

C - Antibiotic that binds to penicillin-binding protein 3

Explanation Why

[Monobactam antibiotics](#) (e.g., [aztreonam](#)) have intrinsic [β-lactamase](#) activity (which binds [PBP](#)) but are effective against gram-negative [bacilli](#) only. [C. diphtheriae](#) is a gram-positive rod, so it is not susceptible to [monobactam antibiotics](#). Appropriate [antibiotics](#), such as [penicillin](#) (or other [beta-lactam antibiotics](#)), should be given to patients with [diphtheria](#) to decrease the transmissibility and improve the course of the acute infection.

D - Salicylate that inhibits prostaglandin synthesis

Explanation Why

[Aspirin](#) is a [salicylate](#) that works by inhibiting [prostaglandin](#) synthesis. It can be used in the treatment of [Kawasaki disease](#) to decrease the risk of [coronary artery aneurysms](#). Although [Kawasaki disease](#) can also present with [fever](#), cervical [lymphadenopathy](#), [chest pain](#), and elevated cardiac [troponins](#) (as a result of [myocardial ischemia](#)), this patient lacks other typical features, such as bilateral nonexudative [conjunctivitis](#), mucositis, and rash. Furthermore, the presence of [pharyngitis](#) with pseudomembrane formation is highly suggestive of [diphtheria](#).


E - Immunoglobulin that targets circulating proteins

Explanation Why


Immediate administration of [immunoglobulin](#) that targets circulating [proteins](#) (i.e., [diphtheria antitoxin](#)) provides [passive immunity](#) and is a critical part of [diphtheria](#) treatment. The use of an antitoxin is imperative as the clinical features of [diphtheria](#) are not caused by the pathogen itself but rather by the [exotoxin](#) ([diphtheria toxin](#)) that [C. diphtheriae](#) produces. The antitoxin can only neutralize unbound toxin and should therefore be administered early in the course of the disease.

Question # 3

A 6-year-old boy is brought to the physician by his mother because of a 3-month history of episodic chest pain and shortness of breath on exertion. He is at the 99th percentile for height and 40th percentile for weight. Examination shows a high-arched palate, long and slender upper extremities, and elbows and knees that can be hyperextended. Cardiac examination shows a grade 2/6 late systolic, crescendo murmur with a midsystolic click. Over which of the following labeled areas is the murmur most likely to be heard best?



	Answer	Image
A	Area A	
B	Area B	
C	Area C	

	Answer	Image
D	Area D	
E	Area E	
F	Area F	
G	Area G	

Hint

This boy's tall stature, long and slender arms, joint hypermobility (as evidenced by his hyperextensible elbows and knees), and high-arched palate are consistent with Marfan syndrome, an inherited connective tissue disorder associated with ocular, musculoskeletal, and cardiovascular abnormalities.

Correct Answer

A - Area A

Explanation Why

Area A corresponds to the carotid area. [Aortic stenosis](#) causes a midsystolic, crescendo-decrescendo [murmur](#) that can radiate to this area. Severe [carotid artery stenosis](#) can also cause a [carotid bruit](#) heard over this area. The [murmur](#) most commonly associated with [Marfan syndrome](#) is best heard over a different area.

B - Area B

Explanation Why

Area B corresponds to the [aortic area](#), which is located in the right 2nd intercostal space in the parasternal line. [Aortic stenosis](#) causes a midsystolic, crescendo-decrescendo [murmur](#) best heard over this area. [Marfan syndrome](#), which this patient has, commonly causes [aortic root dilation](#) and, consequently, [aortic regurgitation](#). In [aortic regurgitation](#) due to aortic root disease, an early [diastolic murmur](#) is heard over the [aortic area](#) during [diastole](#). The [systolic murmur](#) in this patient, however, is best heard over a different area.

C - Area C

Explanation Why

Area C corresponds to the [pulmonic area](#), located in the left 2nd parasternal [intercostal space](#). [Pulmonary stenosis](#) causes a [systolic](#), crescendo-decrescendo [murmur](#) best heard over this area. [Pulmonary stenosis](#) is not associated with [Marfan syndrome](#).

D - Area D

Explanation Why

Area D corresponds to the left sternal border. [Aortic regurgitation](#) (AR) due to valvular disease can cause a [murmur](#) that is best heard over this area. AR in [Marfan](#) disease, although a common manifestation, is due to [aortic root dilation](#), which causes a [murmur](#) best heard at the right rather than the left sternal border. Moreover, the [murmur](#) in AR is early [diastolic](#). The [systolic murmur](#) in this patient is best heard over a different area.


E - Area E

Explanation Why

Area E corresponds to the [tricuspid area](#) located in the left 4th parasternal [intercostal space](#). [Ventricular septal defects](#) (the most common congenital cardiac defect) can cause a [holosystolic](#), harsh-sounding [murmur](#) best heard over this area. Tricuspid stenosis causes a [diastolic](#), low-pitched, decrescendo [murmur](#) and [tricuspid regurgitation](#) causes a [holosystolic murmur](#), both of which can also be heard over this area. The [murmur](#) most commonly associated with [Marfan syndrome](#) is best heard over a different area.

F - Area F

Image



Explanation But

[Marfan syndrome](#) can lead to other musculoskeletal abnormalities, including [arachnodactyly](#) and [spinal](#) abnormalities ([scoliosis](#) and [sternal kyphosis](#)). Ocular manifestations can include [ectopia lentis](#) (superiorly and temporally), [myopia](#), [retinal detachment](#), and [glaucoma](#).

Explanation Why

Patients with [Marfan syndrome](#) can have [mitral valve prolapse \(MVP\)](#), which causes a late [systolic](#), crescendo [murmur](#) with a midsystolic click. This [murmur](#) is best heard at the [heart](#) apex ([mitral area](#), located at the left 5th intercostal space in the [midclavicular line](#)) when the patient is in the left [lateral](#) recumbent position. Other cardiovascular abnormalities associated with [Marfan syndrome](#) include aortic disease (e.g., [aortic root dilation](#), [aortic regurgitation](#), [aortic aneurysm](#), [aortic dissection](#)) and [berry aneurysms](#).


G - Area G

Explanation Why

Area G corresponds to the left [midaxillary line](#). [Mitral regurgitation](#) causes a [holosystolic murmur](#) that can radiate to this area, particularly if the patient is in the left [lateral](#) recumbent position. The [murmur](#) most commonly found in patients with [Marfan syndrome](#) is best heard over a different area.

Question # 4

A 75-year-old woman is brought to the emergency department 30 minutes after an episode of dizziness and loss of consciousness that lasted 10 seconds. The symptoms occurred while the patient was exercising in an aerobics class. She has a 3-year history of progressive shortness of breath and chest discomfort on exertion. Radial pulses are decreased and delayed bilaterally. The patient most likely has a murmur starting at which of the following labeled points in the curve shown?



	Answer	Image
A	A	
B	B	
C	C	

	Answer	Image
D	D	
E	E	

Hint

The patient has a history of exertional dyspnea and angina pectoris and presents with exertional syncope. Examination shows pulsus parvus et tardus. These signs and symptoms are most consistent with aortic valve stenosis.

Correct Answer

A - A

Explanation Why

This point on the volume/pressure curve corresponds to the beginning of the [left ventricular](#) contraction and the closure of the [mitral valve](#). During this period, the ventricular pressure rises without changes in volume ([isovolumetric contraction](#)). Hence the [aortic valve](#) remains closed until the pressure between aorta and ventricle is equalized. Since there is no blood flowing through the valve, [aortic stenosis murmur](#) would not be heard at this point.

B - B

Explanation Why

This point on the [left ventricular](#) volume/pressure curve corresponds to the opening of the [aortic valve](#). Patients with [aortic valve stenosis](#) (AS) typically present with a crescendo-decrescendo, [systolic](#), ejection [murmur](#) that begins at this point (shortly after the S1) and finishes before the S2.

C - C

Explanation Why

This point indicates the beginning of [diastole](#) and corresponds to the closure of the [aortic valve](#). When the pressure inside the ventricle falls below the aortic pressure, the [aortic valve](#) closes and a period of [isovolumetric relaxation](#) occurs. The [murmur](#) associated with [aortic stenosis](#) is produced during [systole](#), and would therefore not be found at this point.

D - D

Explanation Why

This point corresponds to the opening of the [mitral valve](#) and indicates the beginning of the passive [filling phase](#) of the ventricle. The [murmur](#) encountered in this patient would occur during [systole](#), as the blood flows from the [left ventricle](#) to the aorta, but not during [diastole](#) while the ventricle is being passively filled.

E - E

Explanation Why

This point represents the left atrial contraction and the [left ventricular](#) rapid [filling phase](#). While the [left ventricle](#) continues to relax during [diastole](#), the left [atria](#) forces more blood flow across the [mitral valve](#). [Mitral valve stenosis murmurs](#) rather than [aortic stenosis murmurs](#) would be heard at this point. [Mitral valve stenosis](#) is associated with [dyspnea](#) on exertion but less commonly with [angina pectoris](#). It is not associated with exertional [dizziness](#), [syncope](#), or [pulsus parvus et tardus](#).

Question # 5

A 62-year-old woman with a history of hypertension and type 2 diabetes mellitus comes to the physician for a routine health maintenance examination. She has smoked 1 pack of cigarettes daily for the last 15 years. Current medications include glyburide and amlodipine. The physician prescribes a medication that decreases the production of mevalonate. Which of the following changes to the serum is most likely to develop as an adverse effect of the prescribed drug?

	Answer	Image
A	Increased creatine kinase concentration	
B	Decreased glucose concentration	
C	Increased lactate concentration	
D	Increased triglyceride concentration	
E	Increased bradykinin concentration	

Hint

This woman has been prescribed a statin as part of a primary prevention strategy for hyperlipidemia. Statins reduce serum cholesterol concentrations by preventing the conversion of HMG-CoA to mevalonate, which is a precursor of cholesterol.

Correct Answer

A - Increased creatine kinase concentration

Explanation But

Strategies to prevent muscle-related adverse effects related to [statin](#) therapy include the use of the lowest effective dose to reach target lipid goals and avoidance of drugs that can increase the risk of developing myopathies, such as [fibrates](#), [niacin](#), [cyclosporine](#), and [glucocorticoids](#). Muscle-related adverse effects typically resolve upon discontinuation of the drug.

Explanation Why

Muscle-related adverse effects can develop within weeks to months after starting [statin](#) therapy. [Statins](#) can decrease the synthesis of coenzyme Q10 and impair energy production within muscle, which can result in myalgia, muscle weakness, and increased serum [creatinine kinase](#) (CK) concentrations due to [myositis](#) and/or [rhabdomyolysis](#).

B - Decreased glucose concentration

Explanation Why

This patient takes [glyburide](#), a [sulfonylurea](#) that can cause [hypoglycemia](#). However, [statin](#) therapy does usually not affect serum glucose concentrations.

C - Increased lactate concentration

Explanation Why

[Statin](#) therapy does usually not affect serum [lactate](#) concentrations. [Lactic acidosis](#) is a rare but serious adverse effect of [metformin](#) therapy.

D - Increased triglyceride concentration

Explanation Why

[Statin](#) therapy lowers serum [triglyceride](#) concentrations by 10–20%. Drugs that commonly increase serum [triglyceride](#) concentrations include [beta blockers](#), [glucocorticoids](#), [cyclosporine](#), and some HIV antiretroviral drugs.

E - Increased bradykinin concentration

Explanation Why

[Statin](#) therapy does usually not affect serum [bradykinin](#) concentrations. The kinin pathway is altered by ACE inhibitors, which increase serum [bradykinin](#) concentrations and can result in adverse effects such as [cough](#) and [angioedema](#).

Question # 6

Seventy-two hours after admission for an acute myocardial infarction, a 48-year-old man develops dyspnea and a productive cough with frothy sputum. Physical examination shows coarse crackles in both lungs and a blowing, holosystolic murmur heard best at the apex. ECG shows Q waves in the anteroseptal leads. Pulmonary capillary wedge pressure is 23 mm Hg. Which of the following is the most likely cause of this patient's current condition?

	Answer	Image
A	Postmyocardial infarction syndrome	
B	Aortic root dilation	
C	Rupture of the chordae tendinae	
D	Rupture of the ventricular free wall	
E	Rupture of the interventricular septum	

Hint

This patient's new murmur (blowing, holosystolic murmur best heard at the apex) and evidence of heart failure (dyspnea, bilateral crackles, elevated capillary wedge pressure) suggest acute mitral regurgitation.

Correct Answer

A - Postmyocardial infarction syndrome

Explanation Why

[Postmyocardial infarction syndrome](#) is a type of [pericarditis](#) that arises from an autoimmune reaction to exposed [myocardial](#) antigens after acute [MI](#). Patients present with [dyspnea](#), [tachypnea](#), and [tachycardia](#). However, this syndrome would manifest 2–10 weeks after an [MI](#) rather than 72 hours as seen in this patient. Furthermore, [fever](#) is usually present, [cardiac auscultation](#) would reveal a [pericardial friction rub](#), and [ECG](#) would show [ST elevations](#) in all leads.

B - Aortic root dilation

Explanation Why

[Aortic root dilation](#) can occur in [postinfarction fibrinous pericarditis](#), which commonly manifests 1–3 days after [MI](#). Patients with [aortic root dilation](#) present with features of [aortic regurgitation](#). These patients have [diastolic](#), not [systolic](#), [murmurs](#). Moreover, this patient is missing other symptoms of [pericarditis](#), such as [chest pain](#) and a friction rub.

C - Rupture of the chordae tendinae

Explanation But

The posteromedial [papillary muscle](#) is supplied only by the [posterior descending artery](#) and is at especially high risk of [ischemia](#) due to lack of collateral supply.

Explanation Why

Chordae tendinae or [papillary muscle rupture](#) most commonly occurs 2–7 days after [acute myocardial infarction](#) and is due to extensive [coagulative necrosis](#) and the formation of [granulation tissue](#) at the area of [infarction](#). This patient's localization of Q waves in the anteroseptal leads (e.g.,

V1–V4) suggests an [infarction](#) of the [left anterior descending artery](#), which supplies the anterolateral [papillary muscle](#). Acute [mitral regurgitation](#) can lead to signs of heart failure, as seen in this patient.

D - Rupture of the ventricular free wall

Explanation Why

Ventricular wall rupture can occur within 2 weeks of [myocardial infarction](#) and typically results in [hemopericardium](#) and [cardiac tamponade](#), which can manifest with profound [right heart failure](#), [shock](#), and often death. While free wall rupture and [cardiac tamponade](#) can also cause acute [pulmonary edema](#), additional expected findings would include [pulsus paradoxus](#) and muffled [heart sounds](#), none of which is observed in this patient.

E - Rupture of the interventricular septum




Explanation Why

Rupture of the interventricular septum may occur within the first 2 weeks after [MI](#) and typically presents with sudden hemodynamic instability ([hypotension](#), [tachycardia](#)), [dyspnea](#), and new-onset [holosystolic murmur](#). However, because blood is shunted from left to right, patients typically present with signs of [right heart failure](#) (e.g., increased [JVP](#), peripheral [edema](#)) rather than the signs of isolated [left heart failure](#) (e.g., [pulmonary edema](#), elevated [PCWP](#)) seen here. Moreover, the expected [murmur](#) would be heard best over the left sternal border, not the apex.

Question # 7

A 2-week-old newborn is brought to the physician because of worsening feeding difficulty since birth. Examination shows a grade 2/6 harsh holosystolic murmur, heard most clearly at the left lower sternal border, and a soft mid-diastolic rumble over the cardiac apex. Echocardiography shows shunting of blood through the ventricular septum during systole. The patient undergoes surgery for closure of the defect. Which of the following sets of changes are expected after successful repair of this cardiac defect?

	Left atrial pressure	Left ventricular pressure	Right ventricular pressure
A	↓	↑	no change
B	↓	↑	↓
C	↑	↑	↑
D	↓	↓	↓
E	↑	↑	↓

	Answer	Image
A	A	
B	B	1  2  3 
C	C	

	Answer	Image
D	D	
E	E	

Hint

The newborn presented with symptoms of decreased cardiac output (poor feeding) and heart murmurs; he was found to have a ventricular septal defect, which allowed for abnormal shunting of blood from the left to the right ventricle. Successful repair of this defect will restore the normal cardiac output.

Correct Answer


A - A

Explanation Why

Mitral stenosis (MS) causes a left atrial outflow obstruction. The left atrium has to contract against the partially closed mitral valve, which results in an increased left atrial pressure. Once repaired, left atrial outflow increases, thereby decreasing the left atrial pressure. The left ventricular input increases, thereby increasing the left ventricular pressure. The right ventricular pressure remains unchanged. These hemodynamic changes would not, however, be expected after the closure of a VSD.

B - B

Image



Explanation Why

A [VSD](#) causes [left-to-right shunting](#) of blood because the [left ventricular](#) pressure is physiologically higher than the [right ventricular](#) pressure due to the thicker ventricular wall and higher force of contraction. Closure of a [VSD](#) would, therefore, increase the amount of blood in the [left ventricle](#) and decrease the amount of blood in the [right ventricle](#) during each [cardiac cycle](#). This results in an increase in [left ventricular](#) pressure and a decrease in [right ventricular](#) pressure as well as [right ventricular](#) output, which translates to a decrease in the amount of blood flowing into the [left atrium](#) and therefore a decrease in left atrial pressure.

C - C

Explanation Why

An increase in the pressure of all the chambers of the [heart](#) would be expected following [fluid resuscitation](#) in patients with [hypovolemic shock](#). These hemodynamic changes would not be expected after the closure of a [VSD](#).

D - D

Explanation Why

A decrease in the pressure in all the chambers of the [heart](#) would be expected following [pericardiectomy](#) for [constrictive pericarditis](#) or following [aspiration](#) of a [hemopericardium](#). These hemodynamic changes would not be expected after the closure of a [VSD](#).

E - E

Explanation Why

Both [atrial septal defect](#) repair, which stops shunting of blood from the [left atrium](#) to the [right atrium](#), and [pulmonary stenosis](#) repair, which removes the right ventricular outflow obstruction, would decrease [right ventricular](#) pressure and increase left atrial as well as [left ventricular](#) pressure by

increasing the amount of blood in the left side of the [heart](#). These hemodynamic changes would not, however, be expected after the closure of a [VSD](#).

Question # 8

A 73-year-old woman comes to the physician because of recurrent episodes of losing consciousness for several seconds upon standing. She has a history of hypertension, which has been treated with hydrochlorothiazide. Her blood pressure is 130/87 mm Hg in the supine position and 100/76 mm Hg 30 seconds after standing up. Cardiac examination shows no abnormalities. Which of the following sets of changes is most likely to occur when the patient stands up?

	Venous return	Carotid sinus baroreceptor activity	Cerebral blood flow
A	↓	↓	↓
B	↑	↑	↑
C	No change	↓	↓
D	↑	↑	↓
E	↓	↑	↓

	Answer	Image
A	A	
B	B	
C	C	
D	D	
E	E	

Hint

Fainting upon standing, combined with a history of advanced age, diuretic use, and the patient's orthostatic blood pressure findings, are best explained by orthostatic hypotension.

Correct Answer

A - A

Explanation Why

[Venous return](#) is diminished upon standing because of blood pooling in the lower extremities. Diminished [venous return](#), in conjunction with reduced blood volumes secondary to [hydrochlorothiazide](#) use, results in reduced arterial blood volumes. In turn, reduced arterial blood volumes decrease both [carotid sinus baroreceptor](#) activity and cerebral blood flow and cause [orthostatic syncope](#). [Autonomic dysfunction](#) often accompanies advanced age, which places elderly patients at increased risk of medication side effects, especially those involving the cardiovascular and [autonomic nervous systems](#) (e.g., α -1 receptor antagonists, [tricyclic antidepressants](#)).

B - B

Explanation Why

An increased cerebral blood flow, [carotid sinus baroreceptor](#) activity, and [venous return](#) are consistent with the physiological changes that a young athlete undergoes during exercise. This constellation would not result in [syncope](#) nor explain this patient's difference in blood pressure upon standing.

C - C

Explanation Why

A decreased cerebral blood flow and [carotid sinus baroreceptor](#) activity, with an unaltered [venous return](#), is consistent with [low blood pressure](#) as a consequence of [LV dysfunction](#). This constellation can be caused by [aortic stenosis](#), which often causes [syncope](#) as a result of reduced [cardiac output](#) and forward [heart failure](#), especially during times of exertion (e.g., such as rising from a chair). However, this patient has no [murmurs](#) to indicate [valvular heart disease](#).

D - D

Explanation Why

A decreased cerebral blood flow, with an increased [carotid sinus baroreceptor](#) activity and [venous return](#), is consistent with a reduction in cerebral perfusion secondary to severe bilateral [carotid stenosis](#). Although bilateral [carotid stenosis](#) can potentially cause [syncope](#), it would be very rare, especially without other previous common signs such as [amaurosis fugax](#) or [transient ischemic attacks/strokes](#).

E - E

Explanation Why

A decreased [venous return](#) and cerebral blood flow, with increased [carotid sinus baroreceptor](#) activity, is consistent with the activation of carotid [baroreceptors](#), which is caused by [carotid sinus massage](#). However, given the patient's history, this is unlikely to be the cause of her [syncope](#).

Question # 9

An 81-year-old man is admitted to the hospital due to acute decompensated heart failure. He has type 2 diabetes mellitus, hypertension, coronary artery disease, and congestive heart failure. Current medications include lisinopril, metformin, and low-dose aspirin. He has smoked one pack of cigarettes daily for 45 years. His temperature is 37.6°C (99.7°F), pulse is 105/min and regular, respirations are 21/min, and blood pressure is 103/64 mm Hg. Laboratory studies show:

Hemoglobin	13.7 g/dL
Leukocyte count	8200/mm ³
Serum	
Na ⁺	128 mEq/L
Cl ⁻	98 mEq/L
K ⁺	4.9 mEq/L
Urea nitrogen	58 mg/dL
Glucose	200 mg/dL
Creatinine	2.2 mg/dL

Which of the following changes in the medication regimen is most appropriate in this patient at this time?

	Answer	Image
A	Begin hydrochlorothiazide therapy	
B	Begin vancomycin therapy	

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

	Answer	Image
C	Discontinue aspirin therapy	
D	Discontinue metformin therapy	
E	Begin nitroprusside therapy	
F	Begin norepinephrine therapy	

Hint

This patient's laboratory findings indicate prerenal acute kidney injury (BUN:creatinine ratio $> 20:1$) secondary to decompensated heart failure (cardiorenal syndrome).

Correct Answer

A - Begin hydrochlorothiazide therapy

Explanation Why

[Diuretics](#) are used to treat fluid congestion in patients with acute [decompensated heart failure](#). However, [loop diuretics](#) are the first-line therapy while [thiazide diuretics](#) are used as an add-on if [loop diuretics](#) fail to relieve congestion.

B - Begin vancomycin therapy

Explanation Why

[Vancomycin](#) is commonly used in the [empiric treatment](#) of suspected [MRSA](#) infections. However, in the absence of [fever](#) and [leukocytosis](#), this patient's respiratory symptoms are more consistent with [pulmonary edema](#) secondary to acutely [decompensated heart failure](#) than an infectious etiology.

C - Discontinue aspirin therapy

Explanation Why

This patient takes [aspirin](#) for his [coronary artery disease](#), which requires low-dose [aspirin](#). While [NSAIDs](#) and high-dose [aspirin](#) are potentially [nephrotoxic](#) agents that should be avoided in patients with [AKI](#), low-dose [aspirin](#) does not impair renal perfusion and should therefore be continued in this patient to prevent complications of his illness.

D - Discontinue metformin therapy

Explanation But

The use of [metformin](#) may be safe in a case of chronic [heart failure](#) with preserved [cardiac output](#). Other important indications for discontinuing [metformin](#) (albeit temporarily) are surgery, [acidemia](#), and the intravenous administration of an iodinated contrast medium (e.g., [CT scans](#), fluoroscopic studies).

Explanation Why

[Metformin](#) is associated with an increased risk of life-threatening [lactic acidosis](#) in patients with predisposing conditions that reduce [lactic acid](#) clearance (i.e., renal and/or hepatic insufficiency), increased [lactic acid](#) production (e.g., [sepsis](#)), or a combination of both (e.g., [heart failure](#) with low [cardiac output](#)). This patient's [renal insufficiency](#) and [heart failure](#) with decreased [cardiac output](#) are both contraindications to [metformin](#) therapy, which should therefore be discontinued immediately. [Insulin](#) should instead be used for inpatient glycemic control.

E - Begin nitroprusside therapy

Explanation Why

In a patient with [acute decompensated heart failure](#), the use of [vasodilatory](#) medications such as [nitroprusside](#) can be considered as an adjunct to [diuretic](#) therapy for [dyspnea](#) relief. However, [nitroprusside](#) should not be used in patients with [hypotension](#).

F - Begin norepinephrine therapy

Explanation Why

[Inotropic](#) agents should be considered in a patient with acute [decompensated heart failure](#) and [hypotension](#). However, [dopamine](#) is the [inotropic](#) drug of choice in patients with [heart failure](#). The use of [norepinephrine](#) should only be considered in patients with a [systolic](#) blood pressure < 70 mm

Hg.

Question # 10

A 3175-g (7-lb) male newborn is delivered at 39 weeks' gestation to a 29-year-old primigravid woman following a spontaneous vaginal delivery. Apgar scores are 8 and 9 at 1 and 5 minutes, respectively. Cardiac examination in the delivery room shows a continuous machine-like murmur. An echocardiogram shows a structure with blood flow between the pulmonary artery and the aorta. This structure is most likely a derivate of which of the following?

	Answer	Image
A	4 th aortic arch	
B	2 nd aortic arch	
C	6 th aortic arch	
D	1 st aortic arch	
E	3 rd aortic arch	

Hint

This newborn's examination shows a structure between the pulmonary artery and aorta with a continuous “machine-like” murmur heard on cardiac auscultation. This is characteristic of a patent ductus arteriosus.

Correct Answer

A - 4th aortic arch

Explanation Why

The 4th [aortic arch](#) develops into the [aortic arch](#) on the left and the [proximal](#) part of the [right subclavian artery](#) on the right. The [ductus arteriosus](#) does not develop from the 4th aortic arch.


B - 2nd aortic arch

Explanation Why

The 2nd [aortic arch](#) develops into the stapedial [artery](#) and the hyoid [artery](#). The [ductus arteriosus](#) does not develop from the 2nd aortic arch.

C - 6th aortic arch

Image



Explanation Why

The 6th [aortic arch](#) is the origin of the [proximal](#) parts of the left and right [pulmonary arteries](#) and the [ductus arteriosus](#) (on the left side only).

D - 1st aortic arch

Explanation Why

The 1st [aortic arch](#) is the origin of the maxillary [artery](#). The [ductus arteriosus](#) does not develop from the 1st aortic arch.


E - 3rd aortic arch

Explanation Why

The 3rd [aortic arch](#) develops into the [common carotid artery](#) and [proximal](#) part of the [internal carotid artery](#). The [ductus arteriosus](#) does not develop from the 3rd aortic arch.

Question # 11

A 75-year-old with hypertension and atrial fibrillation comes to the emergency department because of a 2-hour history of severe abdominal pain and nausea. He has smoked 1 pack of cigarettes daily for the past 45 years. The patient undergoes emergency laparotomy and is found to have dusky discoloration of the hepatic colonic flexure and an adjacent segment of the transverse colon. The most likely cause of his condition is occlusion of a branch of which of the following arteries?

	Answer	Image
A	Median sacral artery	
B	Inferior mesenteric artery	
C	Celiac artery	
D	Internal iliac artery	

	Answer	Image
E	Superior mesenteric artery	

Hint

Approximately 50% of acute mesenteric ischemia cases are caused by acute arterial embolism as a result of cardiac pathologies (e.g., atrial fibrillation with development of atrial thrombi, myocardial infarction, valvular heart disease, endocarditis). The artery in question emerges from the abdominal aorta at the L1 vertebral level, posterior to the neck of the pancreas and pylorus of the stomach.

Correct Answer

A - Median sacral artery

Explanation Why

The [median sacral artery](#) branches off from the [abdominal aorta](#) at the L4 [vertebral](#) level, superior to the [aortic bifurcation](#) into the left and right common iliac arteries. The [median sacral artery](#) supplies the [sacrum](#), [coccyx](#), and the [posterior](#) wall of the [rectum](#). The [ischemic](#) hepatic [colonic](#) flexure in this patient is perfused by another [artery](#).


B - Inferior mesenteric artery

Explanation Why

The [inferior mesenteric artery](#) (IMA) branches off from the [abdominal aorta](#) at the level of L3. This [artery](#) supplies the last third of the [transverse colon](#), the [descending colon](#), the [sigmoid colon](#), and the upper [rectum](#) ([superior rectal artery](#)), all structures that arise from the embryological [hindgut](#). There is a [watershed area](#) on the [transverse colon](#) at the point of the [splenic flexure](#), where limited vascularity increases the risk of [ischemia](#) in cases of hypoperfusion. However, the [ischemic](#) hepatic [colonic](#) flexure in this patient is perfused by another [artery](#).

C - Celiac artery

Image



Explanation Why

The [celiac artery](#) arises from the [abdominal aorta](#) at the level of T12, directly where the aorta enters the abdomen. The [artery](#) supplies the derivatives of the [foregut](#), from the [distal](#) portion of the [esophagus](#) to the descending part of the duodenum. The three major branches of the [celiac artery](#) are the [left gastric artery](#), the [splenic artery](#), and the [common hepatic artery](#). However, the [ischemic](#) hepatic [colonic](#) flexure in this patient is perfused by another [artery](#).

D - Internal iliac artery


Explanation Why

The internal iliac artery branches off from the common iliac artery at the level of the sacral brim, entering the [pelvis](#). The visceral organs that the [internal iliac artery](#) supplies include the [bladder](#),

uterus, [vagina](#), and inferior [rectum](#) (via the inferior and [middle rectal arteries](#)). The [ischemic](#) hepatic [colonic](#) flexure in this patient is perfused by another [artery](#).

E - Superior mesenteric artery

Image



Explanation Why

The [superior mesenteric artery](#) (SMA) is the site of blood flow interruption in 90% of [acute mesenteric ischemia](#) cases. The SMA supplies the derivatives of the embryological [midgut](#), including the [small intestine](#) ([duodenum](#), [jejunum](#), and [ileum](#)), [ascending colon](#) ([right colic artery](#)), and first two-thirds of the [transverse colon](#) ([middle colic artery](#)). The [cecum](#) and [appendix](#) are supplied by the [ileocolic artery](#).

Question # 12

A 73-year-old man comes to the physician because of a 2-month history of intermittent blood in his stool. He has had no pain with defecation. Physical examination shows a 2-cm mass located above the dentate line. Further evaluation of the mass confirms adenocarcinoma. Which of the following best describes the most likely route of hematogenous spread of the malignancy?

	Answer	Image
A	Superior rectal vein → inferior mesenteric vein → hepatic portal vein	
B	Inferior rectal vein → inferior mesenteric vein → splenic vein	
C	Inferior rectal vein → internal pudendal vein → internal iliac vein	
D	Superior rectal vein → superior mesenteric vein → splenic vein	
E	Superior rectal vein → superior mesenteric vein → hepatic portal vein	

	Answer	Image
F	Inferior rectal vein → internal pudendal vein → external iliac vein	


Hint

The anal canal is anatomically divided by the pectinate line into the upper two thirds, which is embryologically derived from the hindgut (endoderm), and the lower third, which is derived from ectoderm. The artery supply, venous drainage, lymphatic drainage, and nerve supply of the region of the anal canal above the dentate line (pectinate line) are, therefore, different from those of the region below this line.

Correct Answer

A - Superior rectal vein → inferior mesenteric vein → hepatic portal vein

Image



Explanation Why

The region of the [anal canal](#) above the [dentate line](#) drains into the [portal venous](#) system via the following pathway: [superior rectal vein](#) → [inferior mesenteric vein](#) → [splenic vein](#) → [hepatic portal vein](#). [Adenocarcinomas](#) typically arise in this region and usually result in hepatic [metastases](#). The region of the [anal canal](#) below the [dentate line](#) drains into systemic venous circulation via the following pathway: [inferior rectal vein](#) → [internal pudendal vein](#) → [internal iliac vein](#) → [inferior vena cava](#). [Squamous cell carcinomas](#) typically arise in this region. The principles of the upper and lower [anal canal](#)'s venous drainage are not only relevant for predicting the most likely route of hematogenous spread of malignancies, but also for understanding [first pass metabolism](#) in rectal drug administration. [First pass metabolism](#) is higher if the drug drains via the [superior rectal vein](#) into the hepatic portal circulation.

B - Inferior rectal vein → inferior mesenteric vein → splenic vein

Explanation Why

The inferior rectal vein does not drain into the [inferior mesenteric vein](#).

C - Inferior rectal vein → internal pudendal vein → internal iliac vein

Explanation Why

“Inferior rectal vein → internal pudendal vein → [internal iliac vein](#) (→ [inferior vena cava](#))” describes the venous pathway of dissemination for tumors located below the [dentate line](#). This patient's [tumor](#) is located [above the dentate line](#).

D - Superior rectal vein → superior mesenteric vein → splenic vein

Explanation Why

The [superior mesenteric vein](#) does not drain into the [splenic vein](#) unless retrograde flow due to [portal hypertension](#) is present. This patient does not show signs of [portal hypertension](#).

E - Superior rectal vein → superior mesenteric vein → hepatic portal vein

Explanation Why

Blood from the [superior rectal vein](#) does not drain into the [superior mesenteric vein](#).

F - Inferior rectal vein → internal pudendal vein → external iliac vein

Explanation Why

Blood from the internal pudendal vein does not drain into the external iliac vein.

Question # 13

A 78-year-old man comes to the emergency department because of a 4-hour history of abdominal pain. Abdominal examination shows guarding with diffuse rebound tenderness. A CT scan of the abdomen shows an infrarenal abdominal aortic aneurysm that extends up to the level of the L4 vertebra and a partial filling defect in the anterior wall of the aneurysm. Which of the following bowel regions is at greatest risk for requiring resection?

	Answer	Image
A	Splenic flexure, descending colon, and sigmoid colon	
B	Hepatic flexure, transverse colon, and splenic flexure	
C	Cecum, ileum, and distal jejunum	
D	Cecum, ascending colon, and hepatic flexure	
E	Ascending colon, cecum, and distal ileum	


Hint

The renal arteries originate from the lateral surface of the aorta at the level of the L2 vertebra. The inferior mesenteric artery (IMA) arises from the anterior surface of the aorta at the level of the L3 vertebra. Occlusion at the opening of the IMA by a thrombus within an aortic aneurysm (suggested by the filling defect) would cause acute mesenteric ischemia and, potentially, necrosis of the bowel segments supplied by this artery.

Correct Answer

A - Splenic flexure, descending colon, and sigmoid colon

Image



Explanation Why

The [splenic flexure](#), [descending colon](#), and [sigmoid colon](#), which are supplied by the [inferior mesenteric artery](#) (IMA), are most at risk for requiring surgical resection in this patient's case. The IMA also supplies the [rectum](#) and the region of the [anal canal above the pectinate line](#) via the [superior rectal artery](#).

B - Hepatic flexure, transverse colon, and splenic flexure

Explanation Why

The hepatic flexure, [transverse colon](#), and [splenic flexure](#) are predominantly supplied by the [superior](#)

[mesenteric artery](#) (SMA) via the [middle colic artery](#) and the left colic branch of the [inferior mesenteric artery](#) (IMA). The SMA arises anteriorly at the level of the L1 [vertebra](#), above the level of the [renal arteries](#). Although this patient is likely to have an occlusion of the IMA, the SMA would not be affected because his [aneurysm](#) and filling defect are infrarenal. Therefore, this combination of bowel regions is less likely to be affected in this patient's case.

C - Cecum, ileum, and distal jejunum

Explanation Why

The [cecum](#), [ileum](#), and [distal jejunum](#) are supplied by the [superior mesenteric artery](#) (SMA), which arises anteriorly from the aorta at the level of the L1 [vertebra](#), above the level of origin of the [renal arteries](#). As this patient's [aneurysm](#) and filling defect are infrarenal, this combination of bowel regions would not be affected.

D - Cecum, ascending colon, and hepatic flexure

Explanation Why

The [cecum](#), [ascending colon](#), and hepatic flexure are supplied by the right colic branch of the [superior mesenteric artery](#) (SMA). The SMA arises anteriorly from the aorta at the level of the L1 [vertebra](#), above the origin of the [renal arteries](#). As this patient's [aneurysm](#) and filling defect are infrarenal, this combination of bowel regions would not be affected.

E - Ascending colon, cecum, and distal ileum

Explanation Why

The [ascending colon](#), [cecum](#), and [distal ileum](#) are supplied by the ileocolic and right colic branches of the [superior mesenteric artery](#) (SMA), which arises anteriorly from the aorta at the level of the L1 [vertebra](#), above the level of origin of the [renal arteries](#). As this patient's [aneurysm](#) and filling defect are infrarenal, this combination of bowel regions would not be affected.

Question # 14

A 29-year-old man is brought to the emergency room 6 hours after the onset of severe epigastric pain and vomiting. His heart rate is 110/min and blood pressure is 98/72 mm Hg. He is diagnosed with acute pancreatitis, and fluid resuscitation with normal saline is initiated. Which of the following is the most likely immediate effect of fluid resuscitation in this patient?

	Answer	Image
A	Increase in cardiac afterload	
B	Increase in volume of distribution	
C	Increase in myocardial oxygen demand	
D	Increase in plasma oncotic pressure	
E	Increase in glomerular filtration fraction	

Hint

Fluid resuscitation would increase venous return and right ventricular end diastolic volume (EDV).

Correct Answer

A - Increase in cardiac afterload

Explanation Why

Cardiac [afterload](#) represents the pressure against which the [heart](#) must pump during [systole](#). A large, rapid infusion of [normal saline](#) can lead to transient hemodilution, decreasing blood viscosity, and causing a decrease in [afterload](#), not an increase.


B - Increase in volume of distribution

Explanation Why

[Volume of distribution](#) (V_D) is an intrinsic measured property of a drug, not a physiological property of the body. Although the infusion of [normal saline](#) given to this patient will temporarily increase his [total body water](#), the V_D of any particular drug would remain unaffected.

C - Increase in myocardial oxygen demand

Image



Explanation Why

Myocardial oxygen demand depends on the [heart rate](#), [heart](#) contractility, and wall tension (affected by [afterload](#)). An increase in [end-diastolic volume](#), as occurs in [fluid resuscitation](#), causes an increase in cardiac [preload](#) by stretching cardiac [myocyte sarcomeres](#), which results in increased [myocardial contractility](#) based on the [Frank-Starling mechanism](#). Increased contractility also increases [cardiac output](#) (CO) and cardiac work, and hence myocardial oxygen demand as well. Note, however, that [fluid resuscitation](#) is beneficial in [hypovolemic](#) patients because it also increases [myocardial](#) oxygen supply by improving coronary perfusion pressure.

D - Increase in plasma oncotic pressure

Explanation Why

Plasma [oncotic pressure](#) is exerted by [proteins](#) (primarily [albumin](#)) within blood vessels. A transient increase in plasma [oncotic pressure](#) would occur with the infusion of [colloids](#) but not with [normal saline](#) infusion. [Normal saline](#) is isotonic to serum and generally redistributes to replenish depleted [extracellular fluid](#), only minimally affecting plasma [oncotic pressure](#). A large, rapid infusion of [normal saline](#) can decrease plasma [oncotic pressure](#) transiently by diluting the concentration of protein intravascularly.

E - Increase in glomerular filtration fraction

Explanation Why

The [glomerular filtration fraction](#) (FF) is the [ratio](#) of [GFR](#) to RPF. [Fluid resuscitation](#) would increase [renal blood flow](#) (and hence RPF) as well as [GFR](#). However, with a normally functioning [kidney](#), the percentage increase in [GFR](#) (ΔGFR) is always less than the percentage increase in RPF (ΔRPF) at normal or [low blood pressures](#) because of [tubuloglomerular feedback](#). Therefore, the [FF](#) would decrease rather than increase in this patient. The [FF](#) would increase in [dehydration](#), which causes a decrease in RPF that is disproportionate to the accompanying decrease in [GFR](#), as well as at [high blood pressures](#) or fluid overload, in which pressure [natriuresis](#) occurs and ΔGFR is greater than ΔRPF .

Question # 15

A 56-year-old man comes to the physician for a follow-up examination 1 week after a temporal artery biopsy was performed to evaluate headache and blurry vision. He has a 12-year history of hypertension treated with lisinopril. Results of the testing show decreased concentrations of arginine in vascular endothelial cells. Which of the following is the most likely explanation for this finding?

	Answer	Image
A	Increased binding of calcium to calmodulin	
B	Increased phosphorylation of myosin	
C	Increased production of nitric oxide	
D	Decreased production of cyclic guanosine monophosphate	
E	Decreased concentration of bradykinin	

Hint

Lisinopril, like other ACE inhibitors, causes relaxation of vascular smooth muscle cells.

Correct Answer

A - Increased binding of calcium to calmodulin

Explanation Why

Increased binding of calcium to [calmodulin](#) results in the activation of [myosin-light-chain kinase](#), which [phosphorylates myosin](#) to cause contraction, not relaxation, of vascular [smooth muscle cells](#). Calcium and [calmodulin](#) do not play a role in vascular [smooth muscle cell](#) relaxation.

B - Increased phosphorylation of myosin

Explanation Why

The [phosphorylation](#) of [myosin](#) by [myosin-light-chain kinase](#) results in contraction, not relaxation, of vascular [smooth muscle cells](#). ACE inhibitor therapy causes relaxation of vascular [smooth muscle cells](#) by promoting the dephosphorylation of [myosin](#) by [myosin-light-chain phosphatase](#).

C - Increased production of nitric oxide

Explanation Why

[Angiotensin-converting enzyme](#) (ACE) inactivates [bradykinin](#), an inflammatory mediator that increases intracellular calcium in [endothelial](#) cells. Intracellular calcium stimulates [nitric oxide synthase](#) to convert [arginine](#) to [nitric oxide](#), which mediates the relaxation of vascular [smooth muscle](#) cells to cause [vasodilation](#). Thus, ACE inhibitor therapy, which increases [bradykinin](#) concentrations, leads to increased production of [nitric oxide](#) from [arginine](#), which leads to decreased concentrations of [arginine](#) in vascular [endothelial](#) cells.

D - Decreased production of cyclic guanosine monophosphate

Explanation Why

[Cyclic guanosine monophosphate \(cGMP\)](#) is a compound that mediates the [vasodilatory](#) effects of ACE inhibitor therapy by acting on [myosin-light-chain phosphatase](#) to dephosphorylate [myosin](#), which leads to vascular [smooth muscle cell](#) relaxation. ACE inhibitor therapy increases, not decreases, the production of [cGMP](#).

E - Decreased concentration of bradykinin

Explanation Why

[Bradykinin](#), an inflammatory mediator, is inactivated by [angiotensin-converting enzyme \(ACE\)](#). Thus, ACE inhibitor therapy increases [bradykinin](#) concentrations.

Question # 16

A 62-year-old man with coronary artery disease and type 2 diabetes mellitus comes to the emergency department because of frequent episodes of palpitations for 4 days. His pulse is 165/min and blood pressure is 98/70 mm Hg. An ECG shows monomorphic ventricular tachycardia. Intravenous pharmacotherapy is begun with a highly lipophilic drug that prolongs the duration of the myocardial action potential and refractory period and inhibits adrenergic stimulation of the myocardium. The patient's ventricular tachycardia converts to normal sinus rhythm. He is discharged home with an oral formulation of the same drug. Which of the following adverse effects is most commonly associated with long-term use of this medication?

	Answer	Image
A	Priapism	
B	Acne vulgaris	
C	Angle-closure glaucoma	
D	Hyperthyroidism	
E	Hepatic adenoma	
F	Basilar predominant emphysema	

Hint

The patient was started on amiodarone, which is often used to control hemodynamically-stable monomorphic ventricular tachycardia. Amiodarone is a lipophilic drug that prolongs the duration of the cardiac action potential and refractory period and inhibits adrenergic stimulation of the myocardium.

Correct Answer

A - Priapism

Explanation Why

[Priapism](#) is an adverse effect most commonly associated with the use of [trazodone](#) and [PDE5 inhibitors](#). Genitourinary side effects of [amiodarone](#), on the other hand, include [sexual dysfunction](#), sterile [epididymitis](#), and scrotal [pain](#) and swelling.

B - Acne vulgaris

Explanation Why

[Acne](#) is a cutaneous side effect of several medications, including [lithium](#), [glucocorticoids](#), and [anabolic steroids](#). Cutaneous manifestations of [amiodarone](#) toxicity, on the other hand, include [photosensitivity](#) reactions and bluish discoloration of the [skin](#) (blue man syndrome)

C - Angle-closure glaucoma

Explanation Why

[Angle-closure glaucoma](#) can be exacerbated by drugs with [anticholinergic](#) activity such as tricyclic [antidepressants](#) (TCAs), [antihistamines](#), decongestants, and [sympathomimetics](#). Ocular side effects related to [amiodarone](#) toxicity, on the other hand, include corneal microdeposits and optic neuropathy.

D - Hyperthyroidism

Explanation Why

A common adverse effect of [amiodarone](#) is [thyroid tissue](#) damage, which can cause [hyperthyroidism](#) or [hypothyroidism](#). It is especially problematic in patients with pre-existing [thyroid](#) conditions. For this reason, [thyroid](#) function should be monitored closely in patients taking [amiodarone](#).

E - Hepatic adenoma

Explanation But

[Amiodarone](#) should be discontinued in patients with [liver](#) enzyme levels more than twice the normal value.

Explanation Why

[Hepatic adenomas](#) are an adverse effect of [oral contraceptive pill \(OCP\)](#) use. A transient increase in hepatic enzymes may be seen with [amiodarone](#) use, which can cause [cirrhosis](#) and hepatic failure in rare cases. However, there is no known association between [amiodarone](#) and [hepatic adenoma](#).


F - Basilar predominant emphysema

Explanation Why

Basilar predominant [emphysema](#) is associated with [alpha-1 antitrypsin deficiency](#) but not [amiodarone](#) use. Although [amiodarone](#) can lead to pulmonary complications, it increases rather the risk of chronic [interstitial pneumonitis](#), which can manifest with an insidious onset of a nonproductive [cough](#), [dyspnea](#), weight loss, and fatigue.

Question # 17

A 67-year-old woman comes to the emergency department because of a 4-month history of fatigue, shortness of breath with exertion, and dizziness. She has a history of atrial fibrillation and had a single-chamber pacemaker placed five years ago after an episode of syncope. Her pulse is 66/min and blood pressure is 98/66 mm Hg. An x-ray of the chest is shown. The x-ray confirms termination of the pacemaker lead in which of the following structures?



	Answer	Image
A	Superior vena cava	
B	Left ventricle	

	Answer	Image
C	Right ventricle	
D	Left atrium	
E	Right atrium	



Hint

This patient's history and presentation are consistent with pacemaker syndrome, in which there is dysynchrony between the atria and ventricles. The image shows a posterioranterior chest x-ray with a single pacemaker lead.

Correct Answer

A - Superior vena cava

Explanation Why

Pacemaker wires travel through the [superior vena cava \(SVC\)](#) in order to enter the [heart](#). They should, however, terminate in a structure other than the [SVC](#) to be effective. On a posteroanterior [x-ray](#) of the chest, the [SVC](#) is superior to the right [lateral](#) aspect of the cardiac contour. The lead in this patient's [x-ray](#) terminates inferior and to the left of the [SVC](#).

B - Left ventricle

Explanation But

Biventricular pacemakers are used to treat abnormal intraventricular conduction, such as a [left bundle branch block](#) or advanced [systolic heart failure](#).

Explanation Why

[Left ventricular](#) pacemaker leads are almost exclusively seen with biventricular pacemakers, which have leads terminating in both the right and the left ventricles. During biventricular pacemaker placement, the [left ventricular](#) lead is passed through the [coronary sinus](#) into a coronary [vein](#) (usually the [lateral](#) or posterolateral branch), from which the lead then paces the [left ventricle](#) transvenously. On a posteroanterior [x-ray](#) of the chest, a [left ventricle](#) lead would be visualized as crossing from the [right atrium](#) into the [coronary sinus](#) and then following a [lateral](#) course along the left [heart](#) contour. This patient's finding of a single pacer lead and history of [atrial fibrillation](#) are not consistent with a [left ventricular](#) pacemaker.

C - Right ventricle

Image



Explanation Why

On a posteroanterior [x-ray](#) of the chest, the [right ventricle](#) makes up the [anterior](#) contour of the [heart](#) and is centered on the [anterior mediastinal](#) plane. The [right ventricle](#) does not contribute to any of the visible [heart](#) contours seen on a PA [x-ray](#) (the right [heart](#) contour is composed primarily of the [right atrium](#)), which can make its identification difficult. However, this patient's chest [x-ray](#) clearly shows her single-chamber pacing lead course through the right [subclavian vein](#), [superior vena cava](#), and [right atrium](#) before terminating in the region of the [right ventricle](#).

D - Left atrium

Explanation Why

On a posteroanterior [x-ray](#) of the chest, the [left atrium](#) sits posteriorly and therefore is not readily

visible apart from the [left atrial appendage](#), which is visualized at the superior left [lateral](#) border of the [heart](#). The lead in this patient's [x-ray](#) terminates inferior to and to the right of the [left atrial appendage](#).

E - Right atrium

Explanation Why

A single chamber pacemaker with a lead placed in the [right atrium](#) can be used in the treatment of [sick sinus syndrome](#). On a posteroanterior [x-ray](#) of the chest, the [right atrium](#) makes up the right [heart](#) contour and is located to the [lateral](#) right of the [mediastinum](#). The lead in this patient's [x-ray](#), however, terminates to the left of the [anterior mediastinum](#).

Question # 18

A 67-year-old woman comes to the physician for chest tightness, shortness of breath, and lightheadedness. She has experienced these symptoms during the past 2 weeks while climbing stairs but feels better when she sits down. She had a cold 2 weeks ago but has otherwise been well. She appears short of breath. Her respirations are 21/min and blood pressure is 131/85 mmHg. On cardiovascular examination, a late systolic ejection murmur is heard best in the second right intercostal space. The lungs are clear to auscultation. Which of the following mechanisms is the most likely cause of this patient's current condition?

	Answer	Image
A	Inflammatory constriction of the bronchioles	
B	Narrowing of the coronary arterial lumen	
C	Cellular injury of the esophageal epithelium	
D	Critical transmural hypoperfusion of the myocardium	
E	Increased left ventricular oxygen demand	

Hint

This patient is presenting with chest tightness consistent with angina. She also has a late ejection systolic murmur on examination, which should raise suspicion of aortic stenosis.

Correct Answer

A - Inflammatory constriction of the bronchioles

Explanation Why

A combination of [inflammation](#) and constriction of the [bronchioles](#) describes the mechanism behind virus-induced bronchospasm. Common respiratory viruses (as this patient likely had) can cause [epithelial](#) damage to the [airways](#), induce [inflammation](#) and bronchospasm, and subsequently trigger wheezing in both patients with and without underlying [asthma](#). While her recent cold and exertional dyspnea could be consistent with bronchospasm, her [presyncopal](#) symptoms and [heart murmur](#) suggest a different mechanism.

B - Narrowing of the coronary arterial lumen

Explanation Why

Narrowing of the [coronary arteries](#) describes the mechanism of [coronary artery disease](#) (CAD), which can result in [myocardial ischemia](#) due to inadequate oxygen supply to the [myocardium](#) during exertion. While this patient clearly has [angina](#), as well as strong [risk factors](#) for CAD (age), her [presyncopal](#) symptoms and [heart murmur](#) are not explained by CAD alone and a different mechanism is more likely to be the cause of her symptoms.

C - Cellular injury of the esophageal epithelium

Explanation Why

Injury to the esophageal [epithelium](#) is the mechanism behind [GERD](#) and is secondary to acid exposure from reflux of gastric contents. [GERD](#) should always be considered in the differential for [chest pain](#), as it may commonly present with atypical [chest pain](#). However, this patient presents with typical [chest pain](#) (e.g., exertional chest tightness that resolves with rest) consistent with [angina](#), and her [presyncopal](#) symptoms and [heart murmur](#) suggest a different mechanism.

D - Critical transmural hypoperfusion of the myocardium

Explanation Why

Critical transmural hypoperfusion of the [myocardium](#) describes the mechanism behind [myocardial infarction](#) (MI), a form of [acute coronary syndrome](#). This patient's symptoms, while concerning, are transient and resolve with rest, making acute MI very unlikely. Her [heart murmur](#) make a different mechanism more likely.

E - Increased left ventricular oxygen demand

Explanation But

The presence of symptoms with [aortic stenosis](#) (AS) is an indication for surgical repair, as symptomatic AS carries a poor prognosis.

Explanation Why

The narrowed opening of a stenotic [aortic valve](#) during [systole](#) obstructs blood flow from the [left ventricle](#) (LV), which over time results in [left ventricular hypertrophy](#) (LVH). LVH subsequently results in increased LV oxygen demand, leading to [myocardial ischemia](#), which manifests as [angina](#) especially during periods of strenuous exercise or stress. The reduction in [cardiac output](#) during exertion also causes transient cerebral hypoperfusion, which explains her [presyncopal](#) symptoms.

Question # 19

A 49-year-old man comes to the physician because of a 5-month history of progressive fatigue and exertional dyspnea. Cardiac examination shows a loud S₂ in the 2nd left intercostal space. Right heart catheterization shows a pulmonary artery pressure of 32 mm Hg. Treatment with bosentan is initiated. The beneficial effect of this drug is due to binding to which of the following?

	Answer	Image
A	L-type voltage-gated calcium channels	
B	Phosphodiesterase-5	The diagram illustrates the signaling pathway for smooth muscle relaxation. In the endothelium, stimuli like ACh, histamine, or shear stress activate eNOS, which converts L-arginine to NO. Nitrates also produce NO. NO then enters the smooth muscle cell and activates Guanylyl cyclase, which converts GTP to cGMP. cGMP activates Protein kinase G (PKG), which in turn inhibits the MLCK/Myosin cycle and the sarcoplasmic reticulum Ca²⁺ release. This leads to a decrease in intracellular Ca²⁺, resulting in smooth muscle relaxation and vasodilation. PDE5 is shown breaking down cGMP to GMP. PDE5 inhibitors block this breakdown. The diagram is color-coded: red for effects of cGMP, yellow for breakdown of cGMP, and green for physiological NO production.
C	Prostacyclin receptor	
D	Adenosine receptors	
E	Endothelin receptors	

Hint

The target of this drug also plays an important role in hemostasis.

Correct Answer


A - L-type voltage-gated calcium channels

Explanation Why

[Calcium channel blocking agents](#) (e.g., [nifedipine](#)) can block [L-type voltage-gated calcium channels](#). While this type of agent can be used to treat [pulmonary artery hypertension](#), [bosentan](#) has a different mechanism of action.

B - Phosphodiesterase-5

Image



Explanation Why

[Phosphodiesterase-5 inhibitors](#) (e.g., [sildenafil](#)) prevent the degradation of [cGMP](#), which normally promotes [vasodilation](#) and relaxation of [smooth muscles](#). Although [phosphodiesterase-5 inhibitors](#)

can be used to treat [pulmonary arterial hypertension](#), [bosentan](#) has a different mechanism of action.

C - Prostacyclin receptor

Explanation Why

[Prostacyclin](#) analogs (e.g., [iloprost](#)) have an [agonist](#) effect on the [prostacyclin](#) receptor. [Prostacyclin](#) is a potent [vasodilator](#) with antiplatelet effects. Although [prostacyclin](#) receptors can be administered in [pulmonary arterial hypertension](#), [bosentan](#) has a different mechanism of action.

D - Adenosine receptors

Explanation Why

Adenosine is a drug commonly used to terminate [supraventricular tachycardia](#) because it hyperpolarizes the [AV node](#) by increasing potassium conductance. Adenosine receptor antagonists (e.g., methylxanthines) induce [smooth muscle](#) relaxation and bronchodilation and are therefore used in the treatment of [asthma](#). [Bosentan](#), however, does not bind to the adenosine receptor.

E - Endothelin receptors

Explanation Why

[Endothelin receptor antagonists](#) (e.g., [bosentan](#)) competitively antagonize endothelin-1 receptors. Endothelin normally promotes vasoconstriction and can increase resistance in pulmonary vessels, leading to [pulmonary arterial hypertension](#). Targeting the endothelin receptor, therefore, prevents pulmonary vasoconstriction.

Question # 20

A 38-year-old woman comes to the physician because of a 10-month history of nonbloody diarrhea and recurrent episodes of flushing and wheezing. She does not take any medications. Physical examination shows a hyperpigmented rash around the base of her neck. Cardiac examination shows a grade 3/6, holosystolic murmur at the left lower sternal border. Echocardiography shows right-sided endocardial and valvular fibrosis with moderate tricuspid regurgitation; there are no septal defects or left-sided valvular defects. Urinalysis shows increased 5-hydroxyindoleacetic acid concentration. Further evaluation of this patient is most likely to show which of the following findings?

	Answer	Image
A	Tumor in the pancreas without metastasis	
B	Tumor in the terminal ileum with hepatic metastasis	
C	Tumor in the left atrium with hepatic metastasis	
D	Tumor in the appendix without metastasis	
E	Tumor in the lung without metastasis	
F	Tumor in the descending colon with hepatic metastasis	

Hint

This patient's presentation (e.g., cutaneous vasomotor flushing, diarrhea, wheezing, and cardiac involvement) is highly suggestive of carcinoid syndrome. 5-hydroxyindoleacetic acid (5-HIAA) is a metabolite of serotonin, the primary molecule produced by carcinoid tumors. Carcinoid heart disease is typically caused by valvular and endocardial fibrosis that affects the right heart, leading to right-sided valvular disease (e.g., tricuspid regurgitation).

Correct Answer

A - Tumor in the pancreas without metastasis

Explanation Why

While a [tumor](#) in the [pancreas](#) could be a [serotonin](#)-producing neuroendocrine neoplasm that causes [carcinoid syndrome](#), systemic symptoms are normally not seen in the absence of hepatic [metastasis](#). Metabolites produced by a [pancreatic carcinoid tumor](#) would be degraded by the [liver](#) ([first pass effect](#)), making this diagnosis unlikely in this patient. Other more common neuroendocrine [pancreatic](#) tumors such as [somatostatinoma](#), [glucagonoma](#), and [insulinoma](#) do not secrete [serotonin](#) and, therefore, would not manifest with [carcinoid syndrome](#).

B - Tumor in the terminal ileum with hepatic metastasis

Explanation But

In patients with a [metastasized carcinoid tumor](#) with a [patent foramen ovale](#), high amounts of [tumor](#) products pass directly from the [right atrium](#) to the [left atrium](#), bypassing the [pulmonary circulation](#), and causing damage to the left side of the [heart](#).

Explanation Why

The terminal [ileum](#) is a common location of [carcinoid tumors](#). A gastrointestinal [carcinoid tumor](#) that has [metastasized](#) to the [liver](#) (or beyond) causes systemic symptoms of [carcinoid syndrome](#), as large quantities of vasoactive metabolites ([serotonin](#), [histamine](#), [VIP](#)) circulate throughout the body without being degraded in the [liver](#). In the [heart](#), [serotonin](#) can cause endocardial [fibrosis](#), resulting in [carcinoid heart disease](#). In patients with [carcinoid heart disease](#), the right side of the [heart](#) is typically affected as the [lungs](#) enzymatically inactivate the metabolites; therefore, the left side of the [heart](#) is usually spared. Increased [serotonin](#) metabolism explains this patient's elevated [5-HIAA](#) levels and has resulted in [pellagra](#) ([niacin deficiency](#)), which classically manifests with the triad of [diarrhea](#), [dementia](#), and [dermatitis](#).

C - Tumor in the left atrium with hepatic metastasis

Explanation Why

[Atrial myxoma](#) is the most common primary cardiac [tumor](#) in adults. A myxoma growing in the [mitral area](#) could cause a [murmur](#), which is seen in this patient. However, the [murmur](#) caused by an [atrial myxoma](#) is usually [diastolic](#) and heard over the apex of the [heart](#) (similar to [mitral stenosis](#)) rather than [systolic](#) and heard at the left lower border. Additionally, [atrial myxomas](#) are not neuroendocrine tumors and therefore would not cause elevated [5-HIAA](#) levels and [carcinoid syndrome](#). Finally, [cardiac myxomas](#) are usually benign and do not [metastasize](#).

D - Tumor in the appendix without metastasis

Explanation Why

Although the [appendix](#) is the most common site of [carcinoid tumors](#), systemic symptoms are not normally seen in the absence of hepatic [metastasis](#). Metabolites produced by a [carcinoid tumor](#) in the [appendix](#) are degraded by the [liver](#) via the [first pass effect](#).

E - Tumor in the lung without metastasis

Explanation Why

While a [tumor](#) in the [lung](#) could be a [serotonin](#)-producing [carcinoid tumor](#) that causes [carcinoid syndrome](#), systemic symptoms are rarely seen in [lung](#) neuroendocrine tumors in the absence of [metastasis](#). Neuroendocrine tumors of the [lung](#) secrete smaller amounts of vasoactive metabolites than neuroendocrine tumors of the [gastrointestinal tract](#). Additionally, a neuroendocrine tumor of the [lung](#) would affect the left [heart](#) rather than the right [heart](#). Because the [lung](#) inactivates neuroendocrine metabolites, [carcinoid heart disease](#) is exceedingly rare in patients with [lung](#) neuroendocrine tumors.

F - Tumor in the descending colon with hepatic metastasis

Explanation Why

A [tumor](#) in the [descending colon](#) with hepatic [metastasis](#) would most likely be an [adenocarcinoma](#), the most common type of [colorectal cancer](#). This patient's symptoms, however, are consistent with [carcinoid syndrome](#) caused by a [carcinoid tumor](#). Although [carcinoid tumors](#) can occur in the [gastrointestinal tract](#), they would typically be found in the terminal [ileum](#) or the [appendix](#).

Question # 21

A 58-year-old man comes to the physician because of a sore throat and painful lesions in his mouth for the past few days. Six weeks ago, he underwent cardiac catheterization and stent implantation of the left anterior descending artery for treatment of acute myocardial infarction. Pharmacotherapy with dual antiplatelet medication was started. His temperature is 38.1°C (100.6°F). Oral examination shows several shallow ulcers on the buccal mucosa. Laboratory studies show:

Hematocrit	41.5%
Leukocyte count	1,050/mm ³
Segmented neutrophils	35%
Platelet count	175,000/mm ³

Which of the following drugs is most likely responsible for this patient's current condition?

	Answer	Image
A	Abciximab	
B	Ticlopidine	
C	Clopidogrel	
D	Apixaban	
E	Enoxaparin	

	Answer	Image
F	Aspirin	

Hint

This patient presents with oral ulcers, sore throat, fever, and a low neutrophil count. Given that he was recently started on new medications, the most likely diagnosis is drug-induced neutropenia.

Correct Answer

A - Abciximab

Explanation Why

[Abciximab](#) is indicated only in high-risk patients with [unstable angina/NSTEMI](#) planned for [PCI](#) within 24 hours to prevent thrombotic complications. Side effects include acute profound [thrombocytopenia](#) and hemorrhage. However, [abciximab](#) does not cause [neutropenia](#) ([agranulocytosis](#)), which is seen in this patient's case.

B - Ticlopidine

Explanation But

Other [P2Y12 receptor antagonists](#) have been shown to be safer than [ticlopidine](#) when used for dual antiplatelet therapy.

Explanation Why

[Ticlopidine](#) is a [P2Y12 receptor antagonist](#) ([ADP receptor blocker](#)) that blocks [platelet aggregation](#). Side effects include [allergic reactions](#) (rash, [pruritus](#), [anaphylaxis](#)), hemorrhage, and gastrointestinal complications. A rare side effect is [neutropenia](#) ([agranulocytosis](#)), which is most commonly seen during the first 3 months of treatment. Other severe hematologic adverse effects of [ticlopidine](#) include [thrombotic thrombocytopenic purpura](#) and [aplastic anemia](#). Weekly monitoring of complete blood counts is indicated during the initial phase of treatment with [ticlopidine](#).

C - Clopidogrel

Explanation Why

[Clopidogrel](#) is used for dual antiplatelet therapy with [aspirin](#) as well as an alternative to [aspirin](#) in cases of intolerance. Side effects include [allergic reactions](#) (rash, [pruritus](#), [anaphylaxis](#)), hemorrhage,

and gastrointestinal complications. However, [clopidogrel](#) use does not cause [neutropenia \(agranulocytosis\)](#), which is seen in this patient.

D - Apixaban

Explanation Why

[Apixaban](#) is used for the treatment and prophylaxis of [venous thromboembolism](#). However, it is not an [antiplatelet agent](#) and therefore is not used in dual antiplatelet therapy. Although side effects of [apixaban](#) include bleeding and [thrombocytopenia](#), [neutropenia \(agranulocytosis\)](#), as seen in this patient, would not be expected.

E - Enoxaparin

Explanation Why

[Enoxaparin](#) is primarily used for prophylaxis and treatment of [venous thromboembolism](#). However, it is not an [antiplatelet agent](#) and therefore is not used in dual antiplatelet therapy. Side effects include [allergic reactions](#), [heparin-induced thrombocytopenia](#), and [osteoporosis](#). [Enoxaparin](#), however, does not cause [neutropenia \(agranulocytosis\)](#), which is seen in this patient's case.

F - Aspirin

Explanation Why

Low-dose [aspirin](#) is used in the management and prophylaxis of cardiovascular disease. It is often used in combination with other drugs for dual antiplatelet therapy. Serious side effects include [peptic ulcers](#), [allergic reactions](#), [interstitial](#) nephritis, and hemorrhage. However, [aspirin](#) does not cause isolated [neutropenia \(agranulocytosis\)](#), which is seen in this patient's case.

Question # 22

A 66-year-old man with coronary artery disease and hypertension comes to the emergency department because of intermittent retrosternal chest pain, lightheadedness, and palpitations. He has smoked one pack of cigarettes daily for 39 years. His pulse is 140/min and irregularly irregular, respirations are 20/min, and blood pressure is 108/60 mm Hg. An ECG shows an irregular, narrow-complex tachycardia with absent P waves. A drug with which of the following mechanisms of action is most likely to be effective in the long-term prevention of embolic stroke in this patient?

	Answer	Image
A	Binding and activation of antithrombin III	
B	Irreversible inhibition of cyclooxygenase	
C	Interference with carboxylation of glutamate residues	
D	Activation of the conversion of plasminogen to plasmin	
E	Irreversible blockade of adenosine diphosphate receptors	

Hint

The patient's lightheadedness, palpitations, and irregular rhythm with no visible P waves on ECG are highly suggestive of atrial fibrillation (Afib). Warfarin is commonly used for long-term anticoagulation in patients with Afib who are at increased risk of embolic stroke.

Correct Answer

A - Binding and activation of antithrombin III

Explanation Why

[Heparin](#) binds to and potentiates the effects of [antithrombin III](#), which inactivates [thrombin](#) and inhibits the [coagulation factors](#) II, VII, IX, X, XI, and XII. [Heparin](#) has a rapid onset of action and is used for anticoagulation in the acute management of thromboses, emboli, and [unstable angina](#). While it may also be used initially for patients with [Afib](#), it is not a good long-term option as it is only administered subcutaneously or intravenously. A different agent would be indicated in this setting.

B - Irreversible inhibition of cyclooxygenase

Explanation Why

[Acetylsalicylic acid](#) (ASA) irreversibly inhibits [cyclooxygenases](#) 1 and 2 through covalent bonding (via [acetylation](#) of a [serine](#) hydroxyl group near the [active site](#) of the enzyme). In this manner, it decreases the synthesis of [thromboxane A2](#) and [prostaglandins](#), thereby inhibiting [platelet aggregation](#). While it does provide some anti-thrombotic effect, it would not provide reliable long-term protection against embolic [stroke](#) and a different agent is indicated.

C - Interference with carboxylation of glutamate residues

Explanation Why

[Warfarin](#) is a commonly used medication for the long-term prevention of embolic [stroke](#) in patients with [Afib](#) and embolic [stroke risk factors](#) (e.g., elevated [CHADS-VASC](#) score). Like all [coumarins](#), [warfarin](#) inhibits hepatic [vitamin K epoxide reductase](#), thereby blocking synthesis (recycling) of the active, reduced form of [vitamin K](#). The result is decreased synthesis of [coagulation factors](#) II, VII, IX, and X, which require reduced [vitamin K](#) to ensure γ -[carboxylation](#) of their [glutamate](#) residues to function. This reduction of functioning [coagulation factors](#) gives [warfarin](#) its anticoagulatory effect.

D - Activation of the conversion of plasminogen to plasmin

Explanation Why

[Thrombolytics](#), including [streptokinase](#) and [tPA](#), activate the conversion of [plasminogen](#) to [plasmin](#), resulting in the cleavage of [thrombin](#) and [fibrin](#) clots. This class of drugs is indicated for the acute management of some thromboses (e.g., [massive PE](#), large vessel [ischemic stroke](#)) but is not used for the long-term mitigation of embolic [stroke](#) risk in patients with [Afib](#).

E - Irreversible blockade of adenosine diphosphate receptors

Explanation Why

[ADP receptor antagonists](#) like [clopidogrel](#) and [prasugrel](#) irreversibly block the [ADP](#) (P2Y₁₂) receptor on [platelets](#), thereby preventing the expression of [glycoproteins IIb/IIIa](#) on the [platelet](#) surface and inhibiting [platelet activation](#) and aggregation. Indications for [ADP receptor antagonists](#) include [acute coronary syndrome](#), coronary stenting, or as secondary prophylaxis following a thrombotic [stroke](#). A different agent is indicated for long-term embolic [stroke](#) prevention in patients with [Afib](#).

Question # 23

A 55-year-old man comes to the physician because of intermittent palpitations that occur when he is stressed, exercising, or when he drinks alcohol. Physical examination shows an irregularly irregular pulse. An ECG shows irregular QRS complexes without any discrete P waves. Pharmacotherapy with carvedilol is initiated for his condition. Compared to treatment with propranolol, which of the following adverse effects is most likely?

	Answer	Image
A	Hyperkalemia	
B	Hypotension	
C	Bradycardia	
D	Hyperglycemia	
E	Bronchospasm	

Hint

This patient presents with symptoms that indicate atrial fibrillation with rapid ventricular response (palpitations, irregular pulse, ECG showing irregular QRS complexes, and indiscernible P-waves). Carvedilol is a nonselective beta blocker with additional α -blocking action.

Correct Answer

A - Hyperkalemia

Explanation Why

[Hyperkalemia](#) can occur as a side effect of [beta blockers](#), especially in patients with pre-existing impaired potassium excretion or with [beta blocker overdose](#). It is caused by two different mechanisms: First, [β-adrenergic](#) blockade impairs urinary potassium excretion by inhibiting the secretion of [renin](#), which in turn decreases [aldosterone](#) synthesis. Second, [beta blockers](#) directly decrease cellular uptake of potassium by inhibiting [Na⁺/K⁺-ATPase](#). As both [carvedilol](#) and [propranolol](#) are non-selective [beta blockers](#), [hyperkalemia](#) is a possible side effect of both.

B - Hypotension

Explanation Why

As [beta blockers](#) are [class II antiarrhythmic drugs](#), [carvedilol](#) can be administered to lower [heart rate](#) by reducing AV conduction velocity. However, because of its α -blocking action, [carvedilol](#) also inhibits α_1 -adrenergic receptors of blood vessels, which leads to relaxation of [smooth muscle](#) and thus [vasodilation](#). This decreases [peripheral vascular resistance](#), which in turn reduces [preload](#) and [afterload](#) and might cause [orthostatic hypotension](#). This adverse effect is not observed in [propranolol](#), which does not have an α -blocking effect, but only blocks β_1 , β_2 , and β_3 receptors.

C - Bradycardia

Explanation Why

Inhibition of β_1 receptors on [cardiac muscle](#) tissue decreases [heart rate](#) and can lead to [bradycardia](#). However, this effect may be observed in all [beta blockers](#), including [carvedilol](#) and [propranolol](#), since they all block β_1 receptors.

D - Hyperglycemia

Explanation Why

Blockade of [\$\beta_2\$ receptors](#) in [pancreatic](#) cells impairs [insulin](#) release, which lowers cellular glucose uptake. Through this mechanism, non-selective [beta blockers](#) can cause [hyperglycemia](#) with [impaired glucose tolerance](#) or even new-onset [diabetes](#). As both [carvedilol](#) and [propranolol](#) are non-selective [beta blockers](#), [hyperglycemia](#) is a possible side effect of both.



E - Bronchospasm

Explanation Why

Non-selective [beta blockers](#) inhibit β_1 , β_2 , and β_3 receptors. Since [\$\beta_2\$ receptors](#) are found in bronchiolar [smooth muscle](#), these [beta blockers](#) can cause bronchospasm. As both [carvedilol](#) and [propranolol](#) are non-selective [beta blockers](#), bronchospasm is a possible side effect of both.

Question # 24

A 54-year-old man is brought to the emergency department 1 hour after the sudden onset of shortness of breath, severe chest pain, and sweating. He has hypertension and type 2 diabetes mellitus. He has smoked one pack and a half of cigarettes daily for 20 years. An ECG shows ST-segment elevations in leads II, III, and avF. The next hospital with a cardiac catheterization unit is more than 2 hours away. Reperfusion pharmacotherapy is initiated. Which of the following is the primary mechanism of action of this medication?

	Answer	Image
A	Blocking of adenosine diphosphate receptors	 The diagram illustrates the pathways of platelet activation and aggregation. On the left, under 'Activation', ADP is shown binding to the P2Y12 receptor (P2Y12R). This triggers a signal transduction cascade involving G-proteins (Gp12b/132aR) and the conversion of arachidonic acid (AA) to thromboxane A2 (TXA2) by the enzyme cyclooxygenase (COX). TXA2 then binds to its receptor, leading to 'Vasoconstriction'. On the right, under 'Aggregation', TXA2 and ADP are shown promoting platelet aggregation. This process involves the activation of glycoprotein IIb/IIIa receptors, which bind to fibrinogen, leading to the formation of a 'Platelet plug'. The diagram also shows 'Irreversible cyclooxygenase inhibitors' and 'Glycoprotein IIb/IIIa inhibitors' as points of intervention.
B	Conversion of plasminogen to plasmin	 The diagram illustrates the fibrinolytic system. It shows the conversion of Plasminogen to Plasmin. Plasminogen can be converted to Plasmin by tPA (tissue plasminogen activator), urokinase, or streptokinase. tPA is labeled as 'Present physiologically and used pharmacologically', while urokinase and streptokinase are labeled as 'Only used pharmacologically'. Plasminogen is also converted to Plasmin by Prokinase, which is formed from Prokallikrein and Kallikrein. Kallikrein is formed from XIIa. Plasmin is then used to degrade Fibrinogen into 'Fibrinogen degradation products' and to break down a 'Fibrin mesh' into 'Fibrin degradation products'. The diagram also shows 'PAI (plasminogen activator inhibitor)' and 'α2-Antiplasmin' as inhibitors of the system.

	Answer	Image
C	Potentialiation of antithrombin III action	The diagram illustrates the coagulation cascade. In the liver, uncarboxylated precursor factors II, VII, IX, and X are converted to carboxylated factors by γ-Glutamyl carboxylase, a process dependent on reduced vitamin K (active). Oxidized vitamin K (inactive) is recycled back to its active form by epoxide reductase, which is inhibited by Vitamin K antagonists. The extrinsic pathway involves Factor VII being converted to VIIa, which then activates Factor X to Xa. The intrinsic pathway involves Factor XII being converted to XI, then IX, and finally IXa, which also activates Factor X to Xa. Factor Xa, along with Factor V (converted to Va), converts Prothrombin II to Thrombin IIa. Thrombin IIa then converts Fibrinogen I to Fibrin Iα. Various inhibitors are shown: Direct oral factor Xa inhibitors block Xa; Direct oral thrombin inhibitors block Thrombin IIa; AT3 (Antithrombin III) blocks both Xa and Thrombin IIa; LMWH (Low molecular weight heparin), Heparinoid, and Synthetic heparin block Thrombin IIa.
D	Prevention of thromboxane formation	
E	Inhibition of glutamic acid residue carboxylation	This diagram is identical to the one in row C, showing the coagulation cascade and the liver's role in carboxylation of factors II, VII, IX, and X. It details the extrinsic pathway (VII to VIIa), intrinsic pathway (XII to XI to IX to IXa), and combined pathway (X to Xa). Xa converts Prothrombin II to Thrombin IIa, which then converts Fibrinogen I to Fibrin Iα. Inhibitors like AT3, LMWH, Heparinoid, and Synthetic heparin are shown blocking various steps.
F	Binding to glycoprotein receptor IIb/IIIa	The diagram illustrates the fibrinolytic system. Plasminogen is converted to Plasmin by tPA (tissue plasminogen activator), urokinase, or streptokinase. tPA is also inhibited by PAI-1 (plasminogen activator inhibitor). Plasmin then degrades Fibrinogen to Fibrin degradation products and Fibrin mesh. Fibrin mesh is further degraded by plasmin to Fibrin degradation products. The diagram also shows the role of PAI-1 (plasminogen activator inhibitor) and α2-Antiplasmin in regulating the system. A legend indicates that green boxes represent factors present physiologically and used pharmacologically, while purple boxes represent factors only used pharmacologically.

	Answer	Image
G	Direct inhibition of thrombin activity	The diagram illustrates the coagulation cascade and the vitamin K cycle. On the left, the Liver section shows the γ-Glutamyl carboxylase cycle: Uncarboxylated precursor factors II, VII, IX, X are converted to Carboxylated factors II, VII, IX, X using Reduced vitamin K (active). This process is inhibited by Vitamin K antagonists. Oxidized vitamin K (inactive) is converted back to the active form by Epoxide reductase. The Extrinsic pathway starts with VII converting to VIIa, which then activates X to Xa. The Intrinsic pathway starts with XII converting to XIIa, which activates XI to XIa, then IX to IXa, and finally IXa activates X to Xa. The Combined pathway shows Xa activating Prothrombin II to Thrombin IIa. Thrombin IIa then activates Fibrinogen I to Fibrin Ia. Direct oral factor Xa inhibitors block the activation of Xa. Direct oral thrombin inhibitors block the conversion of Prothrombin II to Thrombin IIa. LMWH, Heparinoid, Synthetic heparin and AT3 are shown as inhibitors of Thrombin IIa. UFH is shown as an inhibitor of Fibrinogen I.


Hint

This patient's ECG findings indicate an acute ST-elevation myocardial infarction (STEMI) of the inferior myocardial wall. Because the next percutaneous coronary intervention (PCI) facility is > 120 minutes away, he has received immediate fibrinolytic therapy (e.g., with alteplase).

Correct Answer

A - Blocking of adenosine diphosphate receptors

Image




Explanation Why

The [ADP](#) receptor inhibitors [clopidogrel](#), [prasugrel](#), and [ticagrelor](#) inhibit [platelet aggregation](#) by blocking the adenosine diphosphate receptors and preventing the [ADP](#)-mediated formation of the [GpIIb/IIIa receptor](#) complex. Based on this mechanism, [ADP](#) receptor inhibitors prevent coagulation but cannot lyse already existing clots to reperfuse blocked [coronary arteries](#). Patients with acute [STEMI](#) who undergo [PCI](#) and stent implantation should receive [ADP](#) receptor inhibitors and [aspirin](#) as dual antiplatelet therapy to decrease the likelihood of [stent thrombosis](#).

B - Conversion of plasminogen to plasmin

Image




Explanation Why

Fibrinolytic therapy (e.g., [alteplase](#)) cleaves [plasminogen](#) to form [plasmin](#), the main enzyme responsible for [fibrinolysis](#). It is indicated in early [MI](#) (symptom onset within the last 12 hours) if [PTCA](#) cannot be performed within 90–120 minutes, as is the case here, as long as there are no contraindications (e.g., recent major surgery or trauma, active bleeding, prior [intracranial hemorrhage](#)). [Fibrinolysis](#) is also used in early [ischemic stroke](#) (< 3 hours) and severe [PE](#). The main adverse effect of [fibrinolytics](#) is an increased risk of bleeding. In [MI](#), [reperfusion injury](#) of the [myocardium](#) may also result in [arrhythmias](#) following [thrombolysis](#).

C - Potentiation of antithrombin III action

Image



Explanation Why

[Heparin](#) potentiates the action of [antithrombin III](#), a [serine](#) protease inhibitor that rapidly inactivates factors IIa, IXa, Xa, XIa, and XIIa and prevents coagulation. However, [heparin](#) is not capable of lysing an already existing clot to reperfuse blocked [coronary arteries](#). Patients with a [STEMI](#) who receive [fibrinolytic therapy](#) are usually also given [unfractionated heparin](#) to prevent reinfarction.

D - Prevention of thromboxane formation


Explanation Why

[Aspirin](#) prevents the formation of thromboxane by irreversibly inhibiting [COX-1](#) in [platelets](#), thus lowering thromboxane synthesis and inhibiting [platelet aggregation](#). Application of [aspirin](#) is indicated in stable and [unstable angina pectoris](#) and in [acute myocardial infarction](#), such as a [STEMI](#),

to prevent further clot formation in the coronaries of these patients. However, it does not have thrombolytic effects and is not used for [thrombolysis](#) in patients with [STEMI](#).

E - Inhibition of glutamic acid residue carboxylation

Image




Explanation Why

[Vitamin K antagonists](#) (e.g., [warfarin](#)) inhibit [vitamin K epoxide reductase](#), which [carboxylates glutamic acid](#) residues on the coagulation [factors II](#), VII, IX, and X. This mechanism inhibits production of the [coagulation factors](#) and causes chronic anticoagulation. [Vitamin K antagonists](#) are used for [venous thromboembolism](#) prophylaxis or prevention of [stroke](#) in [atrial fibrillation](#). They have no thrombolytic effect and are not given for immediate [thrombolysis](#) in patients with [STEMI](#).

F - Binding to glycoprotein receptor IIb/IIIa

Image




Explanation Why

[Glycoprotein IIb/IIIa inhibitors](#) bind to the [glycoprotein IIb/IIIa](#) receptor and thereby prevent [fibrinogen](#) binding to [platelets](#) and subsequent [platelet aggregation](#). [Glycoprotein IIb/IIIa inhibitors](#) are indicated in high-risk patients with [unstable angina](#) or [non-ST-segment elevation myocardial infarction \(NSTEMI\)](#) who are scheduled to undergo [PCI](#) within the next 24 hours to prevent thrombotic complications. However, [glycoprotein IIb/IIIa inhibitors](#) are not used as [fibrinolytic therapy](#) in patients with [STEMI](#).

G - Direct inhibition of thrombin activity

Image



Explanation Why

[Direct thrombin inhibitors](#), such as [dabigatran](#) and [argatroban](#), bind to [thrombin](#) and so decrease its activity. This mechanism has an anticoagulative but not a thrombolytic effect. [Direct thrombin inhibitors](#) are not generally used in the treatment of [STEMI](#).

Question # 25

Several hours after vaginal delivery, a male newborn delivered at full-term develops tachycardia and tachypnea. His blood pressure is within normal limits. Pulse oximetry on room air shows an oxygen saturation of 79% in the right hand and 61% in the left foot. Physical examination shows bluish discoloration of the face and trunk, supraclavicular and intercostal retractions, and a machine-like murmur over the precordium. Bedside echocardiography shows pulmonary and systemic circulation are in parallel rather than in series. What is the most appropriate pharmacotherapy for this patient?

	Answer	Image
A	Sildenafil	
B	Alprostadil	
C	Metoprolol	
D	Indomethacin	
E	Dopamine	

Hint

The findings of cyanosis early in a neonate's life is concerning for a congenital heart defect. This patient's echocardiogram shows two separate circulations connected by a patent ductus arteriosus (machine-like murmur). This indicates transposition of the great vessels.

Correct Answer

A - Sildenafil

Explanation Why

[Sildenafil](#), a [phosphodiesterase type 5 inhibitor](#), is used to dilate pulmonary vessels in [neonates](#) with [pulmonary hypertension](#) until the underlying cause is addressed. In this patient, dilation of pulmonary vessels could decrease right to left shunting across the [PDA](#) and, thus, worsen his symptoms.

B - Alprostadil

Explanation Why

[Alprostadil](#) ([prostaglandin E1](#)) is used in [newborns](#) with congenital [heart](#) defects to maintain a [patent ductus arteriosus](#). In this [infant](#), who has [transposition of the great vessels](#), only deoxygenated blood is pumped from the [right ventricle](#) into the aorta. Therefore, it is important that oxygenated and deoxygenated blood are mixed through a shunt flow until definitive treatment (e.g., surgery) is performed. Other ductal-dependent [heart](#) defects include [tetralogy of Fallot](#), critical [pulmonic stenosis](#), and pulmonary atresia.

C - Metoprolol

Explanation Why

[Beta blockers](#), such as [propranolol](#) and [metoprolol](#), are used for the treatment of neonatal [hypertension](#) and congenital [long QT syndrome](#). In this patient, [tachycardia](#) is a compensatory mechanism and thus should not be treated with [beta blockers](#).

D - Indomethacin

Explanation Why

[Indomethacin](#), an inhibitor of [prostaglandin](#) synthesis, is used to close a [patent ductus arteriosus](#) in [premature infants](#). In patients with [transposition of the great vessels](#), survival depends on communication between the two parallel circulations. Hence, closure of the [patent ductus arteriosus](#) would have a deleterious effect in this [infant](#).

E - Dopamine

Explanation Why

[Dopamine](#) can be used to raise systemic blood pressure in [infants](#) with cardiogenic [shock](#). This [infant](#) does not need a vasopressor as he is not hypotensive.

Question # 26

A 41-year-old man comes to the physician because of a 10-day history of generalized weakness. He has also had headaches and night sweats and is a known user of illicit intravenous drugs. His temperature is 39.1°C (102.4°F), pulse is 110/min, respirations are 17/min, and blood pressure is 127/78 mm Hg. There are several track marks on his forearms. A holosystolic murmur that increases on inspiration is heard along the left sternal border. Laboratory studies show a leukocyte count of 13,900/mm³ and an erythrocyte sedimentation rate of 58 mm/h. Which of the following is the most likely consequence of this patient's condition?

	Answer	Image
A	Hematuria	
B	Painless macules and papules on palms and soles	
C	Rupture of nail-bed capillaries	
D	Retinal hemorrhages	

	Answer	Image
E	Pulmonary embolism	
F	Painful nodules on pads of the fingers	

Hint

The patient uses illicit IV drugs and presents with fever, tricuspid regurgitation, and leukocytosis. This combination is highly suggestive of acute right-sided infective endocarditis (IE).

Correct Answer

A - Hematuria

Explanation Why

[Hematuria](#) is caused by a wide range of conditions, including trauma, infections, [malignant tumors](#), and [ischemia](#). Renal damage in IE is primarily caused by two mechanisms: [Septic embolization](#) can lead to [renal artery](#) occlusion with subsequent [ischemia](#) and renal damage, while [immunoglobulin](#) and complement deposition in the [glomerulus](#) can lead to glomerulonephritis. Although a [septic embolism](#) can cause [renal injury](#) in IE, this complication is more common in left-sided involvement.

B - Painless macules and papules on palms and soles

Image



Explanation Why

In IE, painless [macules](#) and [papules](#) on the palms and soles are known as [Janeway lesions](#). These occur more commonly in left-sided IE than right-sided IE.

C - Rupture of nail-bed capillaries

Image



Explanation Why

[Splinter hemorrhages](#) are caused by the rupture of [capillaries](#) in the [nail](#) bed and are seen in various conditions, including trauma, IE, [antiphospholipid antibody syndrome](#), and [trichinosis](#). Patients with left-sided [endocarditis](#) are more likely to have [splinter hemorrhages](#) than those with [right-sided endocarditis](#).

D - Retinal hemorrhages

Explanation Why

Retinal hemorrhages occur in a variety of diseases, including [central retinal vein occlusion](#), [malignant hypertension](#), [retinal detachment](#), [subarachnoid hemorrhage](#), and IE (where they are known as [Roth spots](#)). However, peripheral manifestations such as retinal hemorrhages are not commonly observed in patients with right-sided IE.

E - Pulmonary embolism

Explanation Why

IV drug users such as this patient are at increased risk of IE because the use of nonsterile syringes can lead to [bacteremia](#). Bacterial [colonization](#) of the [heart valves](#) can lead to the formation of thrombi and subsequent [bacterial emboli](#). While right [heart](#) involvement is rare in IE, IV drug users are at an increased risk of tricuspid or multiple valvular involvement. [Thrombi](#) from the right [heart](#) most frequently dislodge into the [lung](#), which puts the patient at risk of [pulmonary embolism](#). Patients with left-sided ([aortic valve](#) or [mitral valve](#)) [endocarditis](#) more commonly present with emboli to the [retina](#) (i.e., [Roth spots](#)), extremities (e.g., [Janeway lesions](#), [Osler nodes](#), [splinter hemorrhages](#)), [kidney](#), brain, and [spleen](#).

F - Painful nodules on pads of the fingers

Image



Explanation Why

Painful nodules on the pads of the fingers (also known as [Osler nodes](#)), which are caused by the deposition of [immune complexes](#) in the [skin](#), are a rare but highly suggestive finding in IE. However, peripheral manifestations such as [Osler nodes](#) are not commonly observed in patients with right-sided [endocarditis](#). Moreover, [Osler nodes](#) typically occur in protracted cases of IE, whereas this patient presents with acute IE.

Question # 27

A 47-year-old woman comes to the physician because of a 6-week history of fatigue and low-grade fever. She has no history of serious illness except for a bicuspid aortic valve, diagnosed 10 years ago. She does not use illicit drugs. Her temperature is 37.7°C (99.9°F). Physical examination shows petechiae under the fingernails and multiple tender, red nodules on the fingers. A new grade 2/6 diastolic murmur is heard at the right second intercostal space. Which of the following is the most likely causal organism?

	Answer	Image
A	Enterococcus faecalis	
B	Staphylococcus epidermidis	
C	Streptococcus pyogenes	
D	Streptococcus sanguinis	
E	Streptococcus gallolyticus	
F	Staphylococcus aureus	

	Answer	Image
G	Candida albicans	
H	Streptococcus pneumoniae	

Hint

Osler nodes, subungual hemorrhage, and a new murmur in a patient with a 6-week history of fatigue and low-grade fever suggest subacute infective endocarditis. In patients with an underlying valve disease, infection is typically caused by a gram-positive, catalase-negative, coagulase-negative, α -hemolytic, optochin-resistant coccus.

Correct Answer

A - *Enterococcus faecalis*

Explanation Why

E. faecalis is a common cause of [subacute endocarditis](#) that is often associated with [nosocomial urinary tract infection](#) or manipulation of the genitourinary tract (e.g., [cystoscopy](#), abortion, delivery). However, this patient has not had a urinary infection or undergone any procedures, making *E. faecalis* unlikely.

B - *Staphylococcus epidermidis*

Explanation Why

S. epidermidis can cause [endocarditis](#), especially in patients with peripheral or [central venous catheters](#) and/or surgical implants and devices, where it forms a [biofilm](#) over the implanted material. In these cases and in patients who have undergone [prosthetic valve](#) surgery (< 60 days post-surgery), *S. epidermidis* is a common cause of [acute endocarditis](#). This patient, however, has a subacute course of [endocarditis](#) with symptoms that have persisted for more than 2 months, and she has had no invasive treatment.

C - *Streptococcus pyogenes*

Explanation But

Damage to the [heart valves](#) caused by [rheumatic heart disease](#) is a [risk factor](#) for [infective endocarditis](#).


Explanation Why

[Pharyngitis](#) or tonsillitis caused by *S. pyogenes* can result in [rheumatic fever](#), which can manifest with fatigue, low-grade [fever](#), a new [murmur](#), and nodules. However, these nodules are typically

nontender and located subcutaneously on extensor surfaces. In the case of [rheumatic fever](#), subungual hemorrhage would not be expected, whereas other classic signs of [rheumatic fever](#) (e.g., migratory polyarthritis, [erythema marginatum](#)) would be present. Moreover, [rheumatic fever](#) typically affects children and [adolescents](#), not adults.

D - Streptococcus sanguinis

Image



Explanation Why

S. sanguinis is a gram-positive member of the [viridans streptococci](#) group, which are oral [commensal](#) bacteria. Further [viridans streptococci](#) include *S. mutans* and *S. mitis*. [Viridans streptococci](#) are the most common cause of [subacute endocarditis](#) in predamaged native valves (such as this patient's [bicuspid aortic valve](#)) and [endocarditis](#) in [prosthetic valves](#) occurring > 60 days post-operation. [Viridans streptococci](#) are also a common cause of [endocarditis](#) after oral procedures, which frequently lead to transient [bacteremia](#). In patients with predamaged native valves, circulating bacteria may subsequently adhere to and infect [platelet-fibrin](#) aggregates that form at sites of endocardial injury, leading to vegetation formation.

E - Streptococcus gallolyticus

Explanation Why

S. gallolyticus is an uncommon cause of [subacute endocarditis](#) that is usually associated with [colon cancer](#). This patient does not have any features to suggest [colon cancer](#) (e.g., weight loss, [constipation](#), [GI bleeding](#)).

F - Staphylococcus aureus

Explanation Why

S. aureus is the most common cause of [acute bacterial endocarditis](#) in all patient groups. It progresses rapidly and often affects previously healthy valves. However, this patient has had a subacute course with symptoms occurring over the past 2 months. *S. aureus* is therefore unlikely to be the causative organism.

G - Candida albicans

Explanation Why

C. albicans can cause [endocarditis](#) in patients who have recently undergone cardiac surgery. [Immunocompromised](#) individuals and intravenous drug users are also at risk of *C. albicans* infection. This patient has none of these [risk factors](#).

H - Streptococcus pneumoniae

Explanation Why

While *S. pneumoniae* can cause [infective endocarditis](#), affected patients tend to have [risk factors](#) such as [cirrhosis](#) or [HIV](#) infection. This patient has no history of serious illness, making *S. pneumoniae* an unlikely cause of her [endocarditis](#).

Question # 28

A 55-year-old man comes to the emergency department because of left-sided chest pain and difficulty breathing for the past 30 minutes. His pulse is 88/min. He is pale and anxious. Serum studies show increased cardiac enzymes. An ECG shows ST-elevations in leads I, aVL, and V5-V6. A percutaneous coronary intervention is performed. In order to localize the site of the lesion, the catheter must pass through which of the following structures?

	Answer	Image
A	Posterior coronary sinus → left circumflex artery	
B	Left coronary artery → left circumflex artery	
C	Right coronary artery → right marginal artery	

	Answer	Image
D	Left coronary artery → posterior descending artery	
E	Aorta → right coronary artery	
F	Right coronary artery → posterior descending artery	
G	Left coronary artery → left anterior descending artery	


Hint

This patient presents with classic signs and symptoms of myocardial infarction (acute left-sided chest pain, dyspnea, pallor). His ECG suggests acute lateral wall infarction.

Correct Answer

A - Posterior coronary sinus → left circumflex artery

Image




Explanation Why

The [coronary sinus](#) is the venous drainage system of the [coronary veins](#). It cannot be used to gain access to [coronary arteries](#).

B - Left coronary artery → left circumflex artery

Image



Explanation Why

Acute [lateral](#) wall [infarction](#), which manifests with [ST elevations](#) in the [lateral](#) (I, aVL) and [lateral anterior precordial leads](#) (V5, V6), is most often caused by occlusion of the left circumflex [coronary artery](#) (LCX). The LCX supplies the [lateral](#) and [posterior](#) walls of the [left ventricle](#) as well as the anterolateral [papillary muscle](#). In [percutaneous coronary intervention](#), the catheter will need to pass through the aortic sinus and the [left coronary artery](#) before entering the LCX.

C - Right coronary artery → right marginal artery

Explanation Why

The [right marginal artery](#), which arises from the [right coronary artery](#), supplies the [anterior](#) and [posterior right ventricle](#). Occlusion of this [artery](#) causes [right ventricular infarction](#) and can lead to

acute [right heart failure](#). An [ECG](#) augmented by right [chest leads](#) (V_{4R}–V_{6R}) is necessary to confirm the diagnosis. This patient's [ECG](#) changes are consistent with [lateral](#) wall [infarction](#) caused by occlusion of a different vessel.

D - Left coronary artery → posterior descending artery

Explanation Why

The [left coronary artery](#) gives rise to the [posterior descending artery](#) in about 10–15% of individuals. The [PDA](#) supplies the [posterior](#) walls of the ventricles, the [posterior](#) third of the interventricular septum, the posteromedial [papillary muscle](#), and the [AV node](#). Occlusion of this [artery](#) causes [posterior myocardial infarction](#) and may lead to [bradycardia](#) or [heart block](#) due to [AV node](#) dysfunction. The [posterior myocardium](#) is not directly visualized by standard 12-lead [ECG](#), but reciprocal changes such as [ST depression](#) and pathological R waves can be seen in the anteroseptal leads V₁–V₃. Extended [precordial leads](#) (V₇–V₉) added to the standard leads will demonstrate [ST elevations](#). This patient's [lateral](#) wall [infarction](#) is caused by occlusion of a different vessel.

E - Aorta → right coronary artery

Explanation Why

The [coronary arteries](#) leave the aorta just above the [aortic valve](#) in the aortic sinus. The [anterior](#) aortic sinus gives rise to the [right coronary artery](#) ([RCA](#)), which in most people (~ 80%) is the more dominant of the two major [coronary arteries](#), supplying the [right atrium](#) and ventricle, parts of the [left ventricle](#), the sinus node (SA), and the [AV node](#). Occlusion of this [artery](#) causes inferior [myocardial infarction](#) and may lead to [bradycardia](#) or [heart block](#) due to SA or [AV node](#) dysfunction. An [ECG](#) would show [ST elevations](#) in leads II, III, and aVF. This patient's [lateral](#) wall [infarction](#) is caused by occlusion of a different vessel.

F - Right coronary artery → posterior descending artery

Explanation Why

The [right coronary artery](#) gives rise to the [posterior descending artery](#) ([PDA](#)) in ~ 80% of

individuals. The [PDA](#) supplies the [posterior](#) walls of the ventricles, the [posterior](#) third of the interventricular septum, the posteromedial [papillary muscle](#), and the [AV node](#). Occlusion of this [artery](#) causes [posterior myocardial infarction](#) and may lead to [bradycardia](#) or [heart block](#) due to [AV node](#) dysfunction. The [posterior myocardium](#) is not directly visualized by standard 12-lead [ECG](#), but reciprocal changes such as [ST depression](#) and pathological R waves can be seen in the anteroseptal leads V₁–V₃. Extended [precordial leads](#) (V₇–V₉) added to the standard leads will demonstrate [ST elevations](#). This patient's [lateral](#) wall [infarction](#) is caused by occlusion of a different vessel.

G - Left coronary artery → left anterior descending artery

Explanation Why

The [left anterior descending artery](#) (LAD), which arises from the [left coronary artery](#), supplies the [anterior](#) surface of the [left ventricle](#), the anterolateral [papillary muscle](#), and the [anterior](#) $\frac{2}{3}$ of the ventricular septum. Occlusion of this [artery](#) causes [anterior](#) wall [myocardial infarction](#), leading to [ST elevations](#) in leads I, aVL, and [anterior](#) wall leads V₁–V₆. [Myocardial infarction](#) (MI) due to occlusion of the LAD is the most common type of [MI](#) and has a poor prognosis owing to the large volume of [myocardium](#) supplied by this vessel. This patient's [lateral](#) wall [infarction](#) is caused by occlusion of a different vessel.

Question # 29

An 80-year-old man is admitted to the hospital after the sudden onset of sub-sternal chest pain and shortness of breath while sitting in a chair. He has hypertension and type 2 diabetes mellitus. He has smoked 1 pack of cigarettes daily for 42 years. Four days after admission, he becomes tachycardic and then loses consciousness; the cardiac monitor shows irregular electrical activity. Cardiac examination shows a new systolic murmur at the apex. Despite appropriate measures, he dies. Microscopic evaluation of the myocardium is most likely to show which of the following?

	Answer	Image
A	Coagulative necrosis with dense neutrophilic infiltrate	
B	Wavy myocardial fibers without inflammatory cells	
C	Low cellularity with dense, non-contractile scar tissue	
D	Dense granulation tissue with collagenous scar formation	
E	Hyperemic granulation tissue with abundance of macrophages	

Hint

This patient's murmur is consistent with acute mitral regurgitation and in the setting of recent chest pain, is most likely due to papillary muscle rupture from myocardial infarction. Other complications that can occur 4 days after myocardial infarction include free wall rupture and interventricular septal rupture.

Correct Answer

A - Coagulative necrosis with dense neutrophilic infiltrate

Explanation Why

[Coagulative necrosis](#) with dense neutrophilic infiltrate is typically seen 1–3 days post-[MI](#). The most important complication during this period is post-[infarction](#) fibrinous pericarditis.

B - Wavy myocardial fibers without inflammatory cells

Explanation Why

Wavy [myocardial](#) fibers without inflammatory cells are typically seen within the first 4 hours post-[MI](#) and are followed by [coagulative necrosis](#) (4-72 hours post-[MI](#)). Common complications during this period include [ventricular arrhythmia](#), [heart failure](#), and [cardiogenic shock](#).

C - Low cellularity with dense, non-contractile scar tissue

Explanation Why

Low cellularity with dense, non-contractile [scar](#) tissue is typically seen 2 months post-[MI](#). Common complications during this period include [heart failure](#), [arrhythmia](#), true [ventricular aneurysm](#), and [Dressler syndrome](#).

D - Dense granulation tissue with collagenous scar formation

Explanation Why

Dense [granulation tissue](#) with collagenous [scar](#) formation is typically seen 2–8 weeks post-[MI](#). Common complications during this period include true [ventricular aneurysm](#) and [Dressler syndrome](#).

E - Hyperemic granulation tissue with abundance of macrophages

Explanation Why

Hyperemic [granulation tissue](#) with abundance of [macrophages](#) is typically seen 3–10 days post [myocardial infarction](#). The complications during this period usually arise due to [macrophage](#)-mediated removal of [necrotic](#) tissue. They include free wall rupture, left ventricular pseudoaneurysm, [papillary muscle rupture](#), and interventricular septal rupture.

Question # 30

A previously healthy 61-year-old man comes to the physician because of a 6-month history of morning headaches. He also has fatigue and trouble concentrating on his daily tasks at work. He sleeps for 8 hours every night; his wife reports that he sometimes stops breathing for a few seconds while sleeping. His pulse is 71/min and blood pressure is 158/96 mm Hg. He is 178 cm (5 ft 10 in) tall and weighs 100 kg (220 lb); BMI is 31.6 kg/m². Which of the following is the most likely cause of this patient's hypertension?

	Answer	Image
A	Nocturnal upper airway obstruction	
B	Hypophyseal neoplasm	
C	Hypersecretion of aldosterone	
D	Overproduction of cortisol	
E	Proliferation of adrenal chromaffin cells	
F	Low circulating free thyroxine levels	

Hint

Polysomnography will confirm the diagnosis.

Correct Answer

A - Nocturnal upper airway obstruction

Explanation But

[Obesity](#), especially around the neck, is the most important [risk factor](#) for developing [OSA](#), and weight loss will often improve symptoms.

Explanation Why

Nocturnal upper [airway](#) obstruction resulting from the collapse of the pharyngeal muscles is the pathological mechanism underlying [obstructive sleep apnea](#). Typical features of this disease include snoring, morning [headaches](#), daytime fatigue, and poor concentration. Individuals with [OSA](#) are hypercapnic due to increased retention of carbon dioxide (a consequence of the [apneic episodes](#)); this, in turn, causes an increased [sympathetic](#) response that leads to [secondary hypertension](#).

B - Hypophyseal neoplasm

Explanation Why

Neoplasms of the [hypophysis](#), such as pituitary [adenomas](#), can cause [mass effect](#) leading to [headaches](#) and visual disturbances. Increased secretion of [ACTH](#) ([Cushing disease](#)), [growth hormone](#) ([acromegaly](#)), or [TSH](#) (central [hyperthyroidism](#)) could also cause increased blood pressure. However, [pituitary adenomas](#) usually cause [visual field](#) defects, such as [bitemporal hemianopsia](#), and it would not explain this patient's nocturnal breathing problems.

C - Hypersecretion of aldosterone

Explanation Why

Hypersecretion of [aldosterone](#), seen in primary hyperaldosteronism, can present with [hypertension](#) and [headaches](#). High systemic [aldosterone](#) levels result in increased sodium reabsorption and

potassium secretion in the [kidney](#). This, in turn, leads to the retention of water along with sodium, causing [hypertension](#) and [hypokalemia](#). However, this patient lacks other common findings of [hypokalemia](#), like muscle weakness, cramping, and [polyuria/polydipsia](#), making this diagnosis less likely. [Primary hyperaldosteronism](#) would also not account for his nocturnal breathing problems.

D - Overproduction of cortisol

Explanation Why

[Cushing syndrome](#), due to endogenous overproduction of [cortisol](#) by the [adrenal gland](#), could explain this patient's [hypertension](#), [obesity](#), and poor concentration. However, patients with [Cushing syndrome](#) typically have moon facies, buffalo hump, and abdominal striae, all of which are side effects of [systemic glucocorticoid](#) excess and are absent in this patient.

E - Proliferation of adrenal chromaffin cells

Explanation Why

A [proliferation](#) of the [adrenal chromaffin cells](#) (e.g., [pheochromocytoma](#)) could explain this patient's persistent [headaches](#) and [hypertension](#). This [tumor](#) can present with sustained [hypertension](#), but the [headaches](#) are often paroxysmal and accompanied by significant sweating, [tachycardia](#), and pallor, making this diagnosis less likely. It also would not account for his nocturnal breathing problems.

F - Low circulating free thyroxine levels

Explanation Why

Low circulating free [thyroxine](#) levels, seen in [hypothyroidism](#), could explain this patient's impaired [cognition](#), [hypertension](#), and [obesity](#). However, this patient lacks the more salient features of [hypothyroidism](#), such as cold intolerance, [constipation](#), and [skin](#) changes (cold, dry [skin](#) and [hair](#) loss), making this diagnosis unlikely. It also would not explain his nocturnal breathing problems.

Question # 31

A 45-year-old woman comes to the emergency department for fatigue and swelling in her legs. She has had a 10-kg (22-lb) weight loss in the past 8 months despite normal appetite. She has no history of serious medical illness, and has not seen a physician in over 15 years. Physical examination shows a brown-black, asymmetrical macule on her left shoulder. Cardiac examination shows jugular venous distention and an S₃ gallop best heard over the apex. Pitting edema is present in the bilateral lower extremities up to the mid-shin. A transthoracic echocardiogram shows a nonmobile mass protruding from the interatrial septum into the right atrium that partially occludes the tricuspid valve. A biopsy of the mass is most likely to show which of the following?

	Answer	Image
A	Nonencapsulated groups of well-differentiated skeletal muscle cells	
B	Clusters of bland cells without mitotic activity	
C	Sheets of large, atypical B cells with prominent nucleoli	
D	Mixture of cells from different germinal layers	
E	Nests of dysplastic melanocytes	

Hint

This patient has evidence of right-sided heart failure (JVD, S₃ heart sound, pitting edema) due to a right atrial mass that is most likely causing obstruction of the tricuspid valve. Cardiac tumors are most commonly seen in the setting of metastatic disease.

Correct Answer

A - Nonencapsulated groups of well-differentiated skeletal muscle cells

Explanation Why

Nonencapsulated groups of well-differentiated [skeletal muscle](#) cells are observed in biopsy samples of [rhabdomyoma](#). These are the most common primary cardiac tumors in children but are significantly less common in adults. The majority of intracardiac [rhabdomyomas](#) arise in association with [tuberous sclerosis](#), of which this patient has no clinical manifestations.

B - Clusters of bland cells without mitotic activity

Explanation Why

Clusters of bland cells without [mitotic](#) activity suggest [cardiac myxomas](#), the most common primary cardiac tumors. Histologic findings include mesenchymal cells within a mucoid, gelatinous stroma. [Cardiac myxomas](#) can cause systemic symptoms such as fatigue and weight loss (due to production of [interleukin 6](#)), as seen here. However, the vast majority of these tumors occur as a mobile, pedunculated mass in the [left atrium](#), rather than the right. Typically, [cardiac myxomas](#) manifest with symptoms of obstruction of blood flow through the [heart](#) (e.g., [syncope](#), [dyspnea](#), [palpitations](#)), a mid-diastolic [murmur](#) (similar to [mitral stenosis](#)), and/or a [tumor](#) plop on auscultation. Furthermore, primary cardiac neoplasms (e.g., [cardiac myxoma](#)) are not the most likely cause of cardiac tumors.

C - Sheets of large, atypical B cells with prominent nucleoli

Explanation Why

Sheets of large, atypical [B cells](#) with prominent nucleoli are consistent with [non-Hodgkin lymphoma](#). Lymphomatous lesions can arise in cardiac tissue, but these are quite rare, accounting for ~ 1% of cardiac tumors. This patient's weight loss is suspicious for cancer; however, a diagnosis other than [non-Hodgkin lymphoma](#) is more likely.

D - Mixture of cells from different germinal layers

Explanation Why

A mixture of cells from different germinal layers is the characteristic finding of a [teratoma](#). Cardiac involvement in [teratoma](#) cases is exceptionally rare, and, when encountered, seen most often in children. Moreover, [teratomas](#) tend to develop from the [pericardium](#) or the [mediastinum](#) rather than the endocardial lining. This patient's [echocardiogram](#) has identified an intracardiac neoplasm, which makes a [teratoma](#) very unlikely.


E - Nests of dysplastic melanocytes

Explanation Why

Nests of [dysplastic melanocytes](#) are seen in [metastatic melanoma](#). Cardiac tumors can be classified as primary or secondary ([metastatic](#)). [Metastatic](#) tumors are approximately 20–30 times more common than primary cardiac tumors and more commonly occur on the right side of the [heart](#). This patient's history of weight loss in combination with the asymmetrical mole raises suspicion of an occult [melanoma](#) that has [metastasized](#) to the [heart](#). [Melanoma](#) is the most frequently implicated cancer, followed by [lung](#) and [breast cancer](#).

Question # 32

A 40-year-old man comes to the physician because of a 1-week history of palpitations and dizziness. His pulse is 65/min and regular. An ECG is shown. A drug is administered that inhibits non-dihydropyridine calcium channels in the heart and his symptoms improve. The drug administered to the patient most likely has which of the following effects on the cardiac conduction system?



	Answer	Image
A	Prolongation of Purkinje fiber refractory period	
B	Decrease in bundle of His refractory period	
C	Delay in ventricular myocardial repolarization	

	Answer	Image
D	Decrease in ventricular myocardial action potential duration	
E	Prolongation of AV node repolarization	

Hint

This patient's symptoms and ECG findings (narrow QRS complexes with an atrial rate of 240/min in a sawtooth pattern) are most consistent with atrial flutter. Verapamil and diltiazem are non-dihydropyridine calcium channel blockers that are commonly used as first-line therapy for atrial flutter.

Correct Answer

A - Prolongation of Purkinje fiber refractory period

Explanation Why

Prolongation of the [Purkinje fiber refractory period](#) can be seen in patients taking [class IA antiarrhythmic drugs](#) and [class III antiarrhythmics](#). Calcium-channel blockers have little or no direct effect on ventricular or His-Purkinje tract conduction or refractoriness and rely on a different effect on the [cardiac conduction system](#) for the treatment of [arrhythmias](#).

B - Decrease in bundle of His refractory period

Explanation Why

A decrease in the [bundle of His refractory period](#) would not be expected with the administration of a drug that inhibits non-dihydropyridine calcium channels. [Calcium-channel blockers](#) have very little (if any) effect on ventricular or His-Purkinje tract conduction velocity or refractoriness.

C - Delay in ventricular myocardial repolarization

Explanation Why

Prolongation of ventricular [myocyte repolarization](#) can be caused by [class III antiarrhythmic drugs](#), which act primarily by blocking cardiac potassium channels. This prolonged [repolarization](#) manifests as a prolonged [QT interval](#) on [ECG](#). [Class III antiarrhythmic drugs](#) may also be used for [atrial flutter](#), as well as for [atrial fibrillation](#) and [ventricular tachycardias](#). However, [calcium channel blockers](#) rely on a different effect on the [cardiac conduction system](#) to prevent [arrhythmias](#).

D - Decrease in ventricular myocardial action potential duration

Explanation Why

A decrease in [myocardial action potential](#) duration is seen in patients taking [class IB antiarrhythmic drugs](#) like [lidocaine](#) or [mexiletine](#). These drugs have a weak blockade effect on [voltage-gated sodium channels](#) and are able to decrease [action potential](#) duration. Because agents in this class have minimal [vagal](#), nodal, and atrial effects, they would not be effective options for decreasing sinoatrial or [atrioventricular node automaticity](#). They are most effective at treating [ventricular tachyarrhythmias](#).


E - Prolongation of AV node repolarization

Explanation Why

Prolongation of [AV node repolarization](#) is seen in patients taking [class IV antiarrhythmic drugs](#) like [verapamil](#) and [diltiazem](#). Because the [depolarization](#) of nodal tissues (e.g., sinoatrial and [atrioventricular nodes](#)) is dependent upon the movement of [calcium ions](#), [non-dihydropyridine calcium channel blockers](#) primarily affect these tissues. The end result is prolonged [AV node](#) conduction time (decrease in conduction velocity) and [refractory period](#) (an increase in [effective refractory period](#)). This unique mechanism allows [class IV antiarrhythmic drugs](#) to effectively decrease SA/[AV node automaticity](#) in [arrhythmias](#) like [atrial flutter](#), [atrial fibrillation](#), and [supraventricular tachycardias](#). [Class IV antiarrhythmic drugs](#) also increase the [PR interval](#).

Question # 33

An investigator is studying the effects of different drugs on the contraction of cardiomyocytes. The myocytes are able to achieve maximal contractility with the administration of drug A. The administration of drug B produces the response depicted in the graph shown. Which of the following drugs is most likely to produce a response similar to that of drug B?



	Answer	Image
A	Albuterol	
B	Propranolol	
C	Pindolol	

	Answer	Image
D	Phenoxybenzamine	
E	Isoproterenol	

Hint

Cardiomyocyte contractility is mediated by beta-1 adrenergic receptors. The graph shows that drug B has lower efficacy than drug A.

Correct Answer

A - Albuterol

Explanation Why

[Albuterol](#) is a full [beta adrenergic receptor agonist](#) that has a higher affinity for beta-2, as opposed to beta-1, receptors. Despite the greater affinity for beta-2, this agent is also a full [agonist](#) for beta-1 receptors and would therefore be expected to follow the curve of drug A in the graph above as opposed to curve of drug B.

B - Propranolol

Explanation Why

[Propranolol](#) is a reversible [competitive antagonist](#) of [beta adrenergic receptors](#). As a reversible [competitive antagonist](#), it inhibits the effect of a [beta adrenergic agonist](#), requiring a higher concentration of that [agonist](#) to achieve a given efficacy. In other words, a [competitive antagonist](#) reduces the [potency](#) of a drug but has no effect on the efficacy. This would be reflected by shifting the curve on the graph to the right.

C - Pindolol

Explanation Why

[Pindolol](#) is a [partial agonist](#) of beta-1 [adrenergic receptors](#). Although partial [agonists](#) (drug B) typically have the same effect as complete [agonists](#) (drug A), they have a lower efficacy (a lower maximal drug effect). [Pindolol](#) would be consistent with drug B in the graph, as the plateau of maximal effect is lower for drug B than for drug A.

D - Phenoxybenzamine

Explanation Why

[Phenoxybenzamine](#) is a [noncompetitive antagonist](#) of [alpha adrenergic receptors](#). Although administration of a [noncompetitive antagonist](#) does reduce the efficacy of a given [agonist](#), this study is evaluating the effects of drugs on the contraction of [cardiomyocytes](#). [Alpha receptors](#) do not play a significant role in [cardiac muscle](#) contraction.

E - Isoproterenol

Explanation Why

[Isoproterenol](#) is a full [agonist](#) of [beta adrenergic receptors](#). Administration of this drug would correspond to the tracing for drug A, not drug B.

Question # 34

A 29-year-old woman, gravida 1, para 0 at 11 weeks' gestation comes to the physician because of a 2-day history of left lower extremity pain and swelling. Her temperature is 37.9°C (100.2°F). Physical examination shows a tender, palpable cord on the lateral aspect of the left lower leg. The overlying skin is erythematous and indurated. Duplex ultrasound shows vascular wall thickening and subcutaneous edema. Which of the following is the most likely diagnosis?

	Answer	Image
A	Deep vein thrombosis	
B	Erysipelas	

	Answer	Image
C	Erythema nodosum	
D	Varicose vein	
E	Lymphedema	
F	Superficial thrombophlebitis	

Hint

A variant of this patient's condition is Trousseau syndrome, in which palpable, tender cords recur at different locations on the body.

Correct Answer

A - Deep vein thrombosis

Image



Explanation Why

[DVT](#) manifests with diffuse swelling, tenderness, and redness of the affected extremity. There is a higher risk of [DVT](#) during [pregnancy](#) because of an increase in coagulability and [venous stasis](#). However, [DVT](#) would not show signs of [inflammation](#) (a tender, palpable cord with local [erythema](#) and induration or elevated temperature), and [duplex ultrasound](#) would not show vascular wall thickening or subcutaneous [edema](#).

B - Erysipelas

Image



Explanation Why

Erysipelas is a superficial skin infection typically caused by group A streptococci (*S. pyogenes*). Patients present with fever and localized, tender streaks on the skin. However, these streaks manifest as sharply-demarcated, bright-red patches, and not as a palpable cord. Additionally, duplex ultrasound might show inflammation of lymphatic vessels but not vascular wall thickening.

C - Erythema nodosum

Image



Explanation Why

[Erythema nodosum](#) is an [inflammation](#) of [subcutaneous fat](#) leading to the formation of painful nodules on the lower extremities and possibly an elevated temperature. It can occur during [pregnancy](#), likely due to a [delayed-type hypersensitivity reaction](#). However, [erythema nodosum](#) usually manifests with [skin](#) nodules on the shins, not with a palpable cord on the [lateral](#) aspect of the left lower leg. Additionally, [duplex ultrasound](#) would not show vascular wall thickening.

D - Varicose vein

Explanation Why

[Varicose veins](#) are common during [pregnancy](#) because the growing [uterus](#) places increased pressure on pelvic [veins](#). However, they manifest as dark purple or blue, twisted, bulging, soft, nontender

cords and usually show no signs of [inflammation](#). Additionally, [duplex ultrasound](#) might show subcutaneous [edema](#) but not vascular wall thickening.

E - Lymphedema

Image



Explanation Why

[Lymphedema](#) is a type of [nonpitting edema](#) caused by insufficient [lymphatic drainage](#) that occurs more commonly during [pregnancy](#) ([pregnancy hormones](#) induce changes in vascular permeability, promoting extravasation of fluids). [Lymphedema](#) is typically diffusely localized over the toes and feet; it would not manifest as a tender, palpable cord on the [lateral](#) aspect of the left lower leg. Additionally, [duplex ultrasound](#) would show subcutaneous [edema](#) but not vascular wall thickening.

F - Superficial thrombophlebitis

Explanation Why

[Superficial thrombophlebitis](#) is usually a benign [inflammation](#) of [veins](#) and overlying tissue due to thrombosis. The inflamed [veins](#), impacted with thrombotic material, manifest as palpable, nodular cords that remain visible when the limb is raised. There is a greater risk of thrombosis during [pregnancy](#) due to a combination of [hypercoagulability](#) (increased circulating clotting factors and [RBCs](#)) and elevated [venous stasis](#) (compression of enlarged [uterus](#) on pelvic [veins](#)).

Question # 35

An investigator is studying cardiovascular changes in athletes during exercise. Participants run on a treadmill at low, moderate, and high intensity in 5-minute intervals. After each interval, an echocardiogram is performed and blood samples are collected. Which of the following sets of findings would be present in an athlete following a moderate intensity interval?

	Cardiac output	PaO ₂	PaCO ₂	Venous blood pH
A	Decreased	normal	normal	normal
B	Increased	normal	normal	decreased
C	Decreased	decreased	decreased	decreased
D	Normal	normal	decreased	increased
E	Increased	decreased	decreased	increased

	Answer	Image
A	A	
B	B	
C	C	
D	D	
E	E	

Hint

During exercise, the body experiences physiological changes as its demands for energy and oxygen increase.

Correct Answer

A - A

Explanation Why

Decreased [cardiac output](#) with normal [PaO₂](#), [PaCO₂](#), and venous [blood pH](#) can be seen in the case of [systolic heart failure](#) without [pulmonary edema](#) but not in an exercising individual.

B - B

Explanation Why

In exercising individuals, [cardiac output](#) increases due to an elevated [stroke volume](#) and [heart rate](#). The depth and rate of respiration also increase, but [PaCO₂](#) and [PaO₂](#) remain normal because the increase in oxygenation and CO₂ washout is offset by higher O₂ consumption and CO₂ production in the exercising muscles. Therefore, the venous blood of exercising patients will have low O₂ tension and high CO₂ tension. The pH of the venous blood also decreases as a result of the high CO₂ tension and [lactic acid](#) production.

C - C

Explanation Why

A decrease in [cardiac output](#), [PaO₂](#), [PaCO₂](#), and venous [blood pH](#) is characteristic of [massive pulmonary embolism](#). In affected patients, the large [pulmonary embolus](#) obstructs pulmonary blood flow and thereby decreases [cardiac output](#). [Hyperventilation](#) caused by the hypoxic ventilatory drive results in increased CO₂ washout. Finally, the pH of venous blood decreases as a result of [lactic acidosis](#) induced by poor tissue perfusion. A different set of findings would be expected in an exercising individual.

D - D

Explanation Why

Decreased [PaCO₂](#) and increased venous [blood pH](#) with normal [cardiac output](#) and [PaO₂](#) can be seen in cases of voluntary [hyperventilation](#) in resting individuals but not in an exercising individual.


E - E

Explanation Why



Increased [cardiac output](#) with decreased [PaO₂](#), decreased [PaCO₂](#), and increased venous [blood pH](#) can be seen in patients with [altitude sickness](#). In such patients, [hyperventilation](#) caused by the hypoxic ventilatory drive results in increased CO₂ washout and [respiratory alkalosis](#). Stimulation of pacemaker cells and carotid [chemoreceptors](#) by [hypoxemia](#) increases the [heart rate](#) and thereby increases [cardiac output](#). A different set of findings would be expected in an exercising individual.



Question # 36

A 56-year-old man comes to the emergency department because of nausea and shortness of breath that started while he was at work one hour ago. He also reports lightheadedness that started earlier in the day. He has type 2 diabetes mellitus, for which he takes metformin and sitagliptin. He has smoked 1 pack of cigarettes daily for 20 years. He appears pale and diaphoretic. A 12-lead ECG is shown. Which of the following is the most likely underlying cause of this patient's symptoms?



	Answer	Image
A	Acute posterior myocardial infarction	

	Answer	Image
B	Viral myocarditis	
C	Ventricular fibrillation	Courtesy of Jason E. Reodiger, CCT, CRAT 
D	Acute pericarditis	
E	Acute inferior myocardial infarction	
F	Atrial fibrillation	

	Answer	Image
G	Saddle pulmonary embolus	 The ECG displays 12 leads: I, II, III, aVR, aVL, aVF, V1, V2, V3, V4, V5, and V6. The rhythm is regular. The most notable findings are in the inferior leads (II, III, and aVF), where the T waves are deeply inverted and have a characteristic 'M' shape, which is highly suggestive of a saddle pulmonary embolus. There is also a slight ST-segment depression in these leads.
H	Second-degree AV block	 The ECG shows a regular sinus rhythm with a PR interval of approximately 0.20 seconds. There are four P waves visible. The second, fourth, and sixth P waves are followed by a QRS complex, while the third and fifth P waves are not followed by a QRS complex, indicating dropped QRS complexes. This pattern is characteristic of second-degree AV block (Mobitz II).


Hint

The ECG shows ST elevations in II, III, and aV_F.

Correct Answer

A - Acute posterior myocardial infarction

Image



Explanation Why

An acute [posterior myocardial infarction](#) can be diagnosed based on the presence of [ST elevations](#) on special leads V7–V9. On a standard 12-lead [ECG](#), a [posterior MI](#) could manifest with reciprocal [ST depression](#) in leads V1–V4 or an R:S wave [ratio](#) > 1 in leads V1 and V2 due to an increase in the R-wave amplitude. This patient only has an [ST depression](#) in lead V2 and an R:S [ratio](#) < 1 in leads V1 and V2. It is therefore unlikely that this patient has had a [posterior wall MI](#).

B - Viral myocarditis

Explanation Why

Viral [myocarditis](#) can cause [dyspnea](#) and nonspecific [ST elevations](#) on [ECG](#), but it would typically be preceded by a viral [prodrome](#), with [fever](#), myalgia, and upper respiratory symptoms. This patient has no history of a preceding viral infection, making viral [myocarditis](#) an unlikely diagnosis. Furthermore, his [ECG](#) findings are more consistent with a different diagnosis.

C - Ventricular fibrillation

Image




Explanation Why

A patient with [ventricular fibrillation](#) would be unresponsive, with an [ECG](#) showing disorganized, high-frequency (usually > 300/min), sawtooth-like waves with indiscernible [p waves](#) or [QRS complexes](#). This patient is responsive, with distinct [P waves](#) as well as [QRS complexes](#) and a [heart](#)

[rate](#) of 60–70/min on [ECG](#).

D - Acute pericarditis

Image



Explanation Why

[Acute pericarditis](#) can cause [shortness of breath](#) and [ST elevations](#) on [ECG](#). However, the [ST elevations](#) in [pericarditis](#) typically occur diffusely in all leads, in contrast to this patient's [ECG](#) findings. Moreover, PR segment depression and clinical features such as [fever](#), a [pericardial friction rub](#), and/or retrosternal [chest pain](#) would be expected in a patient with [acute pericarditis](#).

E - Acute inferior myocardial infarction

Explanation Why

Acute inferior [myocardial infarction](#) (MI) is the most likely diagnosis in a patient with [dyspnea](#),

[lightheadedness](#), [ST elevation](#) in the [inferior leads](#) (II, III, aVF), and reciprocal [ST depression](#) in leads I and aVL. This patient has significant [risk factors](#) for atherosclerotic disease, including type 2 [diabetes mellitus](#) and a 20 [pack-year](#) smoking history. Acute [STEMI](#) is usually a clinical diagnosis based on the initial [ECG](#) and the presence of typical [ischemic chest pain](#). However, diabetic patients and elderly women may have an atypical presentation with little or no [chest pain](#) and predominantly autonomic symptoms (e.g., nausea).


F - Atrial fibrillation

Explanation Why

[Atrial fibrillation](#) can manifest with nausea, [shortness of breath](#), and light-headedness. However, in the case of [atrial fibrillation](#), the [ECG](#) would have revealed a variable RR interval and absent [P waves](#). This patient has a constant RR interval and clearly discernible [P waves](#) on [ECG](#), making [atrial fibrillation](#) unlikely.

G - Saddle pulmonary embolus

Image




Explanation Why

A saddle [pulmonary embolism](#) can manifest with [shortness of breath](#), light-headedness, and nonspecific [ST elevations](#) on [ECG](#). However, this patient is lacking features of [right ventricular strain](#) on [ECG](#) (e.g., S1Q3T3 pattern, increased R-wave amplitude and [ST depression](#) in V1 and V2) and the absence of [tachycardia](#) and major [risk factors](#) (e.g., prolonged immobilization, recent trauma, surgery, [malignancy](#)) make this diagnosis unlikely. This patient's [ECG](#) findings are more consistent with a different etiology.

H - Second-degree AV block

Image



Explanation Why

A [second-degree AV block](#) can manifest as either a [Mobitz type 1](#) or a [Mobitz type 2](#) block. [Mobitz type 1](#) manifests with a progressive prolongation of the [PR interval](#) that culminates in a non-conducted [P wave](#) (“dropped beat”) on [ECG](#). [Mobitz type 2 AV block](#) is characterized by a constant [PR interval](#) and a single or intermittent non-conducted [P wave](#) without subsequent [QRS complexes](#). This patient's [ECG](#) shows [P waves](#) before every [QRS complex](#) and a constant [PR interval](#).

Question # 37

A 39-year-old woman is brought to the emergency department 30 minutes after her husband found her unconscious on the living room floor. She does not report having experienced light-headedness, nausea, sweating, or visual disturbance before losing consciousness. Three weeks ago, she was diagnosed with open-angle glaucoma and began treatment with an antiglaucoma drug in the form of eye drops. She last used the eye drops 1 hour ago. Examination shows pupils of normal size that are reactive to light. An ECG shows sinus bradycardia. This patient is most likely undergoing treatment with which of the following drugs?

	Answer	Image
A	Latanoprost	
B	Timolol	
C	Dorzolamide	
D	Apraclonidine	
E	Tropicamide	
F	Pilocarpine	

Hint

This drug decreases adenylate cyclase activity. It is used to treat glaucoma because it decreases aqueous humor synthesis.

Correct Answer


A - Latanoprost

Explanation Why

[Latanoprost](#), a [prostaglandin F2 \$\alpha\$ agonist](#) analog, decreases [intraocular pressure](#) by increasing the outflow of [aqueous humor](#). Common side effects include discoloration of the [iris](#), eyelids, and [conjunctiva](#) as well as blurred vision, watering/burning, and foreign body sensation. [Bradycardia](#) is not an adverse effect of [latanoprost](#).

B - Timolol

Image



Explanation But

[Nonselective beta blockers](#) also cause bronchoconstriction, especially in patients with reactive

[airway](#) disease, and should be used with caution in patients with [asthma](#).

Explanation Why

[Timolol](#) is a nonselective [beta blocker](#) that is used as an ophthalmic formulation for the treatment of [open-angle glaucoma](#). It blocks β_2 receptors in the [ciliary body](#) of the [eye](#), decreasing [aqueous humor](#) production and consequently lowering [intraocular pressure](#). However, because [timolol](#) is passively absorbed into the [systemic circulation](#) from the [eye](#), it has high [bioavailability](#), potentially leading to systemic side effects such as the blockade of β_1 receptors in the [heart](#) and subsequent negative inotropic and [chronotropic](#) effects. In excess, these effects can cause [bradyarrhythmias](#) and [orthostatic hypotension](#), both of which can lead to [syncope](#), as seen in this patient.

C - Dorzolamide

Explanation Why

Dorzolamide, a [carbonic anhydrase inhibitor](#), reduces [aqueous humor](#) production. Common side effects include an ocular burning sensation, [pruritus](#), lacrimation, blurred vision, and a [bitter taste](#), but [bradycardia](#) is not a known side effect.

D - Apraclonidine

Explanation Why

[Apraclonidine](#) reduces [intraocular pressure](#) by decreasing [aqueous humor](#) production. While [apraclonidine](#) is associated with [bradycardia](#) in rare cases, other side effects, e.g., ocular hyperemia, [pruritus](#), foreign body sensation, and blurry vision, are more common.

E - Tropicamide

Explanation Why

[Tropicamide](#), an [anticholinergic](#) medication, is used commonly as a [mydriatic](#) agent. Pupillary dilation can precipitate [angle-closure glaucoma](#) in predisposed individuals and is contraindicated in

patients with [glaucoma](#). Moreover, [tachycardia](#), not [bradycardia](#), is a side effect of [tropicamide](#), making a different agent more likely.

F - Pilocarpine

Explanation Why

[Pilocarpine](#), a [cholinergic agonist](#) that primarily acts on [M3 muscarinic receptors](#), reduces [intraocular pressure](#) by increasing the [trabecular outflow](#) of [aqueous humor](#). [Pilocarpine](#) can cause [bradycardia](#), which could explain this patient's [syncope](#). However, [pilocarpine](#) also causes sweating, nausea, blurred vision and pupillary constriction, none of which are seen in this patient.

Question # 38

A 72-year-old man with type 2 diabetes mellitus, hypertension, and systolic heart failure comes to the physician because of a 5-day history of progressively worsening shortness of breath at rest. Physical examination shows jugular venous distention, diffuse crackles over the lower lung fields, and bilateral lower extremity edema. As a part of treatment, he is given a derivative of a hormone that acts by altering guanylate cyclase activity. This drug has been found to reduce pulmonary capillary wedge pressure and causes systemic hypotension as an adverse effect. The drug is most likely a derivative of which of the following hormones?

	Answer	Image
A	Angiotensin II	
B	Aldosterone	
C	Antidiuretic hormone	
D	Brain natriuretic peptide	
E	Prostacyclin	
F	Somatostatin	

Hint

This patient presents with acute decompensated congestive heart failure (CHF). The drug that was most likely prescribed to this patient is nesiritide.

Correct Answer

A - Angiotensin II

Explanation Why

[Angiotensin II](#) is a renal [hormone](#) that is released as a response to decreased intravascular filling, resulting in increased water and sodium reabsorption. The reduced [cardiac output](#) in [CHF](#) is among other pathomechanisms caused by a [systolic ventricular dysfunction](#) due to an increased [preload](#). An increased water and sodium reabsorption would further increase the [preload](#) and actually worsen this patient's condition. In fact, ACE inhibitors and [angiotensin II receptor blockers](#) are standard treatment options for [CHF](#).

B - Aldosterone

Explanation Why

[Aldosterone](#) is a renal [hormone](#) that regulates sodium and potassium secretion and reabsorption through the [collecting duct](#) system. [Aldosterone](#) analogs, like [spironolactone](#) and [eplerenone](#), are common treatment options for patients with [CHF](#) but are rarely recommended in [acute decompensated CHF](#). Potential side effects of [aldosterone](#) include [hypotension](#) and electrolyte disturbances instead of reduced [pulmonary capillary wedge pressure](#). Moreover, [aldosterone](#) analogs act through [nuclear receptors](#) and do not affect [guanylate cyclase](#) activity.


C - Antidiuretic hormone

Explanation Why

[Antidiuretic hormone](#) ([ADH](#)) mediates renal water reabsorption (via [V2 receptors](#)) and vasoconstriction (via [V1 receptors](#)). Vasopressin receptor antagonists like [tolvaptan](#) have been experimentally used to treat patients with acute [CHF](#). However, they operate via the [cAMP](#) pathway and do not affect [guanylate cyclase](#) activity.

D - Brain natriuretic peptide

Image



Explanation Why

[Brain natriuretic peptide \(BNP\)](#) is a [hormone](#) naturally released by ventricular [heart](#) muscle cells as a response to increased [ventricular filling](#). Derivatives of [BNP](#), like [nesiritide](#), are rarely used today but represent a treatment option in patients with [decompensated CHF](#). [BNP](#) causes [vasodilation](#) through the increase of [cGMP](#) and reduces [pulmonary capillary wedge pressure](#).

E - Prostacyclin

Explanation Why

[Prostacyclin](#) is a [prostaglandin](#) with agonistic activity at the [PGI₂](#) receptor that causes [vasodilation](#) and decreased [platelet aggregation](#). Common adverse effects include [hypotension](#). However, [prostacyclin](#) is used to treat [pulmonary hypertension](#), not acute [CHF](#). Moreover, it acts through [G](#)

[protein-coupled receptors](#) and does not affect [guanylate cyclase](#) activity.


F - Somatostatin

Explanation Why

[Somatostatin](#) is a [hormone](#) that inhibits several gastrointestinal [hormones](#) (e.g., [secretin](#), [histamine](#), [gastrin](#)) and neural [hormones](#) (e.g., [growth hormone](#)). [Octreotide](#) is a derivative of [somatostatin](#) commonly used in clinical practice. However, [octreotide](#) acts through agonistic effects at [somatostatin](#) receptors and not through intracellular [cGMP](#) increase. Moreover, it is used to treat [portal hypertension](#), [acromegaly](#), and [carcinoid tumors](#), not [acute decompensated CHF](#).

Question # 39

A 71-year-old woman comes to the physician because of dizziness and intermittent episodes of heart palpitations for 5 days. During this time, she has also had one episode of syncope. An ECG shows absence of P waves and irregular RR intervals. Treatment with an antiarrhythmic drug is initiated. The effect of the drug on the cardiac action potential is shown. Which of the following cardiac ion channels is most likely targeted by this drug?



	Answer	Image
A	Voltage-gated nonselective cation channels	
B	Voltage-gated sodium channels	
C	Voltage-gated potassium channels	

	Answer	Image
D	Voltage-gated calcium channels	Class IV The graph shows membrane potential on the y-axis and time on the x-axis. A horizontal dashed line represents the threshold potential, and a solid line at 0 mV is also shown. A blue curve represents the baseline action potential, rising from a resting potential, crossing the threshold, peaking, and then repolarizing. A green curve shows the effect of voltage-gated calcium channel activation: the upstroke (phase 0) is less steep (decreased slope), the peak is higher, and the repolarization phase (phase 4) is prolonged, taking more time to return to the resting potential. A pink curve shows the effect of decreased slope of phase 4 and phase 0 depolarization, which is a shallower repolarization phase. A callout box points to the peak of the green curve with the following text: • Decreased slope of phase 4 and phase 0 depolarization • Prolonged repolarization
E	Voltage-gated chloride channels	

Hint

Absent P waves and irregular RR intervals on ECG are diagnostic of atrial fibrillation, which commonly manifests with intermittent palpitations and dizziness. Class III antiarrhythmic drugs can be used for acute termination and rhythm control, as they prolong repolarization and the effective refractory period.

Correct Answer

A - Voltage-gated nonselective cation channels

Explanation Why

Voltage-gated nonselective cation channels play an important role in cardiac pacemaking, with the most well-known example being the [funny current](#). The [funny current](#) mediates the slow, spontaneous [depolarization](#) of [myocytes](#) during [diastole](#), which accounts for the intrinsic pacemaking of the [heart](#). This channel can be blocked pharmacologically by [ivabradine](#), causing prolonged [depolarization](#) (phase 4 of the [cardiac action potential](#)) and reducing the intrinsic firing rate of the [sinoatrial node](#). Blockage of these channels would not cause the slowed [repolarization](#) phase seen in this patient.

B - Voltage-gated sodium channels

Explanation Why

[Voltage-gated sodium channels](#) are the target of class I antiarrhythmic drugs, which can be further subdivided into [class IA antiarrhythmics](#), [class IB antiarrhythmics](#), and [class IC antiarrhythmics](#). Although there are subtle differences in how drugs from each class affect specific aspects of the [action potential](#), they all reduce or block conduction across cardiac [myocytes](#) by impairing phase 0 [depolarization](#) of the [action potential](#). On graphs, this can be seen as a change in slope of the rapid [depolarization](#) phase. None of the class I antiarrhythmic agents reduce the rate of [repolarization](#) or prolong [repolarization](#) like the agent given to this patient.

C - Voltage-gated potassium channels


Explanation Why

[Voltage-gated potassium channels](#) are the target of [class III antiarrhythmic drugs](#) like [sotalol](#), [amiodarone](#), [dofetilide](#), and [ibutilide](#). The markedly prolonged [repolarization](#) on the [action potential](#) curve here indicates potassium channel inhibition, which normally functions to restore resting cardiac [myocyte](#) membrane polarity. This effect prolongs the overall [action potential](#) and the

[effective refractory period](#), which explains the effectiveness of these drugs in treating [tachyarrhythmias](#) such as [atrial fibrillation](#), atrial flutter, and [ventricular tachycardias](#).

D - Voltage-gated calcium channels

Image



Explanation Why

Voltage-gated calcium channels are the target of [class IV antiarrhythmic drugs](#) like [verapamil](#) and [diltiazem](#). These drugs work by inhibiting slow cardiac calcium channels, which slows conduction velocity in the [sinoatrial node \(SA node\)](#) and [atrioventricular node \(AV node\)](#). This causes the slope during phase 0 and 4 of the [cardiac action potential](#) to decrease. The markedly prolonged [repolarization](#) seen in the [action potentials](#) of the cardiac [myocyte](#) here suggests inhibition of another voltage-gated ion channel.

E - Voltage-gated chloride channels

Explanation Why



Voltage-gated [chloride](#) channels are crucial for a host of biological processes. However, they do not play any major role in cardiac [action potential](#) generation. The markedly prolonged [repolarization](#) seen in the [action potential](#) of this patient suggests inhibition of another voltage-gated ion channel.

Question # 40

A 56-year-old man comes to the physician for increasing shortness of breath and retrosternal chest pain on exertion. He has smoked 2 packs of cigarettes daily for 35 years. His blood pressure is 145/90 mm Hg. Cardiac examination is shown. Which of the following is the most likely cause of this patient's auscultation findings?

[Video](#)

	Answer	Image
A	Constrictive pericarditis	
B	Mitral valve prolapse	The image displays two cardiac waveforms. The top waveform is an Electrocardiogram (ECG) showing a normal sinus rhythm with P waves, QRS complexes, and T waves. The bottom waveform is a Phonocardiogram (PCG) showing heart sounds. A mid-systolic click is visible, followed by a late systolic murmur, which is characteristic of mitral valve prolapse. The waveforms are labeled with 'Diastole' and 'Systole' phases, and 'S1' and 'S2' heart sounds.
C	Left ventricular failure	The diagram shows a frontal view of the human torso with the heart and lungs. Numbered circles indicate auscultation points: 1 (Aortic valve), 2 (Pulmonary valve), 3 (Tricuspid valve), 4 (Mitral valve), and 5 (Erb's point). The points are further labeled with 'a' and 'b' to indicate specific locations on the chest.

	Answer	Image
D	Aortic valve sclerosis	 The image displays an ECG (blue line) and a Phonocardiogram (orange line) synchronized with heart cycles. The background is divided into Diastole (grey) and Systole (white) phases. The ECG shows normal QRS complexes. The Phonocardiogram shows a normal aortic valve closure sound (A2) at the S2 position, with no significant diastolic murmur.
E	Ventricular septal rupture	
F	Right ventricular hypertrophy	
G	Mitral valve stenosis	 The image displays an ECG (blue line) and a Phonocardiogram (orange line) synchronized with heart cycles. The background is divided into Diastole (grey) and Systole (white) phases. The ECG shows normal QRS complexes. The Phonocardiogram shows a characteristic diastolic murmur (M2) at the S2 position, indicated by an arrow labeled 'OS' (Opening Snap), which is a key finding in mitral valve stenosis.

Hint

This patient's cardiac auscultation findings are consistent with an S₃ heart sound.

Correct Answer


A - Constrictive pericarditis

Explanation Why

[Constrictive pericarditis](#) may present with [dyspnea](#) and [chest pain](#) on exertion, which this patient also has. However, it is associated with a [pericardial friction rub](#) and/or a [pericardial knock](#), rather than an [S3 gallop](#). In addition, [constrictive pericarditis](#) would manifest with other symptoms such as [jugular vein distention](#) and peripheral [edema](#).

B - Mitral valve prolapse

Image



Explanation Why


[Mitral valve prolapse \(MVP\)](#) may present with [dyspnea](#) and [chest pain](#) on exertion, which this

patient also has. However, [MVP](#) is associated with a midsystolic click and a late [systolic](#) crescendo [murmur](#) heard best at the apex, rather than an [S3](#).

C - Left ventricular failure

Image

- ① Aortic valve
- ② Pulmonary valve
- ③ Tricuspid valve
- ④ Mitral valve
- ⑤ Erb's point



Explanation Why

[Left ventricular failure](#) is the most likely cause of this patient's [S3 heart sound](#), which results from a sudden deceleration of blood flow from the [left atrium](#) into the [left ventricle](#). While it can be a normal variant in young, healthy individuals and during [pregnancy](#), an [S3](#) is often associated with [congestive heart failure](#) and/or volume overload in older individuals such as this patient, especially in the setting of a dilated [left ventricle](#).

D - Aortic valve sclerosis

Image



Explanation Why

[Aortic valve sclerosis](#) can present with [dyspnea](#) and [chest pain](#) on exertion. However, auscultation would likely show a mid-[systolic ejection murmur](#) heard best at the right upper sternal border. An [S3](#) can occur in severe [aortic stenosis](#), but it would not be expected in isolated [aortic valve sclerosis](#).

E - Ventricular septal rupture

Explanation Why

While ventricular septal rupture may manifest with worsening [dyspnea](#) and [chest pain](#), it typically occurs in acutely ill patients following [myocardial infarction](#) and would present with signs of hemodynamic instability. Furthermore, the associated [murmur](#) would be a harsh, [holosystolic murmur](#) heard best at the left lower sternal border, rather than an isolated [S3](#).


F - Right ventricular hypertrophy

Explanation Why

Depending on the underlying etiology (e.g, [pulmonary hypertension](#)), [right ventricular hypertrophy \(RVH\)](#) may manifest with [dyspnea](#) and retrosternal [chest pain](#), which are also seen in this patient. However, [RVH](#) is associated with an [S4 gallop](#), which occurs during late [diastole](#) immediately prior to ventricular contraction, rather than in early [diastole](#).

G - Mitral valve stenosis

Image



Explanation Why

[Mitral valve stenosis](#) can present with [dyspnea](#) and [chest pain](#) on exertion, which this patient has. However, auscultation would reveal an [opening snap](#), followed by a mid-[diastolic murmur](#) that

terminates with the initiation of ventricular contraction rather than an [S3](#).

Question # 1

A 72-year-old woman comes to the emergency department because of a 2-week history of worsening shortness of breath, lower extremity swelling, and a 3-kg (6.6-lb) weight gain. Crackles are heard on auscultation of the chest. Cardiac examination shows a dull, low-pitched early diastolic sound at the 5th left intercostal space that becomes louder in the left lateral decubitus position at end-expiration. Which of the following is the most likely cause of these auscultation findings?

	Answer	Image
A	Decreased left myocardial compliance	
B	Increased ventricular contractility	
C	Increased left ventricular end-systolic volume	
D	Decreased left-ventricular filling pressure	
E	Increased capacity of the pulmonary circulation	

Hint

The patient presents with clinical features of heart failure (shortness of breath, crackles, lower extremity swelling, weight gain); the early diastolic heart sound is S₃, a strong predictor for heart failure.

Correct Answer

A - Decreased left myocardial compliance

Explanation Why

Decreased [myocardial](#) compliance is the cause of the [S4 gallop](#), which is an important sign of [heart failure with preserved ejection fraction](#) ([diastolic heart failure](#)) or other etiologies with reduced [myocardial](#) compliance (e.g., active [ischemia](#), [hypertrophic cardiomyopathy](#)). However, in contrast to the auscultation findings in this patient, [S4 gallop](#) is a late [diastolic](#) sound.


B - Increased ventricular contractility

Explanation Why

[Heart failure](#) due to [systolic dysfunction](#) is due to decreased, not increased, ventricular contractility. Increased ventricular contractility can be seen after [myocardial hypertrophy](#) and is a sign of a well-functioning [heart](#).

C - Increased left ventricular end-systolic volume

Image



Explanation Why

Increased [left ventricular](#) end-[systolic](#) volume is the most likely cause of this patient's [third heart sound](#) ([S₃](#)). In [systolic heart failure](#), the [myocardium](#) is overly compliant and cannot effectively contract to empty during [systole](#), resulting in ventricular volume overload. During the rapid [ventricular filling](#) phase, a sudden deceleration of blood occurs when the ventricle reaches its elastic limit, producing a [third heart sound](#).

D - Decreased left-ventricular filling pressure

Explanation Why

The [left ventricular](#) filling pressure would be decreased in a patient with [hypovolemia](#). In this patient with features of [heart failure](#), LV filling pressure would be increased.

E - Increased capacity of the pulmonary circulation

Explanation Why

An increase in the capacity of the [pulmonary circulation](#) occurs during inspiration as a result of a decrease in intrathoracic pressure. This results in a [split S₂](#) due to early closure of the [aortic valve](#) (A₂) and delayed closure of the [pulmonary valve](#) (P₂). However, it does not result in an S₃, nor is it related to [heart failure](#).

Question # 2

A 65-year-old African American man comes to the physician for a follow-up examination after presenting with elevated blood pressure readings during his last visit. He has no history of major medical illness and takes no medications. He is 175 cm (5 ft 9 in) tall and weighs 68 kg (150 lb); BMI is 22 kg/m². His pulse is 80/min and blood pressure is 155/90 mm Hg. Laboratory studies show no abnormalities. Which of the following is the most appropriate initial pharmacotherapy for this patient?

	Answer	Image
A	Valsartan	
B	Spirolactone	
C	Metoprolol	
D	Chlorthalidone	
E	Aliskiren	
F	Captopril	

Hint

This patient meets the diagnostic criteria for hypertension. Potential adverse effects of the first-line drug for this condition include hypokalemia, hyperglycemia, and hypercalcemia.

Correct Answer

A - Valsartan

Explanation Why

[Valsartan](#), along with other [angiotensin receptor blockers](#), is indicated for the treatment of [hypertension](#) in patients with [diabetes mellitus](#), renal disease, [ischemic heart disease](#), and/or [heart failure](#) who do not tolerate ACE inhibitors. This patient does not meet these criteria. As such, [valsartan](#) is not an appropriate treatment for a African American patient with isolated [hypertension](#).

B - Spironolactone

Explanation Why

[Spironolactone](#) is an [aldosterone](#) ([mineralocorticoid](#)) receptor antagonist that can be used to treat [hypertension](#) due to primary aldosteronism or as adjunct therapy in [resistant hypertension](#). It is not the first-line therapy for [hypertension](#) in African American patients with isolated [hypertension](#).

C - Metoprolol

Explanation Why

[Metoprolol](#) is a second-line therapy for [hypertension](#) that is typically used in patients with cardiovascular comorbidities such as [ischemic heart disease](#), [heart failure](#), or [atrial fibrillation](#). [Metoprolol](#) is not indicated as a first-line therapy for isolated [hypertension](#) in any patient group.

D - Chlorthalidone

Explanation Why

[Chlorthalidone](#) is a preferred first-line drug for African American patients with isolated [hypertension](#). [Thiazide diuretics](#) are considered especially effective antihypertensive agents because impaired salt excretion and subsequent intravascular volume expansion are thought to be the primary drivers of [hypertension](#) in this population. [Hypokalemia](#), [hyperglycemia](#), and [hypercalcemia](#) are all side effects seen with chronic [chlorthalidone](#) use; [hyperglycemia](#), especially, limits the use of [chlorthalidone](#) in patients with [metabolic syndrome](#). [Dihydropyridine calcium channel blockers](#) can also be used as first-line therapy in African American patients, particularly those with [metabolic syndrome](#).

E - Aliskiren


Explanation Why

[Aliskiren](#) is a [direct renin inhibitor](#) that decreases [RAAS](#) activity, which in turn decreases the amount of [angiotensin II](#) and [aldosterone](#) in circulation, lowering blood pressure. [Aliskiren](#) is used to treat [arterial hypertension](#) in situations in which neither ACE inhibitors nor angiotensin-receptor blockers are tolerated. [Aliskiren](#) is not the first-line therapy for [hypertension](#) in African American patients with isolated [hypertension](#).

F - Captopril


Explanation Why

[Captopril](#) is a first-line antihypertensive drug in white patients with [diabetes mellitus](#), renal disease, [ischemic heart disease](#), or [heart failure](#). [Captopril](#), along with other ACE inhibitors, is beneficial in these populations because it reduces circulating levels of [angiotensin II](#) and [aldosterone](#). The pathophysiology of [hypertension](#) in most African American patients is not significantly influenced by the [renin-angiotensin-aldosterone system](#), which limits the effectiveness of ACE inhibitors. For this reason, [captopril](#) is not a first-line therapy in African American patients with isolated [hypertension](#).



Question # 3

A 50-year-old man undergoes parathyroidectomy for treatment-resistant hyperparathyroidism. The procedure is complicated by brisk bleeding from an artery that travels along the external branch of the superior laryngeal nerve. To stop the bleeding, the artery is ligated at its origin. Which of the following is most likely the origin of the artery that was injured in this patient?

	Answer	Image
A	Subclavian artery	
B	Internal carotid artery	
C	External carotid artery	 The diagram illustrates the arterial and venous systems of the neck. On the left side, the external carotid artery is shown originating from the common carotid artery and branching into the superior thyroid artery, inferior thyroid artery, thyrocervical trunk, and subclavian artery. On the right side, the internal carotid artery is shown ascending. The venous system includes the superior, middle, and inferior thyroid veins, the internal jugular vein, and the brachiocephalic veins. The superior vena cava is also visible at the base of the neck.
D	Thyrocervical trunk	
E	Ascending pharyngeal artery	

Hint

The superior thyroid artery travels along the superior laryngeal nerve.

Correct Answer

A - Subclavian artery

Explanation Why

The [subclavian artery](#) gives rise to the [inferior thyroid artery](#) via the [thyrocervical trunk](#). The [inferior thyroid artery](#) pierces the lower pole of the [thyroid gland](#) and is closely associated with the [recurrent laryngeal nerve](#). This patient has an injury of a vessel traveling with the [superior laryngeal nerve](#) rather than with the RLN.


B - Internal carotid artery

Explanation Why

The cervical portion of the [internal carotid artery](#) (ICA) has no branches and neither of the [thyroid arteries](#) arise from the ICA. As a result, ligation of the ICA would not affect this patient's bleeding. The first branches of the [internal carotid](#) arise in the petrous segment of the ICA. Before it terminates in the brain, the major branches include the [ophthalmic artery](#) and the superior hypophyseal [artery](#), along with several smaller branches.

C - External carotid artery

Image



Explanation Why

The superior [thyroid artery](#) is a direct branch of the [external carotid artery](#). After the carotid [artery](#) bifurcates in the [carotid triangle](#), the superior [thyroid artery](#) arises as the first branch of the [external carotid artery](#) and supplies the [cranial](#) half of the [thyroid gland](#) before forming an anastomotic network with the [inferior thyroid artery](#). The [artery](#) travels with the external branch of the [superior laryngeal nerve](#) and pierces the [thyroid gland](#) at the superior pole. Therefore, bleeding from the superior [thyroid artery](#) would be controlled by ligating it at its origin from the [external carotid artery](#).

D - Thyrocervical trunk

Explanation Why

The [thyrocervical trunk](#) originates from the [subclavian artery](#) and gives rise to the [inferior thyroid](#)

[artery](#). The [inferior thyroid artery](#) pierces the lower pole of the [thyroid gland](#) and is closely associated with the [recurrent laryngeal nerve](#). This patient has an injury of a vessel traveling with the [superior laryngeal nerve](#) rather than with the RLN.

E - Ascending pharyngeal artery

Explanation Why

The [ascending pharyngeal artery](#) arises from the [external carotid artery](#) and gives rise to a pharyngeal and neuromeningeal trunk. Branches of the [ascending pharyngeal artery](#) supply the pharyngeal constrictor muscles, [tympanic cavity](#), and [meninges](#) but they do not travel along the external branch of the [superior laryngeal nerve](#). Bleeding from the [ascending pharyngeal artery](#) or one of its branches would be unlikely during a [parathyroidectomy](#).

Question # 4

Three days after undergoing laparoscopic colectomy, a 67-year-old man reports swelling and pain in his right leg. He was diagnosed with colon cancer 1 month ago. His temperature is 38.5°C (101.3°F). Physical examination shows swelling of the right leg from the ankle to the thigh. There is no erythema or rash. Which of the following is likely to be most helpful in establishing the diagnosis?

	Answer	Image
A	D-dimer level	
B	Transthoracic echocardiography	
C	CT pulmonary angiography	
D	Blood cultures	
E	Compression ultrasonography	<pre> graph TD Start[Suspected DVT] --> Wells[Calculate Wells score for DVT] Wells --> Low[Low pretest probability] Wells --> Int[Intermediate pretest probability] Wells --> High[High pretest probability] Low --> PE[Provoked risk factor, surgery, or active malignancy?] PE --> DVT_L[Whole-leg compression US with venous duplex] PE --> DVT_L DVT_L --> DVT_L_Pos[DVT diagnosed] DVT_L --> DVT_L_Neg[DVT ruled out] DVT_L_Pos --> Consider_Venography_L[Consider venography, CT venography, or MR venography] DVT_L_Neg --> Consider_Venography_L Int --> DVT_I[Whole-leg compression US with venous duplex] DVT_I --> DVT_I_Pos[DVT diagnosed] DVT_I --> DVT_I_Neg[DVT ruled out] DVT_I_Pos --> Consider_Venography_I[Consider venography, CT venography, or MR venography] DVT_I_Neg --> Consider_Venography_I High --> DVT_H[Whole-leg compression US with venous duplex] DVT_H --> DVT_H_Pos[DVT diagnosed] DVT_H --> DVT_H_Neg[DVT ruled out] DVT_H_Pos --> Consider_Venography_H[Consider venography, CT venography, or MR venography] DVT_H_Neg --> Repeat[Repeat US within a week if no alternate diagnosis] Repeat --> DVT_H </pre>

Hint

This patient has swelling and pain in his right lower extremity and fever with no erythema. Using the Wells score for deep vein thrombosis (DVT), his pretest probability of a DVT is at least 3 points (i.e., major surgery within 4 weeks, malignancy, leg swelling), which indicates a high probability of DVT.

Correct Answer

A - D-dimer level

Explanation Why

[D-dimer](#) level testing is frequently used to rule out [DVT](#). However, it should only be used when [pretest probability](#) is low (i.e., [Wells score](#) ≤ 1) rather than high, which is seen here. In addition, [D-dimer](#) testing is not recommended in postoperative patients because the concentration of [D-dimer](#) is rarely normal in this state (leading to frequent [false positives](#))

B - Transthoracic echocardiography

Explanation Why

[Transthoracic echocardiography](#) would help in the diagnosis of [right heart failure](#) due to a [massive pulmonary embolism \(PE\)](#), which could also result in a high [Wells score](#). However, a [massive PE](#) typically manifests with severe [dyspnea](#), [chest pain](#), [hypotension](#), and [shock](#), none of which are seen in this patient. A [TTE](#) would not be helpful in the diagnosis of this patient's [DVT](#).

C - CT pulmonary angiography

Explanation Why

CT [pulmonary angiography](#) is the test of choice for the diagnosis of [pulmonary embolism \(PE\)](#), for which the pretest possibility can also be calculated using the [Wells score](#). However, [PE](#) typically presents with [chest pain](#), [shortness of breath](#), and/or [tachycardia](#), none of which this patient has. A [CT pulmonary angiography](#) would not be helpful in diagnosing this patient's likely [DVT](#).


D - Blood cultures

Explanation Why

Blood cultures are useful in diagnosing [bacteremia](#), which can be caused by [cellulitis](#). [Cellulitis](#) typically manifests with [fever](#) as well as local swelling and [pain](#). However, it also manifests with [erythema](#), which is not seen in this patient. Moreover, [cellulitis](#) does not have a high [Wells score](#).

E - Compression ultrasonography

Image



Explanation Why

[Compression ultrasonography](#) is the test of choice for diagnosing [DVT](#). It is affordable, reliable, and safe, making it an ideal first-line test. [Sonographic](#) findings that suggest a [thrombus](#) include a noncompressible [vein](#), visible hyperechoic mass, and/or absent or abnormal venous flow.

Question # 5

A 55-year-old man with hypertension, hyperlipidemia, type 2 diabetes mellitus, and asthma comes to the physician because of a 2-month history of intermittent dry, hacking cough. He does not have fever, chest pain, or shortness of breath. He does not smoke cigarettes. Current medications include simvastatin, metformin, albuterol, and ramipril. His temperature is 37°C (98.6°F), pulse is 87/min, and blood pressure is 142/88 mm Hg. Cardiopulmonary examination shows no abnormalities. Which of the following is the most appropriate initial action by the physician?

	Answer	Image
A	Start dextromethorphan and increase frequency of albuterol	
B	Stop simvastatin and start atorvastatin	
C	Stop ramipril and start candesartan	The diagram illustrates the RAAS pathway. Renin converts Angiotensinogen to Angiotensin I. ACE converts Angiotensin I to Angiotensin II. Angiotensin II causes arteriolar vasoconstriction and efferent arteriole constriction, maintaining GFR. It also stimulates Aldosterone release from the adrenal cortex and ADH release from the posterior pituitary. Aldosterone and ADH increase water and sodium reabsorption in the distal tubule and collecting duct, leading to increased blood pressure and cardiac output. ACE inhibitors like ramipril block the conversion of Angiotensin I to Angiotensin II, which can cause a dry cough. ARBs like candesartan block the effects of Angiotensin II without affecting the cough.
D	Stop ramipril and start lisinopril	
E	Stop albuterol and start salmeterol	

Hint

Increased local concentration of bradykinin and substance P is thought to be responsible for the symptom this patient is experiencing.

Correct Answer

A - Start dextromethorphan and increase frequency of albuterol

Explanation Why

[Dextromethorphan](#) is an [NMDA glutamate](#) receptor antagonist used as an over-the-counter [cough suppressant](#) and [albuterol](#) is a [beta-2 agonist](#) used in the treatment of [asthma](#). While [dextromethorphan](#) might temporarily mitigate this patient's [cough](#), it would not address the root cause of it. Additionally, there is no other evidence that this patient's [cough](#) is associated with an asthma exacerbation (such as [dyspnea](#), wheezing, or chest tightness); increasing [albuterol](#) is therefore unlikely to resolve this problem.


B - Stop simvastatin and start atorvastatin

Explanation Why

Both [simvastatin](#) and [atorvastatin](#) are [statins](#), which are indicated to treat [hyperlipidemia](#) and act by decreasing [LDL](#) levels. Classic side effects of [statins](#) include [myopathy](#) and [rhabdomyolysis](#) but not [cough](#). Therefore, changing the patient's medication from [simvastatin](#) to [atorvastatin](#) would be unlikely to resolve his [cough](#).

C - Stop ramipril and start candesartan

Image



Explanation But

Other common adverse effects of ACE inhibitors include [hypotension](#), [hyperkalemia](#), a decrease in [glomerular filtration rate](#), and [angioedema](#).

Explanation Why

[Ramipril](#) is an ACE inhibitor, a class of drugs that is known to cause a characteristic dry [cough](#). ACE inhibitors decrease blood pressure by inhibiting the [renin-angiotensin-aldosterone](#) system (RAAS). Inhibition of [ACE](#) also causes an increase in [bradykinin](#), which can lead to bronchoconstriction and dry [cough](#). Patients presenting with [cough](#) caused by ACE inhibitors should be changed to an [angiotensin receptor blocker](#) (ARB) such as [candesartan](#). [Cough](#) is a rare adverse effect of [ARBs](#), as these agents do not influence the breakdown of kinins.

D - Stop ramipril and start lisinopril

Explanation Why

[Ramipril](#) and [lisinopril](#) are both ACE inhibitors. A common side effect of treatment with ACE inhibitors is an intermittent, non-productive [cough](#), such as the one seen in this patient. Because all ACE inhibitors are known to cause this [cough](#), switching from one agent to another within this class would be unlikely to resolve the problem.



E - Stop albuterol and start salmeterol




Explanation Why

[Albuterol](#) and [salmeterol](#) are both [beta-2 adrenergic agonists](#) used in the treatment of [asthma](#). While [albuterol](#) is a short-acting agent used for acute [asthma](#) exacerbations, [salmeterol](#) is a long-acting agent used for prophylaxis of [asthma](#) attacks. Although this patient has a history of [asthma](#), his current [cough](#) seems unrelated to it. He does not have a history or findings to suggest exacerbation of [asthma](#) (e.g., exposure to triggers, [dyspnea](#), or [tachypnea](#) on examination). Moreover, stopping [albuterol](#) would not be appropriate if there was any suspicion of an asthma exacerbation.

Question # 6

A 2-year-old boy is brought to the physician for evaluation of delayed onset of speech. Over the past year, he has also had recurrent dizziness and three episodes of syncope. Examination of the ears shows clear auditory canals and intact tympanic membranes with normal light reflexes. Visual reinforcement audiometry shows bilateral sensorineural deafness. Genetic analysis reveals a mutation in the *KCNQ1* gene causing a defect in slow voltage-gated potassium channels. An electrocardiogram of this patient is most likely to show which of the following?

	Answer	Image
A	Pseudo-right bundle branch block	 ECG showing pseudo-right bundle branch block. The QRS complex in leads I, II, III, aVR, aVL, and aVF shows a deep S wave, which is characteristic of this condition. The QRS complex in leads V1, V2, V3, V4, V5, and V6 shows a deep S wave, which is also characteristic of this condition. The QRS complex in leads V1, V2, V3, V4, V5, and V6 shows a deep S wave, which is also characteristic of this condition. 25mm/s 10mm/mV 40Hz 0.05s 12SL 233 CTD 1.5 Courtesy of P.G. Postema, M.D., AMC ECG-JPEDIA.ORG
B	Slurred upstroke of the QRS complex	 ECG showing slurred upstroke of the QRS complex. The QRS complex in leads I, II, III, aVR, aVL, and aVF shows a slurred upstroke, which is characteristic of this condition. The QRS complex in leads V1, V2, V3, V4, V5, and V6 shows a slurred upstroke, which is also characteristic of this condition. HR: 68 QRS: 126 ms QT: 400 ms QTc: 418 ms ST: 120 ms Td: 174 ms P ax: 84 QRS ax: 67 T ax: 71 QRS-T ax(3D): 116 Cardiac V4 37.7 05-02-2021 13:00 1 T: 0.00 - 0.35 25 mm/s 10 mm/mV
C	Epsilon wave following the QRS complex	

	Answer	Image
D	Progressive lengthening of the PR interval	 The image shows a single-lead ECG tracing on a standard grid. It displays four cardiac complexes. The PR interval, which is the time between the start of the P wave and the start of the QRS complex, progressively increases from the first complex to the fourth complex. This is a characteristic finding in second-degree Mobitz II heart block.
E	Absence of P waves	 The image shows a 12-lead ECG tracing. The leads are arranged in four rows: (I, aVR, V1, V4), (II, aVL, V2, V5), (III, aVF, V3, V6), and a fourth row with lead II. In all leads, there is no visible P wave preceding the QRS complex. The QRS complexes appear normal in morphology and timing, which is characteristic of Atrial Fibrillation.
F	Prolongation of the QT interval	 The image shows a 12-lead ECG tracing. The leads are arranged in three rows: (I, aVR, V1, V4), (II, aVL, V2, V5), and (III, aVF, V3, V6). The QT interval, which is the time from the start of the QRS complex to the end of the T wave, is visibly prolonged in all leads. The T waves are also somewhat flattened. This is a characteristic finding in Long QT Syndrome.


Hint

A mutation in the *KCNQ1* gene is the most common cause of Jervell and Lange-Nielsen syndrome, a condition associated with arrhythmias and sensorineural deafness.

Correct Answer

A - Pseudo-right bundle branch block

Image




Explanation Why

Pseudo-right [bundle branch](#) block and [ST elevations](#) in leads V₁ and V₂ are characteristic [ECG](#) findings of [Brugada syndrome](#), which can manifest with [dizziness](#) and [syncope](#). However, [Brugada syndrome](#) is caused by defective cardiac voltage-gated sodium channels (not potassium channels) and is most commonly seen in adults males of southeast Asian descent. Moreover, [Brugada syndrome](#) would not explain this patient's [sensorineural deafness](#).

B - Slurred upstroke of the QRS complex

Image



Explanation Why

A slurred upstroke of the [QRS complex](#) (i.e., delta wave) is a characteristic [ECG](#) finding of [Wolff-Parkinson-White \(WPW\)](#) syndrome, which may manifest with [dizziness](#) and [syncope](#). However, [WPW](#) is caused by an [accessory pathway](#) between the [atria](#) and ventricles and is not due to defective voltage-gated potassium channels. While familial forms of [WPW](#) exist, they are uncommon and are associated with mutations in the [PRKAG2 gene](#). Moreover, [WPW](#) would not explain this patient's [sensorineural deafness](#).

C - Epsilon wave following the QRS complex


Explanation Why

An epsilon wave following the [QRS complex](#) is a highly specific [ECG](#) finding for [arrhythmogenic](#)

[right ventricular cardiomyopathy \(ARVC\)](#), which may cause [dizziness](#) and [syncope](#). However, while this condition can exhibit either [autosomal dominant](#) or [autosomal recessive inheritance](#) and is associated with various genetic mutations, it is not caused by a defect in voltage-gated potassium channels. Moreover, [ARVC](#) would not explain this patient's [sensorineural hearing loss](#).

D - Progressive lengthening of the PR interval

Image




Explanation Why

Progressive lengthening of the [PR interval](#) until a beat is dropped is the characteristic [ECG](#) finding of [Mobitz type I](#), which may manifest with [dizziness](#) and [syncope](#). However, while [heart block](#) has many etiologies (e.g., increased [vagal](#) tone in athletes, [ischemic heart disease](#), medications), [Mobitz I](#) is not caused by defective voltage-gated potassium channels. Moreover, [Mobitz I](#) would not explain this patient's [sensorineural deafness](#).

E - Absence of P waves

Image




Explanation Why

Absence of [P waves](#) and an irregularly irregular rhythm are the characteristic [ECG](#) findings of [atrial fibrillation](#), which may cause [dizziness](#) and [syncope](#). However, while there are many [risk factors for atrial fibrillation](#) (e.g., [valvular heart disease](#), smoking, [hyperthyroidism](#)), it occurs almost exclusively in adults and is not caused by a defect in voltage-gated potassium channels. Moreover, [atrial fibrillation](#) would not explain this patient's [sensorineural deafness](#).

F - Prolongation of the QT interval

Image



Explanation But

Another [congenital LQTS](#) with a pure cardiac [phenotype](#) is [Romano-Ward syndrome](#), an [autosomal dominant](#) condition.

Explanation Why

Prolongation of the [QT interval](#) (typically with a [QTc](#) > 500 ms) is seen on [ECG](#) in patients with [Jervell and Lange-Nielsen syndrome](#), an [autosomal recessive](#) form of congenital [long QT syndrome](#) that is caused by a [loss-of-function](#) mutation in the potassium channel [genes](#) [KCNQ1](#) (90% of cases) or [KCNE1](#). In [cardiomyocytes](#), the defective potassium channels cause prolongation of the [ventricular action potential](#), which can lead to [dizziness](#), [syncope](#), [torsades de pointes](#), and [sudden cardiac death](#). In the [inner ear](#), the defect disrupts the production of endolymph, resulting in congenital [sensorineural deafness](#).

Question # 7

A 61-year-old woman is brought to the emergency department because of crushing substernal chest pain at rest for the past 2 hours. She is diaphoretic. Her temperature is 37.5°C (99.5°F), pulse is 110/min, respirations are 21/min, and blood pressure is 115/65 mm Hg. An ECG shows ST elevation in I, aVL, and V2–V4. Coronary angiography shows an 80% stenosis in the left anterior descending artery. Which of the following is the most likely initial step in the pathogenesis of this patient's coronary condition?

	Answer	Image
A	Intimal smooth muscle cell migration	
B	Intimal monocyte infiltration	
C	Platelet activation	
D	Low density lipoprotein oxidation	

	Answer	Image
E	Endothelial cell dysfunction	The diagram illustrates the progression of atherosclerosis through a series of cross-sections of an artery wall, categorized into three time scales: Asymptomatic, Decades, and Symptomatic. A red arrow at the top indicates the progression from left to right. Asymptomatic: Normal endothelium: Shows a healthy artery wall with a single layer of endothelial cells. Endothelial dysfunction: Shows the endothelium becoming thicker and less permeable. A box labeled "Chronic stress (e.g. due to hypertension, inflammation)" points to this stage. Infiltration of monocytes and LDL resulting in accumulation of macrophages and oxLDL: Shows monocytes and LDL entering the artery wall, where they become macrophages and oxidized LDL (oxLDL). Decades: Foam cell formation resulting in fatty streaks: Macrophages ingest oxLDL, becoming foam cells. Cholesterol crystals begin to form. Proliferation and migration of SMCs and deposition of ECM: Smooth muscle cells (SMCs) migrate from the media to the intima, proliferate, and deposit extracellular matrix (ECM). Symptomatic: Fibrous plaque formation: A fibrous cap forms over the plaque, containing SMCs and ECM. Plaque destabilization: The fibrous cap becomes thinner and more fragile. A box labeled "PDGF" points to this stage. Plaque rupture (e.g., in context of increased matrix degradation): The fibrous cap ruptures, exposing the underlying necrotic core. Thrombus formation: Platelets and fibrin form a thrombus on the ruptured plaque surface. Legend: - LDL (green circle) - Monocyte (purple circle) - Macrophage (yellow circle) - oxLDL (green star) - Foam cell (orange cell) - Platelets (red dots) - Lymphocytes (blue dots) - Extracellular matrix (ECM) (yellow mesh) - Smooth muscle cell (SMC) (pink cell) - Cholesterol crystals (blue crystals) - Cellular debris (brown dots) - PDGF (pink box) - Fibrous cap (pink layer) - Necrotic core (yellow core) - SMC apoptosis, breakdown of the stabilizing ECM (pink box)
F	Intimal foam cell accumulation	
G	Fibrous plaque formation	

Hint

This patient's symptoms (acute, crushing substernal chest pain, diaphoresis) and ECG changes suggest a diagnosis of acute coronary syndrome (ACS). The angiography showing a stenosis of the left anterior descending artery confirms the diagnosis. ACS is caused primarily by atherosclerosis.

Correct Answer

A - Intimal smooth muscle cell migration

Explanation Why

[Intimal smooth muscle cells](#) migration is mediated by [macrophages](#) producing [platelet-derived growth factor \(PDGF\)](#) and [fibroblast](#) growth factor (FGF). However, [intimal smooth muscle cell](#) migration is not the initial step of [atherosclerosis](#).

B - Intimal monocyte infiltration

Explanation Why

[Intimal monocyte](#) infiltration is one of the first steps in the pathogenesis of [atherosclerosis](#). After adhesion to the [endothelium](#), the [monocytes](#) emigrate through the [endothelium](#) into the [intima](#). As the [endothelium](#) is physiologically impermeable to [monocytes](#), [monocyte](#) infiltration is not the initial step of [atherosclerosis](#).

C - Platelet activation

Explanation Why

[Platelet activation](#) releases a variety of inflammatory mediators and thereby recruits inflammatory cells towards the site of the atherosclerotic lesion. However, [platelets](#) are only activated after adhesion to the [endothelium](#) of affected vessels. Therefore, [platelet activation](#) is not the initial step of [atherosclerosis](#).


D - Low density lipoprotein oxidation

Explanation Why

[Low-density lipoprotein \(LDL\)](#) oxidation releases [cholesterol](#), which is then ingested by [macrophages](#), inducing [foam cell](#) formation. However, attraction and activation of [macrophages](#) happen before the formation of [foam cells](#). Therefore, [LDL](#) oxidation is not the initial step of [atherosclerosis](#).

E - Endothelial cell dysfunction

Image



Explanation Why

The development of [atherosclerosis](#) begins with an external factor (e.g., [hypertension](#), smoking, advanced glycation end-products) damaging the [endothelium](#), which increases adhesiveness and leads to local [inflammation](#) and invasion of [smooth muscle cells](#) into the [tunica intima](#). Inflammatory

[cytokines](#) attract [macrophages](#), which then form [foam cells](#) by taking up [cholesterol](#) from oxidized low-density lipids. These [foam cells](#) accumulate to form [fatty streaks](#) (early atherosclerotic lesions). Finally, [collagen](#) production by lipid-laden [macrophages](#) and SMCs form these [fatty streaks](#) into a [fibrous plaque](#).

F - Intimal foam cell accumulation

Explanation Why

[Intimal foam cell](#) accumulation is a consequence of [macrophages](#) infiltrating the [intima](#) and ingesting [cholesterol](#) from oxidized [LDL](#). These [foam cells](#) accumulate and form [fatty streaks](#). However, [intimal foam cell](#) accumulation is not the initial step of [atherosclerosis](#).


G - Fibrous plaque formation

Explanation Why

[Fibrous plaque](#) formation is mediated by [macrophages](#), [lymphocytes](#), and [smooth muscle cells](#). Together with deposited [extracellular matrix](#) (e.g., [collagen](#)), these cell types form the [fibrous plaque](#) in [atherosclerosis](#). However, [fibrous plaque](#) formation is not the initial step of [atherosclerosis](#).

Question # 8

A 67-year-old man with type 2 diabetes mellitus comes to the emergency department because of lightheadedness over the past 2 hours. He reports that he has had similar episodes of lightheadedness and palpitations over the past 3 days. His only medication is metformin. His pulse is 110/min and irregularly irregular. An ECG shows a variable R-R interval and absence of P waves. The patient undergoes transesophageal echocardiography. During the procedure, the tip of the ultrasound probe is angled posteriorly within the esophagus. This view is most helpful for evaluating which of the following conditions?

	Answer	Image
A	Thrombus in the left pulmonary artery	
B	Myxoma in the left atrium	
C	Aneurysm of the descending aorta	
D	Thrombus in the left ventricular apex	
E	Tumor in the right main bronchus	

Hint

Correct Answer

A - Thrombus in the left pulmonary artery

Explanation Why

The [esophagus](#) lies [posterior](#) to the [right pulmonary artery](#), which can be seen in the [anterior](#) view on a [TEE](#). However, the [left pulmonary artery](#) is hardly visible in any view on [TEE](#).


B - Myxoma in the left atrium

Explanation Why

A [tumor](#) in the [left atrium](#), e.g., left atrial myxoma, can be visualized well on [TEE](#), but only when the probe is rotated anteriorly in the mid-[esophagus](#). The [left atrium](#), [right atrium](#), and [interatrial septum](#) can be seen in the [anterior](#) view.

C - Aneurysm of the descending aorta

Image



Explanation Why

The upper [esophagus](#) lies [posterior](#) to the [trachea](#) and [anterior](#) to the [vertebral column](#). Below the bifurcation of the [trachea](#), the [descending aorta](#) lies [posterior](#) to the lower [esophagus](#). An [aneurysm](#) of the [descending aorta](#) can, therefore, be visualized well on [transesophageal echocardiography \(TEE\)](#) through an [ultrasound](#) probe that is angled posteriorly within the [esophagus](#). Other structures lying [posterior](#) and in close proximity to the [esophagus](#) include the [thoracic duct](#) and [azygos vein](#).

D - Thrombus in the left ventricular apex

Explanation Why

A [thrombus](#) in the [left ventricular](#) apex can be visualized well on [transthoracic echocardiography \(TTE\)](#) because of the proximity of the ventricular apex to the [anterior chest wall](#). While a [TEE](#)

allows for better imaging quality when evaluating the cardiac structures that lie in close proximity to the [esophagus](#), more [anterior](#) structures such as the [left ventricular](#) apex would not be seen well.

E - Tumor in the right main bronchus

Explanation Why

The [esophagus](#) lies [posterior](#) to the [trachea](#) and left mainstem [bronchus](#). These structures could be visualized in the [anterior](#) view on a [TEE](#), but the right main [bronchus](#) would not be seen. A chest [x-ray](#) or CT imaging of the chest is indicated if a [bronchial tumor](#) is suspected.

Question # 9

A 62-year-old man comes to the physician because of a 5-day history of swelling in his left arm. Two months ago, he was diagnosed with a deep venous thrombosis in the left calf. He has had a 7-kg (15-lb) weight loss in the last 3 months. He has smoked 1 pack of cigarettes daily for the past 25 years. His only medication is warfarin. Physical examination shows warm edema of the left forearm with overlying erythema and a tender, palpable cord-like structure along the medial arm. His lungs are clear to auscultation bilaterally. Duplex sonography shows thrombosis of the left basilic and external jugular veins. Which of the following is the most appropriate next step to confirm the underlying diagnosis?

	Answer	Image
A	24-hour electrocardiography monitoring	
B	X-ray of the chest	
C	Transesophageal echocardiography	
D	CT scan of the abdomen	
E	Serum antiphospholipid antibody level	
F	Serum D-dimer level	

Hint

This patient's presents with a history of thrombosis and current superficial thrombophlebitis involving different anatomical locations, which is consistent with migratory thrombophlebitis. The presence of this sign should always raise suspicion for an underlying hypercoagulable state.

Correct Answer

A - 24-hour electrocardiography monitoring

Explanation Why

24-hour [electrocardiography](#) monitoring is useful for detecting paroxysmal [atrial fibrillation](#). Affected patients may present with seemingly inexplicable [arterial thromboembolism](#) (manifesting as [transient ischemic attack/stroke](#), [bowel infarction](#), or [acute limb ischemia](#)). The diagnosis is often delayed because [electrocardiogram](#) at the time of presentation may demonstrate normal [sinus rhythm](#), thus obscuring the need for further workup of a cardiogenic cause of the clotting. This patient has venous thromboses, however, which would not be explained by [paroxysmal atrial fibrillation](#) so 24-hour [electrocardiography](#) monitoring is not appropriate.

B - X-ray of the chest

Explanation Why

A chest [x-ray](#) can be used to identify potential [lung](#) malignancies, which could result in a [hypercoagulable state](#). This patient's pulmonary exam is unremarkable and he lacks concerning respiratory signs/symptoms, making a [lung malignancy](#) less likely. Finally, chest [x-ray](#) is no longer a recommended screening tool for patients at high risk for [lung cancer](#). Instead, a low-dose helical [CT scan](#) is the imaging modality of choice. Chest [x-ray](#) is less likely to help confirm the underlying diagnosis.

C - Transesophageal echocardiography

Explanation Why

[Transesophageal echocardiography](#) is useful for diagnosing a cardiogenic etiology of arterial thrombosis (e.g., atrial [thrombus](#) or valvular vegetations). Atrial [thrombi](#) most often arise in the setting of [atrial fibrillation](#) and manifest with recurrent arterial thromboses, not venous thromboses as seen here. Bacterial valvular vegetations can also cause recurrent arterial thrombosis due to [septic microemboli](#), manifesting with [Osler nodes](#), [Janeway lesions](#), and [nail](#) bed hemorrhages, all of which

are absent in this man. This patient presents with [DVT](#) and [superficial thrombophlebitis](#), making a cardiogenic etiology unlikely.

D - CT scan of the abdomen

Explanation Why

A [CT scan](#) of the abdomen is most likely to confirm the diagnosis of [Trousseau syndrome](#), a condition that is characterized by recurrent venous thrombosis in different [veins](#) with evidence of simultaneous [thrombophlebitis](#) (warm, red, swollen [skin](#) with tender cord-like structures) as seen here. This patient's older age, smoking history, and weight loss raise concern for a [malignancy](#)-induced [hypercoagulable state](#). [Trousseau syndrome](#) is most often seen in the setting of a visceral [malignancy](#) (classically, pancreatic adenocarcinoma) due to the production of [procoagulants](#) by the malignant cells.

E - Serum antiphospholipid antibody level

Explanation Why

Serum [antiphospholipid antibody](#) levels are elevated in patients with [antiphospholipid syndrome](#). Patients with APS are at significantly increased risk of developing recurrent venous thromboses, as seen in this man. However, other common manifestations of APS include recurrent arterial thrombosis (possibly causing [transient ischemic attack](#) or [stroke](#)) and [livedo reticularis](#), neither of which are reported or observed here. Finally, significant weight loss and the presence of simultaneous [superficial thrombophlebitis](#) are not typical of APS, making this diagnosis unlikely.

F - Serum D-dimer level

Explanation Why

Serum [D-dimer](#) level is most useful in determining whether or not a patient is experiencing a thrombotic event. Because the [D-dimer](#) value reflects the extent to which [fibrinolysis](#) is occurring, it can serve as indirect evidence of a venous thrombosis. In patients with a low [pretest probability](#) for [DVT](#), the [D-dimer](#) is considered especially useful as a screening tool. In this patient with a confirmed

history of [DVT](#), the [D-dimer](#) would add no useful information in confirming the underlying diagnosis.

Question # 10

A 38-year-old woman comes to the physician because of a 1-month history of palpitations. She does not smoke or drink alcohol. Her pulse is 136/min and irregularly irregular. An ECG shows irregularly spaced QRS complexes with no distinct P waves. Treatment is started with a drug that slows atrioventricular node conduction velocity and prevents voltage-dependent calcium entry into myocytes. The patient is at greatest risk for which of the following adverse effects?

	Answer	Image
A	Tinnitus	
B	Dry mouth	
C	Peripheral edema	
D	Bronchospasm	
E	Gingival hyperplasia	

Hint

The ECG findings in this patient (irregularly spaced QRS complexes with no distinct P waves) are consistent with atrial fibrillation. Non-dihydropyridine calcium channel blockers are commonly used in the treatment of atrial fibrillation.

Correct Answer

A - Tinnitus

Explanation Why

[Tinnitus](#) is a side effect of [quinidine](#), a [class IA antiarrhythmic drug](#) sometimes used to treat [atrial fibrillation](#). [Quinidine](#) does not affect the voltage-dependent calcium channels in the [myocardium](#), meaning it is not the drug in question. Common side effects of [quinidine](#) include [cinchonism](#), prolongation of the [QT interval](#), and [hypotension](#).

B - Dry mouth

Explanation Why

Dry mouth is a classic [anticholinergic side effect](#). However, [anticholinergic drugs](#) are not used in the management of [atrial fibrillation](#).

C - Peripheral edema

Explanation Why

Peripheral [edema](#) is a side effect of [dihydropyridine calcium channel blockers](#) (DHP [CCBs](#)) such as [nifedipine](#). DHP [CCBs](#) act mainly on vascular [smooth muscle](#), causing [vasodilation](#); this mechanism makes them most useful for treating [hypertension](#). They are not indicated for the [treatment of atrial fibrillation](#), as they have a negligible effect on the [cardiac conduction system](#) and [myocardium](#). Their [vasodilatory](#) effect accounts for the classic side effects of [edema](#) and flushing.

D - Bronchospasm

Explanation Why

[Beta blockers](#) are commonly used for rate control in atrial fibrillation and are associated with bronchospasm, especially in patients with [asthma](#) or [COPD](#). However, [beta blockers](#) do not affect calcium channels in the [myocardium](#), meaning the drug in question is not a [beta blocker](#). Other common side effects of [beta blocker](#) use include [bradyarrhythmias](#), [hypotension](#), depression, [dyslipidemia](#), and [impotence](#).

E - Gingival hyperplasia

Explanation Why

Gingival hyperplasia is a side effect of [non-dihydropyridine calcium channel blockers](#) such as [verapamil](#). In patients with [atrial fibrillation](#), [non-dihydropyridine calcium channel blockers](#) act as rate control agents through two distinct mechanisms. First, these agents slow down conduction of atrial electrical impulses through the [AV node](#). Second, inhibition of calcium entry into ventricular [myocytes](#) reduces the ventricular rate. [Verapamil](#), along with other [CCBs](#) like [nifedipine](#) and [diltiazem](#), may cause gingival hyperplasia with long-term use. Other side effects include dose-dependent [bradycardia](#), [AV block](#), and [hypotension](#).

Question # 11

A 79-year-old woman is brought to the emergency department by her husband 20 minutes after losing consciousness. She was walking briskly with her husband when she collapsed suddenly. Her husband says that she regained consciousness after 1 minute. She has had episodes of mild chest pain for the past 2 months, especially when working in the garden. Physical examination shows a grade 3/6 systolic ejection murmur. The intensity of the murmur decreases with the handgrip maneuver and does not change with inspiration. Which of the following is the most likely cause of this patient's symptoms?

	Answer	Image
A	Cystic medial degeneration of the aortic root	
B	Sterile platelet thrombi on the mitral valve	
C	Infected fibrin aggregates on the tricuspid valve	
D	Dystrophic calcification on the aortic valve	
E	Granulomatous nodules on the mitral valve	

	Answer	Image
F	Congenital leaflet fusion of the aortic valve	

Hint

Physical examination of this patient is likely to show pulsus parvus et tardus.

Correct Answer

A - Cystic medial degeneration of the aortic root

Explanation Why

[Cystic medial degeneration](#) of the aortic root, which occurs in [Marfan syndrome](#), can lead to [aortic regurgitation](#). Unlike this patient's [murmur](#), the [murmur](#) of [aortic regurgitation](#) is holodiastolic and accentuated by maneuvers that increase [afterload](#) (e.g., handgrip). Moreover, this patient does not have any features that suggest a diagnosis of [Marfan syndrome](#) (e.g., long extremities, hypermobile [joints](#), [arachnodactyly](#), blurred vision due to [lens dislocation](#)).

B - Sterile platelet thrombi on the mitral valve

Explanation Why

Sterile [platelet thrombi](#) on the [mitral valve](#) are characteristic of [nonbacterial thrombotic endocarditis \(NBTE\)](#), which can occur in individuals with rheumatologic disorders (e.g., [systemic lupus erythematosus](#)) or malignancies. [NBTE](#) can result in [mitral regurgitation](#). Unlike this patient's [murmur](#), the [murmur](#) of [mitral regurgitation](#) is [holosystolic](#). Moreover, this patient lacks manifestations that suggest an underlying rheumatologic disorder (e.g., [fever](#), [joint pain](#), malaise) and does not have a known [malignancy](#).


C - Infected fibrin aggregates on the tricuspid valve

Explanation Why

Infected [fibrin](#) aggregates on [heart valves](#) are seen in individuals with [infective endocarditis \(IE\)](#). [Tricuspid valve](#) involvement (e.g., from intravenous drug use) results in [tricuspid regurgitation](#). Unlike this patient's [murmur](#), the [murmur](#) of [tricuspid regurgitation](#) is [holosystolic](#). Moreover, this patient does not have typical manifestations of IE (e.g., [fever](#), malaise).

D - Dystrophic calcification on the aortic valve

Image



Explanation Why

The most common cause of [aortic valve stenosis](#) is [dystrophic calcification](#) of the [aortic valve](#). [Aortic valve stenosis](#) is the most common valvular abnormality in adults > 65 years of age. Typical symptoms include [syncope](#) (especially during physical exercise), [angina pectoris](#), and [heart failure](#) due to decreased [ejection fraction](#). [Cardiac auscultation](#) in a patient with [aortic stenosis](#) typically reveals a mid-to-late-peaking [systolic](#), crescendo-decrescendo [murmur](#) that is heard best at the [aortic area](#) (right upper sternal border). Maneuvers that increase [afterload](#) (e.g., sustained handgrip) decrease the intensity of the [murmur](#).

E - Granulomatous nodules on the mitral valve

Explanation Why

Granulomatous nodules on the [mitral valve](#) are associated with [rheumatic heart disease](#) (RHD) and can result in [mitral regurgitation](#). RHD typically manifests during [adolescence](#) or early adulthood and is an unlikely underlying cause of a [murmur](#) in this 79-year-old patient. Moreover, the handgrip maneuver increases the intensity of the [murmur](#) of [mitral regurgitation](#).

F - Congenital leaflet fusion of the aortic valve



Explanation Why

A [bicuspid aortic valve](#) due to congenital fusion of the [aortic valve](#) leaflets can lead to [aortic stenosis](#) and manifest with [syncope](#), [chest pain](#), and a late-peaking [systolic murmur](#) that decreases in intensity with the handgrip maneuver. However, [aortic stenosis](#) due to a [bicuspid valve](#) typically becomes symptomatic in the fourth or fifth decade of life. This 79-year-old patient is more likely to have [aortic stenosis](#) of a different etiology.

Question # 12

A 65-year-old man with hypertension comes to the physician for a routine health maintenance examination. Current medications include atenolol, lisinopril, and atorvastatin. His pulse is 86/min, respirations are 18/min, and blood pressure is 145/95 mm Hg. Cardiac examination is shown. Which of the following is the most likely cause of this physical examination finding?

[Video](#)

	Answer	Image
A	Decreased compliance of the left ventricle	 The image contains four vertically stacked graphs. The top graph shows pressure (mmHg) vs. time, with curves for aortic pressure (orange), left ventricular pressure (blue), and left atrial pressure (green). The middle graph shows ventricular volume (ml) vs. time, with curves for stroke volume (blue) and end-diastolic volume (red). The bottom two graphs show ECG (purple) and heart sounds (red). The graphs illustrate that in decreased compliance, the end-diastolic volume is reduced (point 3) and the pressure is higher for a given volume (point 2).
B	Myxomatous degeneration of the mitral valve	 The image shows two vertically stacked graphs. The top graph is an ECG (blue) and the bottom graph is a Phonocardiogram (red). The ECG shows a normal sinus rhythm. The Phonocardiogram shows a mid-systolic click (indicated by an arrow) and a late systolic murmur (indicated by a shaded area). The graphs are labeled with Diastole, Systole, and Diastole. The mid-systolic click is a characteristic finding in myxomatous degeneration of the mitral valve.
C	Inflammation of the pericardium	

	Answer	Image
D	Dilation of the aortic root	The image displays four panels (A, B, C, D) comparing normal aortic valve closure (OS) with aortic root dilation. Each panel shows an ECG trace (blue) and a Phonocardiogram (PCG) trace (orange). The PCG traces are synchronized with the ECG, showing the aortic valve closure (OS) during diastole. In panels A and B, the OS is sharp and occurs early in diastole. In panels C and D, the OS is prolonged and decelerated, indicating aortic root dilation.
E	Thickening of the mitral valve leaflets	The image displays a single panel showing an ECG trace (blue) and a Phonocardiogram (PCG) trace (orange). The PCG trace is synchronized with the ECG, showing the mitral valve closure (OS) during diastole. The OS is prolonged and decelerated, indicating mitral valve disease.


Hint

The auscultatory finding in this patient is an S₄ gallop, which occurs as the atria pump blood into the ventricles. An S₄ gallop is almost always pathological.

Correct Answer

A - Decreased compliance of the left ventricle

Image




Explanation Why

Decreased compliance of the [left ventricle](#) is most likely due to [left ventricular hypertrophy](#) caused by long-standing [hypertension](#) and often manifests with an [S4 gallop](#). Increased blood pressure activates [angiotensin II](#) either by local mechanical stretching of myocytes or as a result of the activation of the [renin-angiotensin-aldosterone](#) system. [Angiotensin II](#) promotes concentric remodeling of the [heart](#), increasing stiffness of the organ, thus, decreasing compliance.

B - Myxomatous degeneration of the mitral valve

Image



Explanation Why

Myxomatous degeneration of the [mitral valve](#) occurs in some [connective tissue diseases](#) (e.g., [Marfan syndrome](#), Ehler-Danlos syndrome) leading to eventual [mitral valve prolapse \(MVP\)](#). MVP is associated with late-[systolic](#) crescendo [murmurs](#) as blood backflows from the [left ventricle](#) to the [left atrium](#), followed by a midsystolic click due to tensing of the chordae tendinae. The [mitral regurgitation](#) and volume overload associated with MVP can also result in an [S3](#), which is an early [diastolic](#) sound, but MVP is not typically associated with an S4, as seen here.

C - Inflammation of the pericardium


Explanation Why

[Pericardial inflammation](#) ([pericarditis](#)), which is typically caused by viral infections, would result in

an auscultatory finding ([pericardial friction rub](#)) but this sound would be heard in both [diastolic](#) and [systolic](#) portions of the [cardiac cycle](#) and it would be best heard along the left sternal border when the patient sits upright, leans forward, and holds his breath. The sound that is heard here is an [S4 gallop](#), which is an unusual finding in [pericarditis](#). Moreover, patients with [pericarditis](#) are rarely asymptomatic; they would usually complain of [chest pain](#) and/or breathlessness. It is, therefore, unlikely that this patient has [pericarditis](#).

D - Dilation of the aortic root

Image




Explanation Why

Dilation of the aortic root, which is caused by age-related degenerative changes, [hypertension](#), [aortitis](#) (e.g., [tertiary syphilis](#)), or [connective tissue disease](#) (e.g., [Marfan syndrome](#)), causes chronic [aortic insufficiency](#) (AI). AI can result in a low-pitched, mid or late [diastolic](#) sound that is best heard at the cardiac apex due to vibration of the ventricular wall by the regurgitant jet. However, a patient with AI would also have an early [diastolic murmur](#) best heard in the right upper sternal border. An isolated low-pitched, late [diastolic](#) sound on [cardiac examination](#) suggests another cause.

E - Thickening of the mitral valve leaflets

Image



Explanation Why

Thickening of the mitral leaflet is seen in [mitral stenosis](#) and is the cause of the [opening snap](#), which is an early [diastolic](#) sound that is best heard at the apex in the left [lateral](#) decubitus position. However, the [opening snap](#) is a high-pitched sound, unlike the low-pitched sound heard here, and it would be accompanied by the late-[diastolic](#), decrescendo [murmur](#) of [mitral stenosis](#).

Question # 13

A 58-year-old woman with type 2 diabetes mellitus comes to the physician because of a 3-month history of right lower extremity pain and burning while walking. The pain subsides with rest. She has smoked one pack of cigarettes daily for the past 30 years. Her current medications include metformin, atorvastatin, and aspirin. Examination shows a lack of hair and decreased skin temperature over the right foot. The right pedal pulse is not palpable. The physician adds a drug to her regimen that causes vasodilation and inhibits the aggregation of platelets and the proliferation of smooth muscle cells. Which of the following drugs was most likely added?

	Answer	Image
A	Dabigatran	
B	Eptifibatide	
C	Bosentan	
D	Clopidogrel	
E	Cilostazol	

Hint

This patient presents with right lower extremity claudication, trophic skin changes (decreased temperature and lack of hair), and absent pedal pulse. These features are suggestive of peripheral arterial disease (PAD).

Correct Answer

A - Dabigatran

Explanation Why

[Dabigatran](#) is an [anticoagulant](#) that directly inhibits [thrombin](#), thereby disrupting the [coagulation cascade](#). Unlike the drug used in this patient, [dabigatran](#) does not affect [platelet aggregation](#), prevent [smooth muscle proliferation](#), or cause [vasodilation](#). Moreover, it is not indicated to treat [PAD](#). [Dabigatran](#) can be used to manage [venous thromboembolism](#), [atrial fibrillation](#), or other conditions requiring long-term anticoagulation.

B - Eptifibatide

Explanation Why

[Eptifibatide](#) exerts its antiplatelet activity through [glycoprotein receptor IIb/IIIa](#) inhibition. While it does inhibit [platelet aggregation](#), [eptifibatide](#) does not cause [vasodilation](#) or inhibit [smooth muscle proliferation](#). This drug is used to treat [unstable angina](#) and to prevent adverse cardiac events during [percutaneous coronary intervention](#).

C - Bosentan

Explanation Why

[Bosentan](#) blocks vascular endothelin receptors, causing [vasodilation](#). It is primarily used to reduce [pulmonary vascular resistance](#) in patients with [pulmonary hypertension](#). Unlike the drug administered here, [bosentan](#) does not prevent [platelet aggregation](#) and is not used to treat [PAD](#).

D - Clopidogrel

Explanation Why

[Clopidogrel](#) is an [antiplatelet agent](#) that acts through [ADP](#) receptor blockade ([ADP receptor inhibitor](#)). While [clopidogrel](#) reduces [platelet aggregation](#), it does not cause [vasodilation](#) or inhibit [smooth muscle proliferation](#). Furthermore, this patient is already on an [antiplatelet agent](#) ([aspirin](#)), and there is no indication for dual antiplatelet therapy (e.g., endovascular placement of a peripheral arterial stent or previous bypass with synthetic graft).


E - Cilostazol

Explanation Why

[Cilostazol](#) is indicated in patients with [PAD](#) to help improve the symptoms of [intermittent claudication](#). The drug acts by inhibiting phosphodiesterase 3, causing an increased intracellular concentration of [cAMP](#) that results in vascular [smooth muscle](#) relaxation and [vasodilation](#). [Cilostazol](#) also reduces [platelet aggregation](#). Common side effects include [headache](#), [diarrhea](#), [palpitations](#), and [tachycardia](#).

Question # 14

A 65-year-old man comes to the physician because of a 10-month history of crampy left lower extremity pain that is exacerbated by walking and relieved by rest. The pain is especially severe when he walks on an incline. He has a 20-year history of type 2 diabetes mellitus, for which he takes metformin. He has smoked 1 pack of cigarettes daily for 40 years. His blood pressure is 140/92 mm Hg. Physical examination shows dry and hairless skin over the left foot. Which of the following is the most likely underlying cause of this patient's symptoms?

	Answer	Image
A	Osteophytic compression of the lumbar spinal canal	
B	Thrombosing vasculitis of the popliteal artery	
C	Intimal plaque in the posterior tibial artery	
D	Fibrin clot in the left popliteal vein	
E	Microvascular compromise of the superficial peroneal nerve	
F	Systemic hyperplastic arteriolosclerosis	

Hint

Ankle-brachial index (ABI) is a common test used to diagnose this condition.

Correct Answer

A - Osteophytic compression of the lumbar spinal canal

Explanation Why

Osteophytic compression of the lumbar [spinal canal](#) may lead to [neurogenic claudication](#), causing lower leg [pain](#) on exertion. However, [neurogenic claudication](#) is relieved by [flexion](#) at the waist, so classically it is less severe when walking uphill. Moreover, this condition does not cause trophic [skin](#) changes such as hairlessness of the foot.

B - Thrombosing vasculitis of the popliteal artery

Explanation Why

Thrombosing [vasculitis](#) of the [popliteal artery](#) could be seen in thromboangiitis obliterans (TAO). TAO is one of the possible causes of lower extremity [claudication](#), and smoking is strongly associated with this condition. However, the symptoms of TAO are commonly bilateral, whereas this patient's symptoms are unilateral. Moreover, the condition classically manifests with [superficial thrombophlebitis](#), [Raynaud phenomenon](#), and early [ischemic](#) ulceration, none of which are present in this case. Finally, TAO typically manifests in patients < 45 years of age, making the diagnosis less likely in this 65-year-old man.

C - Intimal plaque in the posterior tibial artery

Image



Explanation Why

An [intimal](#) plaque in the [posterior tibial artery](#) is the most likely cause of this patient's presentation, which is consistent with [peripheral arterial disease \(PAD\)](#). [PAD](#) is strongly associated with [diabetes](#), [hypertension](#), and smoking, all of which are present in this case. The disease begins with the buildup of lipids and fibrous material beneath the [tunica intima](#) in peripheral [arteries](#). With time, focal accumulation of these substances can cause stenosis and reduced blood flow to downstream tissues, manifesting as episodic [ischemia](#) (in this case, [claudication](#)). Typical [physical exam](#) findings include decreased or absent peripheral pulses, [atrophic skin](#) changes (i.e., dry, shiny, or [hairless skin](#)), or chronically non-healing foot wounds. The diagnosis is confirmed via [ankle-brachial index](#).

D - Fibrin clot in the left popliteal vein

Explanation Why

A [fibrin](#) clot in the left [popliteal vein](#) would manifest as a [deep venous thrombosis \(DVT\)](#). Although smoking and [diabetes](#) are [risk factors](#) for the development of [DVT](#), this condition manifests with acute onset unilateral leg [pain](#), swelling, and [erythema](#). There is most often a history of surgery or inactivity as well that causes [venous stasis](#), neither of which are present in this case. Furthermore, [DVT](#) does not cause [claudication](#), pedal [alopecia](#), or the foot to turn pale on elevation.

E - Microvascular compromise of the superficial peroneal nerve

Explanation Why

[Microvascular](#) compromise of the [superficial peroneal nerve](#) (or any other sensory nerve of the [distal](#) lower extremity) occurs with long-standing [diabetes](#). Through a complex series of pathways, chronic [hyperglycemia](#) induces [microvascular](#) damage in the vessels of the [perineurium](#), causing axonal damage that manifests as [diabetic neuropathy](#). This patient is at significant risk to develop neuropathy as a complication of his [diabetes](#). However, he presents with intermittent unilateral lower extremity [pain](#) and evidence of [skin](#) changes, neither of which are consistent with [diabetic neuropathy](#). Moreover, he does not report [pain](#), burning, tingling, or numbness in the [dorsal](#) foot and toes (the [dermatome](#) of the [superficial peroneal nerve](#)), all of which are characteristic symptoms of [diabetic neuropathy](#).

F - Systemic hyperplastic arteriolosclerosis


Explanation Why

Systemic [hyperplastic arteriosclerosis](#) is a form of small vessel disease that typically arises in the setting of severe [hypertension](#) ($\geq 180/120$ mmHg). The [intimal](#) response to sustained blood pressures in this range results in a characteristic “onion [skin](#)” appearance on histological examination of vascular tissue. This process classically causes [ischemic](#) damage in the [kidneys](#), but it is not implicated in cases of vascular insufficiency in the extremities. Moreover, this patient's blood pressure is only 140/92 mm Hg, which is much lower than what is necessary to cause [hyperplastic](#)

[arteriosclerosis.](#)

Question # 15

A 39-year-old man comes to the physician because of a 2-month history of shortness of breath with exertion. He immigrated to the United States from Ghana 20 years ago. His pulse is 88/min, respirations are 18/min, and blood pressure is 120/80 mm Hg. Echocardiography shows thickened mitral valve leaflets with reduced motion and a narrowed mitral valve orifice diameter. Which of the following auscultation findings is most likely to correlate with the severity of this patient's valvular disease?

	Answer	Image
A	Decreased time between S2 and opening snap	
B	Presence of a mid-systolic click	
C	Radiation of murmur to the axilla	
D	Increased intensity of the murmur	
E	Absence of A2 heart sound	


Hint

This patient's echocardiography findings are consistent with mitral stenosis.

Correct Answer

A - Decreased time between S2 and opening snap

Image



Explanation But

[Atrial fibrillation](#) and [pulmonary hypertension](#) are also associated with a poor prognosis in patients with MS.

Explanation Why

A decrease in the time between the aortic component of S2 and the [opening snap](#) (OS) occurs due to increasing left atrial pressure (LAP) brought on by worsening [mitral stenosis](#) (MS). The greater the LAP, the faster the acceleration of valve leaflets to the maximal point of opening, and the sooner the [opening snap](#) occurs after S2. The most common cause of MS in developing countries is [rheumatic fever](#).

B - Presence of a mid-systolic click

Explanation Why

The presence of a mid-[systolic click](#) followed by a late [systolic murmur](#) is associated with [mitral valve prolapse](#). The click occurs as the chordae tendinae anchoring the prolapsed valve leaflet reach their maximum length and snap tight. However, this patient's [echocardiography](#) findings are consistent with [mitral stenosis](#), which does not manifest with a mid-[systolic click](#).

C - Radiation of murmur to the axilla

Explanation Why

Radiation of a [murmur](#) to the [axilla](#) is most noticeable in [mitral regurgitation](#). However, this patient's [echocardiography](#) is consistent with [mitral stenosis](#), which is best heard from the apex and is not known to radiate.

D - Increased intensity of the murmur

Explanation Why

An increased intensity of a [murmur](#) is correlated with increased disease severity in conditions such as [mitral regurgitation](#) or [tricuspid regurgitation](#). However, this patient's [echocardiography](#) findings are consistent with [mitral stenosis](#), in which [murmur](#) intensity has no [correlation](#) with disease severity. In patients with very severe [mitral stenosis](#) the [murmur](#) may even be inaudible due to the severe limitation of blood flow across the stenotic valve.

E - Absence of A2 heart sound

Explanation Why

Absence of the A2 component of the S2 [heart sound](#) can be a sign of worsening [aortic stenosis](#). S2 is comprised of a louder aortic contribution (A2) and a softer pulmonary contribution (P2). As stenosis of the aortic [semilunar valves](#) worsens, the A2 diminishes until it is no longer audible. However, this patient's [echocardiography](#) is consistent with [mitral stenosis](#), in which there is no change in S2 as the disease increases in severity.

Question # 16

A 77-year-old woman with congestive heart failure is admitted to the hospital for evaluation prior to cardiac transplantation. During her stay at the hospital, the physician prescribes a drug to improve cardiac contractility. The drug works by selectively inhibiting an isoenzyme that is responsible for the degradation of cyclic adenosine monophosphate. Which of the following is the most likely adverse effect of this drug?

	Answer	Image
A	QT interval prolongation	
B	Hyperkalemia	
C	Hypotension	
D	Bradycardia	
E	Hyperglycemia	
F	Bronchospasm	

Hint

The drug that was most likely prescribed by the physician is milrinone, a selective phosphodiesterase type 3 inhibitor that has positive inotropic and chronotropic effects on the heart. Milrinone improves cardiac contractility by increasing cAMP concentration in cardiomyocytes, which stimulates calcium influx into the cells.

Correct Answer

A - QT interval prolongation

Explanation Why

[QT prolongation](#) is a potential side effect of a number of cardiotropic drugs, including [class IA antiarrhythmic](#) and [class III antiarrhythmic drugs](#), but it is rarely triggered by [PDE3 inhibitors](#) such as [milrinone](#) (< 1% of patients).

B - Hyperkalemia

Explanation Why

[Hyperkalemia](#) is a potential adverse effect of ACE inhibitors, [ARBs](#), [digoxin](#), and [potassium-sparing diuretics](#). [PDE3 inhibitors](#) such as [milrinone](#) can cause [hypokalemia](#) (< 1% of patients), but they are not associated with [hyperkalemia](#).

C - Hypotension

Explanation But

[Dobutamine](#), which acts on adrenergic [β1 receptors](#), is also used as an [inotropic](#) agent in the treatment of [decompensated heart failure](#) and [cardiogenic shock](#). It is preferred in patients with [kidney](#) dysfunction because, unlike [milrinone](#), it is not metabolized by the [kidneys](#).

Explanation Why

[Hypotension](#) is a potential side effect of [PDE inhibitors](#) such as [milrinone](#). In peripheral vessels, an increase in [cAMP](#) mediates [smooth muscle](#) relaxation by inhibiting [myosin light chain kinase](#), which results in peripheral [vasodilation](#). Other side effects of [milrinone](#) include [tachycardia](#), [ventricular arrhythmias](#), [headache](#), and GI upset. [Milrinone](#) serves as a bridge to [cardiac transplant](#) and is also indicated in the treatment of advanced [congestive heart failure \(CHF\)](#) and [cardiogenic shock](#).

D - Bradycardia

Explanation Why

[Bradycardia](#) is a potential side effect of a number of cardiotropic drugs, including [beta blockers](#), [digoxin](#), and [non-dihydropyridine calcium channel blockers](#). [PDE3 inhibitors](#) such as [milrinone](#) are associated with [tachycardia](#) rather than [bradycardia](#) due to their positive [chronotropic](#) effects.

E - Hyperglycemia

Explanation Why

[Hyperglycemia](#) is a potential adverse effect of [epinephrine](#), which also improves cardiac contractility by increasing [cAMP](#) concentration in [cardiomyocytes](#). However, [epinephrine](#) increases [cAMP](#) concentration by stimulating [adenylate cyclase](#) activity rather than decreasing the degradation of [cAMP](#). Moreover, [epinephrine](#) is not indicated in the treatment of [decompensated CHF](#) because it is associated with increased [mortality](#) in this patient group.

F - Bronchospasm

Explanation Why

Bronchospasm is a potential side effect of a number of cardiotropic drugs, including adenosine and non-selective [beta blockers](#), but it is rarely triggered by [PDE3 inhibitors](#) such as [milrinone](#) (< 1% of patients).

Question # 17

An 8-year-old boy is brought to the physician by his father because of abnormal movements of his limbs. For the past four days, he has had uncontrolled jerking movements of his arms and legs and has been dropping cups and toys. His symptoms are worse when he is excited and improve while he is asleep. During the same time period, he has become increasingly irritable and tearful. He had a sore throat 6 weeks ago that resolved without treatment. His temperature is 37.3°C (99.2°F). Examination shows occasional grimacing with involuntary jerking movements of his limbs. Muscle strength and tone are decreased in all extremities. When he grips the physician's index and middle fingers with his hands, his grip increases and decreases continuously. This patient is at increased risk for which of the following complications?

	Answer	Image
A	Extraneural metastasis	
B	Early-onset dementia	
C	Attention deficit hyperactivity disorder	
D	Diabetes mellitus	
E	Mitral regurgitation	

Hint

In a pediatric patient with a recent history of a sore throat, random jerking movements, emotional lability, hypotonia, and a “milkmaid's grip” most likely indicate Sydenham chorea.

Correct Answer

A - Extraneural metastasis

Explanation Why

Extraneural [metastasis](#) of a primary [brain tumor](#) such as [medulloblastoma](#) is rare. Large [brain tumors](#) can also present with gait abnormalities but typically would also be accompanied by [features of increased intracranial pressure](#) (e.g., [headaches](#), early morning vomiting, [papilledema](#)), which are not seen in this patient. Additionally, repetitive jerking movements shortly after a pharyngeal infection is most consistent with [Sydenham chorea](#), which is not due to [malignancy](#) and so not associated with [metastasis](#).

B - Early-onset dementia

Explanation Why

[Huntington disease](#) is an [autosomal-dominant](#) neurocognitive disease that is characterized by repetitive movement disorders ([chorea](#)), psychiatric symptoms, and early-onset [dementia](#). It is most commonly seen in patients with a [family history](#) of the disease. Although this patient's repetitive and jerky movements are typical of [chorea](#), the onset of symptoms shortly after a pharyngeal infection suggest [Sydenham chorea](#). [Sydenham chorea](#) is not associated with increased risk of early-onset [dementia](#).

C - Attention deficit hyperactivity disorder

Explanation Why

[Attention deficit hyperactivity disorder](#) is strongly associated with [Tourette syndrome](#), which is a neurological movement disorder characterized by multiple motor [tics](#) and at least one vocal [tic](#). This patient lacks vocal [tics](#) and the motor [tics](#) in [Tourette syndrome](#) usually involves the facial muscles rather than the upper and the lower limbs seen here. This patient's repetitive movements and emotional lability are more consistent with [Sydenham chorea](#), which is not associated with [ADHD](#).

D - Diabetes mellitus

Explanation Why

[Friedreich ataxia](#) is a neurodegenerative disease that most commonly manifests with [diabetes mellitus](#), neurological dysfunction, and [hypertrophic cardiomyopathy](#). The neurological dysfunction seen in [Friedreich ataxia](#) may include weakness, poor coordination, and staggering gait but is not characterized by [choreiform movements](#) that improve with rest. [Sydenham chorea](#), which this patient has, is not associated with the development of [diabetes mellitus](#).


E - Mitral regurgitation

Explanation Why

[Rheumatic fever](#) is a complication of untreated pharyngeal infection with [Streptococcus pyogenes](#) (GAS), which can manifest with [fever](#), arthralgias, [Sydenham chorea](#) (as seen here), and cardiac valve disease, of which [mitral valve](#) involvement is most common. Valve disease occurs as a result of [antibodies](#) against bacterial M [proteins](#) cross-reacting with valvular self-antigens through a process known as [molecular mimicry](#), which results in valvular destruction. The mechanism of [Sydenham chorea](#) is thought to be similar and result from [antibody cross-reactivity](#) with [basal ganglia](#) structures.

Question # 18

A 22-year-old woman comes to the physician for gradual worsening of her vision. Her father died at 40 years of age. She is 181 cm (5 ft 11 in) tall and weighs 69 kg (152 lb); BMI is 21 kg/m². A standard vision test shows severe myopia. Genetic analysis shows an *FBNI* gene mutation on chromosome 15. This patient is at greatest risk of mortality due to which of the following causes?

	Answer	Image
A	Left ventricular outflow obstruction	
B	Obstruction of the superior vena cava lumen	
C	Increased pressure in the pulmonary arteries	
D	Eccentric ventricular hypertrophy	
E	Supraventricular tachyarrhythmia	
F	Intimal tear of the aortic root	 Marfan syndrome Epidemiology Prevalence: 1-5/10,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15; autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissection) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups) Clinical features: - Ectopia lentis, severe myopia - Heart valve defects - Pectus deformity (pectus carinatum or pectus excavatum) - Aortic disease (e.g., aortic dissection, aortic aneurysm) - Anachnodyly: Joint hypermobility - Hyperextensible skin - Tall stature - Berry aneurysms - High-arched palate - Spinal deformities (scoliosis, hyperkyphosis) - Long extremities - Pes planus, hindfoot valgus

Hint

Other abnormal morphometric features of this condition include thoracic wall deformities (e.g., pectus carinatum, pectus excavatum), long extremities, arachnodactyly, subluxation of the lens, and hypermobile joints.

Correct Answer

A - Left ventricular outflow obstruction

Explanation Why

Left ventricular outflow tract obstruction is an important cause of [mortality](#) in patients with [hypertrophic cardiomyopathy](#) (HCM). Hereditary HCM can be caused by an [autosomal dominant](#) defect in [genes](#) responsible for the synthesis of certain [sarcomeric proteins](#) (e.g., MYH7, MYBPC3) or by an [autosomal recessive trinucleotide repeat](#) mutation on the [frataxin gene](#) on [chromosome 9](#) ([Friedreich ataxia](#)). A mutation in the [FBN1 gene](#) is not associated with an increased risk of HCM.

B - Obstruction of the superior vena cava lumen

Explanation Why

[SVC syndrome](#) is most commonly seen in patients with a [mediastinal mass](#) (e.g., [lymphoma](#)), [Pancoast tumor](#), or thrombosis due to an indwelling venous catheter. [FBN1 gene](#) defects are not associated with an increased risk of [SVC syndrome](#).

C - Increased pressure in the pulmonary arteries

Explanation Why

Chronic [pulmonary arterial hypertension](#) (PAH) can lead to [cor pulmonale](#) and [right ventricular failure](#), resulting in death. Genetic causes of PAH include [inactivating mutations](#) on the [BMPR2 gene](#) on [chromosome 2](#) and [connective tissues](#) disease such as [systemic sclerosis](#). However, mutations on the [FBN1 gene](#) are not associated with the development of [pulmonary hypertension](#).

D - Eccentric ventricular hypertrophy

Explanation Why

Eccentric ventricular [hypertrophy](#) is the underlying disease process in [dilated cardiomyopathy](#). Genetic causes of [dilated cardiomyopathy](#) include certain [sarcomeric](#) protein mutations (e.g., TTN gene), [Duchenne muscular dystrophy](#), and hereditary [hemochromatosis](#). Mutations in the [FBN1 gene](#) are not associated with an increased risk of [dilated cardiomyopathy](#).


E - Supraventricular tachyarrhythmia

Explanation Why

[Supraventricular tachycardia \(SVT\)](#) refers to any [tachyarrhythmia](#) that originates above or within the [Bundle of His](#). Causes include a dysfunctional [AV node](#), the presence of an additional electrical pathway ([AVNRT](#)), and the presence of an [accessory pathway](#) between the [atria](#) and ventricles ([AVRT](#), [WPW](#)). This patient's condition is not associated with an increased risk of [supraventricular tachycardia](#).

F - Intimal tear of the aortic root

Image



Explanation But


Severe [mitral valve prolapse](#) is another common cause of early [mortality](#) in patients with [Marfan syndrome](#).

Explanation Why


An [autosomal dominant](#) defect in the [FBN1 gene](#) on [chromosome 15](#) is responsible for [Marfan syndrome](#), which can cause tall stature and [myopia](#), as seen in this patient. [Marfan syndrome](#) is associated with [cystic medial necrosis](#) of the aorta, which can result in aortic root dissection (e.g., [intimal tear](#)), [aortic root dilation](#), and [aortic regurgitation](#). Aortic root disease is the most common cause of [morbidity](#) and [mortality](#) in patients with [Marfan syndrome](#).

Question # 19

A 9-year-old boy from Eritrea is admitted to the hospital for lethargy and increased work of breathing. He has had recurrent episodes of fever, shortness of breath, and fatigue over the past 3 years. His pulse is 132/min and blood pressure is 90/66 mm Hg. Pulse oximetry on room air shows an oxygen saturation of 82%. Auscultation of the chest shows coarse crackles in both lungs and a diastolic murmur at the cardiac apex. Despite appropriate lifesaving measures, he dies. A photomicrograph of a section of myocardium obtained at autopsy is shown. Which of the following is the most likely underlying cause of this patient's cardiac disease?



	Answer	Image
A	Amastigote infiltration	
B	Beta-myosin heavy chain defect	

	Answer	Image
C	Caseating granulomatous inflammation	
D	Type II hypersensitivity reaction	
E	Exotoxin-mediated myonecrosis	

Hint

The photomicrograph shows an Aschoff body, which is composed of T cells, plasma cells, and cardiac histiocytes (Anitschkow cells) surrounding degenerated myofibers. Aschoff bodies are characteristic of rheumatic heart disease.

Correct Answer

A - Amastigote infiltration

Explanation Why

Infiltration of the [myocardium](#) by [amastigotes](#) is seen in patients with [Chagas disease](#), which is caused by the protozoan parasite [Trypanosoma cruzi](#). [Chagas disease](#) can manifest with [fever](#) (early sign), [dyspnea](#), [tachycardia](#), fatigue, [hypoxia](#), and/or a [heart murmur](#) (later stages) and is characterized by eccentric cardiac [hypertrophy](#) with apical [atrophy](#). However, an [Aschoff body](#) would not be expected on histological examination. Moreover, [Chagas disease](#) is [endemic](#) to Central and South America, not Eritrea.


B - Beta-myosin heavy chain defect

Explanation Why

A beta-[myosin heavy chain](#) defect is consistent with [hypertrophic cardiomyopathy \(HOCM\)](#), which can also manifest with [dyspnea](#), [tachycardia](#), fatigue, a decreased [oxygen saturation](#), and a [heart murmur](#). However, [HOCM](#) does not manifest with [fever](#), and histological examination would show myofibrillar [fibrosis](#) and disarray, not an [Aschoff body](#).

C - Caseating granulomatous inflammation

Image



Explanation Why

[Caseating granulomatous inflammation](#) is characteristic of [tuberculosis](#), which should be suspected in a patient with a history of immigration, chronic [fever](#), [dyspnea](#), and fatigue, as seen here. Also, [caseating granulomas](#) have similar histological features, e.g., [lymphocytes](#) and histiocytes surrounding an area of central [necrosis](#). However, [myocardial](#) involvement is extremely rare in [tuberculosis](#). Moreover, the central [necrotic](#) region in [caseating granulomas](#) is composed of amorphous or granular [necrotic](#) debris, not the degenerated wavy fibers seen here.

D - Type II hypersensitivity reaction

Explanation Why

[Rheumatic fever](#) is an inflammatory sequela of untreated infection with group A [streptococcus](#) that

can involve the [heart](#) ([rheumatic heart disease](#)), [joints](#), [skin](#), and [CNS](#). The exact pathogenic mechanism that causes [rheumatic fever](#) is not fully understood, but it is believed to be due to a [type II hypersensitivity reaction](#) induced by [streptococcal proteins](#). The [antibodies](#) formed then react with cardiac [myosin proteins](#) due to [molecular mimicry](#) between [streptococcal](#) M protein and human cardiac [myosin](#), causing tissue [inflammation](#) and injury.


E - Exotoxin-mediated myonecrosis

Explanation Why

[Exotoxin](#)-mediated myonecrosis is characteristic of [myocarditis](#) caused by [Corynebacterium diphtheriae](#) infection. Diphtheritic [myocarditis](#) can manifest with [fever](#), [dyspnea](#), [tachycardia](#), fatigue, [hypoxia](#), and a [heart murmur](#). However, an autopsy would show a distorted [myocardium](#) with loss of striations and broad areas of granular degeneration, not an [Aschoff body](#).

Question # 20

A 79-year-old homeless man is brought to the emergency department by ambulance 30 minutes after being found unresponsive by the police. On arrival, he is apneic and there are no palpable pulses. Despite appropriate life-saving measures, he dies. Examination of the heart during autopsy shows normal ventricles with a sigmoid-shaped interventricular septum. A photomicrograph of a section of the heart obtained at autopsy is shown. Which of the following is the most likely underlying cause for the structure indicated by the arrow?



	Answer	Image
A	Accumulation of iron granules	
B	Clumping of defective mitochondria	
C	Oxidation of phospholipid molecules	

	Answer	Image
D	Aggregation of alpha-synuclein	
E	Deposition of wild-type transthyretin	

Hint

The yellow-brown perinuclear inclusions indicated by the arrow are lipofuscin granules.

Correct Answer

A - Accumulation of iron granules

Explanation Why

Accumulation of [iron](#) granules would be consistent with [hemochromatosis](#), which can also affect the [heart](#) and also appear as brown, sarcoplasmic perinuclear granules. However, other characteristic findings consistent with [hemochromatosis](#) are absent, such as [fibrosis](#) and [cardiomegaly](#) due to restrictive or [dilated cardiomyopathy](#).

B - Clumping of defective mitochondria

Explanation But

The [mitochondrial](#) accumulation that is seen in [mitochondrial myopathy](#) stains red with Gomori Trichrome stain, hence the commonly used descriptive term “[ragged red fibers](#)”.

Explanation Why

Clumping of defective [mitochondria](#) is seen in [mitochondrial myopathy](#), a rare genetic disorder that mainly affects organs with a high energy requirement including the [heart](#). However, [examination of the heart](#) would show ventricular [hypertrophy](#) and/or dilatation, not a sigmoid-shaped interventricular septum. Moreover, the histological examination would show subsarcolemmal and intermyofibrillar [mitochondria](#) accumulation rather than the yellow-brown perinuclear inclusions seen here.

C - Oxidation of phospholipid molecules

Explanation Why

[Lipofuscin](#) granules, also known as wear-and-tear granules, are composed of yellow-brown pigmented granules and occur as a natural process with age. They are formed by the oxidation of phospholipid molecules and can be found in various organs (e.g., [heart](#), [liver](#), [kidneys](#)). [Lipofuscin](#)

accumulates with age because it is never completely eliminated. Age-related mechanisms that can accelerate the accumulation of [lipofuscin](#) include the enhancement of [autophagocytosis](#), a decline in intralysosomal degradation, and/or a decrease in exocytosis. A sigmoid-shaped interventricular septum is also a (physiological) sign of [aging](#).

D - Aggregation of alpha-synuclein

Explanation Why

Aggregation of alpha-synuclein leads to neuronal death and is seen in conditions such as [Parkinson disease](#), [MSA](#), and [Lewy body dementia](#). These [neurodegenerative disorders](#) mainly affect the brain and not the [heart](#), which is the organ involved in this patient. Moreover, histological examination with H&E staining would show eosinophilic (pink) [cytoplasmic](#) inclusions rather than the yellow-brown pigmented inclusions seen here.


E - Deposition of wild-type transthyretin

Explanation Why

Deposition of wild-type [transthyretin](#) is seen in senile amyloidosis and mainly affects the [myocardium](#) in elderly patients. However, senile amyloidosis would manifest with [cardiomegaly](#) due to [myocardial](#) thickening, not with a sigmoid-shaped interventricular septum. Moreover, histological examination with H&E staining would show [amyloid proteins](#), which appear as glassy, eosinophilic, homogeneous deposits, not yellow-brown perinuclear inclusions.

Question # 21

A 52-year-old woman comes to the physician because of a 3-month history of worsening chest pain and breathlessness during exercise. She has no history of serious illness and takes no medications. Vital signs are within normal limits. Auscultation of the chest shows a murmur in the 2nd right intercostal space. A phonocardiogram of the murmur is shown. Which of the following is the most likely underlying cause of this patient's symptoms?



	Answer	Image
A	M protein cross-reactivity	
B	Metastatic valvular calcification	

	Answer	Image
C	Congenital leaflet fusion	The image displays two cardiac waveforms. The top waveform is an Electrocardiogram (ECG) with labels for Diastole, Systole, and Diastole. The bottom waveform is a Phonocardiogram with labels for S1 and S2. The ECG shows aortic regurgitation (aortic valve closure) and the Phonocardiogram shows aortic stenosis (aortic valve opening).
D	Cystic medial necrosis	Marfan syndrome Epidemiology Prevalence: 1-5/10,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15, autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissections) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups) Clinical features shown in the infographic include: Ectopia lentis, severe myopia; Heart valve defects; Pectus deformity (pectus carinatum or pectus excavatum); Aortic disease (e.g., aortic dissection, aortic aneurysm); Arachnodactyly, joint hypermobility; Hyperextensible skin; Tall stature; Berry aneurysms; High-arched palate; Spinal deformities (scoliosis, hyperkyphosis); Long extremities; Pes planus, hindfoot valgus.
E	Sterile platelet thrombi formation	
F	Viridans group streptococci infection	

Hint

The phonocardiogram shows a crescendo-decrescendo murmur during systole. Such a murmur at the aortic area (2nd intercostal space) is characteristic of aortic stenosis, which can manifest with angina, dyspnea, and/or lightheadedness.

Correct Answer

A - M protein cross-reactivity

Explanation Why

M protein [cross-reactivity](#) is the underlying cause of [rheumatic heart disease](#) (RHD). In RHD, [antibodies](#) to [streptococcal M protein](#) cross-react with valvular [proteins](#), resulting in valve [inflammation](#), [fibrosis](#), and stenosis. While RHD is the most common cause of AS in developing countries, it is relatively uncommon in the United States.


B - Metastatic valvular calcification

Explanation Why

[Metastatic calcification](#) of the valves occurs in previously healthy tissue due to chronic [hypercalcemia](#) or [hyperphosphatemia](#) (as opposed to [dystrophic calcification](#)). [Aortic stenosis](#) caused by [metastatic calcification](#) would manifest with [dyspnea](#) and [chest pain](#). However, this patient has no underlying condition that would predispose to [metastatic calcification](#), which includes [primary hyperparathyroidism](#), [multiple myeloma](#), [sarcoidosis](#), and [chronic kidney disease](#).

C - Congenital leaflet fusion

Image




Explanation Why

Congenital [aortic valve](#) leaflet fusion is the most likely cause of [aortic stenosis](#) (AS) in this patient. AS is the most common [valvular heart disease](#) in industrialized countries, with [incidence](#) increasing with age. In patients < 70 years presenting with AS, a congenital [bicuspid aortic valve](#) abnormality is most likely responsible. This pathology predisposes the valve to [dystrophic calcification](#) and degeneration that occurs earlier than calcific [aortic stenosis](#) associated with old age.

D - Cystic medial necrosis

Image



Explanation Why

Cystic medial necrosis is the pathology underlying aneurysmal dilation of the thoracic aorta and the aortic root, which is associated with hypertension, bicuspid aortic valve, and connective tissue disorders such as Marfan syndrome. If the aortic root is sufficiently dilated, these patients can present with aortic regurgitation and symptoms such as dyspnea and chest pain. However, cystic medial necrosis is not associated with aortic stenosis.

E - Sterile platelet thrombi formation

Explanation Why

Sterile platelet thrombi can form on cardiac valves in patients with nonbacterial thrombotic endocarditis (NBTE), which is usually associated with rheumatologic disorders (e.g., systemic lupus

[erythematosus](#)) or malignancies. While patients with [NBTE](#) can present with [dyspnea](#) and [chest pain](#), there is no indication of an underlying pathology in this patient. Furthermore, [NBTE](#) is associated with aortic or [mitral regurgitation](#), not [aortic stenosis](#).


F - Viridans group streptococci infection

Explanation Why

[Viridans group streptococci](#) are associated with [subacute bacterial endocarditis](#), which can cause regurgitant valvular lesions, e.g., mitral or [aortic regurgitation](#), manifesting with [dyspnea](#). However, [infectious endocarditis](#) does not typically cause [aortic stenosis](#). Furthermore, the patient here has no systemic symptoms that could be attributed to [subacute bacterial endocarditis](#), e.g., [fever](#), malaise, and/or [tachycardia](#).

Question # 22

A 76-year-old woman with a history of hypertension and type 2 diabetes mellitus is brought to the emergency department 60 minutes after the acute onset of left-sided abdominal pain and nausea with vomiting. Three weeks ago, she underwent emergency surgical revascularization for acute left lower extremity ischemia. Physical examination shows left upper quadrant tenderness without rebound or guarding. Serum studies show an elevated lactate dehydrogenase level. Laboratory studies, including a complete blood count, basic metabolic panel, and hepatic panel, are otherwise unremarkable. A transverse section of a CT scan of the abdomen is shown. Further evaluation is most likely to show which of the following?



	Answer	Image
A	Absent P waves on electrocardiogram	
B	Non-compressible femoral vein on ultrasonography	

	Answer	Image
C	Infrarenal aortic aneurysm on abdominal CT scan	
D	Right atrial thrombus on transesophageal echocardiography	
E	Schistocytes on peripheral blood smear	

Hint

This woman's CT findings (wedge-shaped splenic hypodensity) and acute onset of left-sided abdominal pain are consistent with acute splenic infarction. This woman has had two different thromboembolic events in the arterial system.

Correct Answer

A - Absent P waves on electrocardiogram

Explanation Why

Absent [P waves](#) and irregularly irregular RR intervals on [electrocardiogram](#) would be consistent with [atrial fibrillation](#), a significant [risk factor](#) for atrial thrombus formation due to the rapid, irregular contraction of the [atria](#) and subsequent blood stasis. [Thrombi](#) formed in the [left atrium](#) or [left ventricle](#) can fragment into emboli and disperse into the arterial circulation, leading to [ischemic](#) events, as seen here. This patient has multiple [risk factors for atrial fibrillation](#), including her age, [hypertension](#), and [diabetes](#).

B - Non-compressible femoral vein on ultrasonography

Explanation Why

[Sonographic](#) evidence of a non-compressible [femoral vein](#) would be consistent with a [deep vein thrombosis \(DVT\)](#) in the [femoral vein](#). While the patient's [acute limb ischemia](#) and splenic [infarction](#) were probably caused by embolic events, the emboli both occurred in the arterial system, whereas [DVTs](#) result in occlusion of the venous system. [Venous emboli](#) from [DVTs](#) can travel to the pulmonary vasculature (resulting in [pulmonary embolism](#)) but would not enter the systemic vasculature without the presence of a significant [right-to-left shunt](#) (e.g., [patent foramen ovale](#)), which is uncommon.

C - Infrarenal aortic aneurysm on abdominal CT scan

Explanation Why

Infrarenal [aortic aneurysms](#) can increase the risk of arterial embolic events at sites [distal](#) to the [aneurysm](#), such as the lower extremities. The [splenic artery](#) and short gastric vessels supplying the [spleen](#) arise from the [celiac trunk](#), which originates above the [renal arteries](#). As a result, the [spleen](#) would not be affected by an infrarenal [aortic aneurysm](#).

D - Right atrial thrombus on transesophageal echocardiography

Explanation Why

A right atrial [thrombus](#) could embolize into the pulmonary vasculature, causing [pulmonary embolism](#). However, embolism from a right atrial [thrombus](#) into the arterial system would only be possible with a concurrent intra-cardiac shunt (e.g., [patent foramen ovale](#)). Given this patient's age and [risk factors](#), a different finding is much more likely to be present.

E - Schistocytes on peripheral blood smear

Explanation Why

[Schistocytes](#) may be present on a [peripheral blood smear](#) as a result of [intravascular hemolysis](#) in conditions such as [hemolytic uremic syndrome \(HUS\)](#) and [thrombotic thrombocytopenic purpura \(TTP\)](#). Both [HUS](#) and [TTP](#) can elevate [LDH](#) levels and cause splenic [infarcts](#). However, in the absence of further signs of [intravascular hemolysis](#) (e.g., [anemia](#), [hyperbilirubinemia](#)), [HUS](#) and [TTP](#) are unlikely causes of the patient's splenic [infarction](#). This patient's elevated [LDH](#) level is most likely a direct result of her splenic [infarct](#).

Question # 23

A 53-year-old woman with hypertension and hyperlipidemia comes to the physician because of a 3-month history of progressively worsening shortness of breath. Her pulse is 92/min, respirations are 22/min, and blood pressure is 110/70 mm Hg. Cardiac examination shows a grade 3/6 holosystolic murmur at the apex that radiates to the axilla and an extra heart sound during early diastole. Assuming all other parameters remain constant, which of the following is most likely to decrease the volume of retrograde blood flow in this patient?

	Answer	Image
A	Increase in atrioventricular orifice size	
B	Decrease in left atrial systolic pressure	
C	Increase in left ventricular inotropy	
D	Decrease in systemic vascular resistance	
E	Increase in left ventricular end-diastolic pressure	

Hint

A holosystolic murmur that is best heard at the cardiac apex and radiates to the axilla is characteristic of mitral regurgitation (MR). The extra heart sound during diastole is most likely an S₃ gallop, which is often heard in patients with moderate to severe MR.

Correct Answer

A - Increase in atrioventricular orifice size

Explanation Why

The left atrioventricular orifice is formed by the [mitral valve](#) annulus. In the case of [mitral regurgitation](#), an increase in the size of this annulus would increase rather than decrease retrograde blood flow (regurgitation).

B - Decrease in left atrial systolic pressure

Explanation Why

The left atrial [systolic](#) pressure acts as a resistance to retrograde blood flow across the [mitral valve](#). Therefore, a decrease in left atrial [systolic](#) pressure would increase rather than decrease the volume of regurgitated blood.

C - Increase in left ventricular inotropy

Explanation But

[Inotropic](#) drugs are used in patients with [cardiogenic shock](#) due to acute [mitral regurgitation](#) but this is done in order to improve the [cardiac output](#), not to decrease retrograde blood flow.

Explanation Why

Increased [left ventricular inotropy](#) would increase the [ejection fraction](#) and thereby decrease the [left ventricular end diastolic volume](#). In a patient with [mitral valve](#) incompetence, as seen here, this phenomenon would be expected to decrease the regurgitant fraction (retrograde blood flow) across the [mitral valve](#) in the next [cardiac cycle](#). However, this effect of increased LV [inotropy](#) is counteracted by the increase in [left ventricular systolic](#) pressure, which tends to increase regurgitant blood flow. Overall, increased LV [inotropy](#) does not result in a specific or predictable change in the

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

regurgitant volume, as has been shown empirically in studies with [dobutamine](#) (an [inotropic](#) drug).

D - Decrease in systemic vascular resistance

Explanation Why

A decrease in [systemic vascular resistance](#) (SVR) would reduce the volume of retrograde blood flow through the [mitral valve](#). In [mitral regurgitation](#) (MR), the left ventricular ejection fraction is divided between the output through the [aortic valve](#) and through the abnormal regurgitant [mitral valve](#). When SVR ([left ventricular afterload](#)) is decreased, a greater volume of blood is ejected through the [aortic valve](#) and less blood flows in a retrograde direction across the [mitral valve](#) (lower regurgitant fraction). This is the rationale for using systemic [vasodilators](#) (e.g., [sodium nitroprusside](#)) to treat [dyspnea](#) in patients with severe acute MR. Conversely, maneuvers such as isometric contraction and a hand grip, which increase SVR, reduce [ejection fraction](#) through the [aortic valve](#) and increase the retrograde flow.


E - Increase in left ventricular end-diastolic pressure

Explanation Why

As retrograde flow is dependent on the pressure gradient between the LV and [left atrium](#), an increase in left ventricular end-diastolic pressure (LVEDP) would increase the driving pressure and increase retrograde blood flow.

Question # 24

A group of epidemiologists is studying the rates of cardiac surgeries performed in several Southeast Asian countries compared to the United States. Results show a significant increase in the number of mitral valve replacements performed in Vietnam in adults aged 30–50 years compared to the same age group in the United States. Which of the following public health interventions is most likely to decrease this number?

	Answer	Image
A	Prompt antibiotic treatment for bacterial pharyngitis	
B	Improved access to drug rehabilitation centers	
C	Genetic screening program for FBN1 gene mutation	 Marfan syndrome Epidemiology Prevalence: 1-5/10,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15; autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissection) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups) Clinical features: Ectopia lentis, severe myopia Heart valve defects Pectus deformity (pectus carinatum or pectus excavatum) Aortic disease (e.g., aortic dissection, aortic aneurysm) Anachondactyly, joint hypermobility Hyperextensible skin Tall stature Berry aneurysms High-arched palate Spinal deformities (scoliosis, hyperkyphosis) Long extremities Pes planus, hindfoot valgus
D	Childhood vaccination against <i>Corynebacterium diphtheriae</i>	
E	Screening echocardiography for prolapsed anterior mitral valve leaflets	

Hint

The underlying cardiac illness in these patients is the most common cause of cardiovascular disease in children and adolescents in developing countries.

Correct Answer

A - Prompt antibiotic treatment for bacterial pharyngitis

Explanation Why

Prompt [antibiotic](#) treatment against established cases of group A [streptococcal pharyngitis](#) prevents the development of [acute rheumatic fever](#) and associated [endocarditis](#). [Acute rheumatic fever](#) is the most common cause of [mitral stenosis](#) (MS). Severe cases of MS may require surgical valve replacement. Therefore, early [antibiotic](#) treatment of [bacterial pharyngitis](#) indirectly decreases the number of [mitral valve](#) replacements. Group A [streptococcal](#) tonsillopharyngitis is diagnosed either by throat culture or rapid strep test and is treated with [penicillin V](#).


B - Improved access to drug rehabilitation centers

Explanation Why

Improved access to drug rehabilitation centers is associated with a decreased [incidence](#) of recreational IV drug use, resulting in decreased septicemia and [endocarditis](#). However, IV drug use-related [endocarditis](#) is primarily associated with [S. aureus](#), *Pseudomonas*, and [Candida](#) infections of the [tricuspid valve](#), not the [mitral valve](#).

C - Genetic screening program for FBN1 gene mutation

Image



Explanation Why

Genetic screening programs for [FBN1 gene mutations](#) allows for early detection of [Marfan syndrome](#). [Marfan syndrome](#) predisposes to the development of [mitral valve prolapse](#), which is susceptible to progress into [mitral regurgitation](#). Severe [mitral regurgitation](#) is treated with [mitral valve](#) replacement. However, early detection of the syndrome neither alters the disease progression nor decreases the need for eventual [mitral valve](#) replacement.

D - Childhood vaccination against *Corynebacterium diphtheriae*

Explanation Why

Childhood [vaccinations](#) against [Corynebacterium diphtheriae](#) prevent the absorption and dissemination of [diphtheria toxin](#) throughout the body. The toxin is associated with cardiac tissue

damage resulting in cardiac dilation and [arrhythmias](#). However, [diphtheria toxin](#) is associated with [myocarditis](#) more than [endocarditis](#) (and subsequent valve destruction). Therefore, treatment of the disease would not significantly reduce the number of [mitral valve](#) replacements compared to other measures.



E - Screening echocardiography for prolapsed anterior mitral valve leaflets

Explanation Why

[Echocardiographic](#) screening for prolapsed [anterior mitral valve](#) leaflets allows for early detection of [mitral valve prolapse \(MVP\)](#) that occurs sporadically (primary [MVP](#)) or secondary to [connective tissue disorders](#). [MVP](#) predisposes mitral valves to the development of [mitral regurgitation](#) and severe [mitral regurgitation](#) is treated with [mitral valve](#) replacement. However, early detection of [MVP](#) does not alter the disease course and would not decrease the number of [mitral valve](#) replacements.

Question # 25

A 72-year-old man comes to the physician because of a 6-month history of intermittent dull abdominal pain that radiates to the back. He has smoked one pack of cigarettes daily for 50 years. His blood pressure is 145/80 mm Hg. Abdominal examination shows generalized tenderness and a pulsatile mass in the periumbilical region on deep palpation. Further evaluation of the affected blood vessel is most likely to show which of the following?

	Answer	Image
A	Accumulation of foam cells in the tunica intima	
B	Obliterative inflammation of the vasa vasorum	
C	Necrotizing inflammation of the entire vessel wall	

	Answer	Image
D	Formation of giant cells in the tunica media	Polymyalgia rheumatica (PMR) Giant cell arteritis (GCA; temporal arteritis) Etiology Autoimmune; granulomatous inflammation of medium and large arteries (GCA esp. affects the carotid arteries). A common pathophysiology is presumed. Epidemiology ♀ > ♂; incidence rises with age Antibody serology No autoantibodies Laboratory analysis ↑↑ ESR, ↑ CRP Complications GCA: 20–30% risk of blindness without therapy Note Distal joints are not involved! GCA: 50% associated with PMR General symptoms 👉 B symptoms 👉 Depressive mood 👉 Weight loss
E	Fragmentation of elastic tissue in the tunica media	Marfan syndrome Epidemiology Prevalence: 1–5/10,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15; autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissections) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups)


Hint

The presence of chronic dull abdominal pain and a pulsatile abdominal mass suggests an abdominal aortic aneurysm (AAA).

Correct Answer

A - Accumulation of foam cells in the tunica intima

Image



Explanation Why

Accumulation of [foam cells](#) in the [tunica intima](#) is the pathological process underlying [atheroma](#) formation ([atherosclerosis](#)), which is the most common cause of aortic wall weakening and formation of an [AAA](#). This patient's history of smoking, [hypertension](#), and advanced age are all [risk factors for atherosclerosis](#).

B - Obliterative inflammation of the vasa vasorum


Explanation Why

Obliterative [inflammation](#) of the [vasa vasorum](#) is characteristic of syphilitic [aortitis](#), which can result

in an [aortic aneurysm](#). However, syphilitic [aortitis](#) affects the [ascending aorta](#), whereas this patient presents with a mass in the abdominal portion of the [descending aorta](#).

C - Necrotizing inflammation of the entire vessel wall

Image




Explanation Why

Necrotizing [inflammation](#) of all the layers of the vessel wall is characteristic of [polyarteritis nodosa](#) ([PAN](#)), which can cause microaneurysms in medium-sized [arteries](#) (e.g., [mesenteric arteries](#) and [renal arteries](#)), episodic abdominal [pain](#) (due to [mesenteric ischemia](#)), and [hypertension](#) (due to renal involvement). However, [PAN](#) would not cause a pulsatile periumbilical mass because it spares large [arteries](#) like the aorta. Other clinical features of [vasculitis](#), such as [fever](#), malaise, weight loss, and cutaneous lesions (e.g., [livedo reticularis](#), [distal](#) ulcerations), would be expected instead.

D - Formation of giant cells in the tunica media

Image




Explanation Why

Formation of [giant cells](#) in the [tunica media](#) is characteristic of [giant cell arteritis \(GCA\)](#) and [Takayasu arteritis](#), both of which are [large vessel vasculitides](#) and can result in an [AAA](#). However, neither condition is the most common cause of [AAA](#), and this patient does not report any symptoms that would otherwise suggest an increased likelihood of [GCA](#) (e.g., [headache](#), scalp tenderness, or [jaw claudication](#)) or [Takayasu arteritis](#) (e.g., limb [claudication](#) or onset of symptoms before the age of 40 years), so a different pathological process is more likely.

E - Fragmentation of elastic tissue in the tunica media

Image



Explanation Why

Fragmentation of elastic tissue in the [tunica media](#) is the pathological process underlying [cystic medial necrosis](#) (CMN), which occurs in certain [connective tissue disorders](#) such as [Marfan syndrome](#). However, for reasons that are not completely understood, CMN typically affects the [thoracic aorta](#) rather than the [abdominal aorta](#). A different pathological process is the most common cause of an [AAA](#). Furthermore, this patient does not have features of [Marfan syndrome](#) (e.g., tall stature, long extremities, [joint](#) hypermobility, [ectopia lentis](#)).

Question # 26

A 30-year-old man with Down syndrome is brought to the physician by his mother for the evaluation of fatigue. Physical examination shows bluish-colored lips and digital clubbing that were not present at his most recent examination. Right heart catheterization shows a right atrial pressure of 32 mmHg. Which of the following is most likely involved in the pathogenesis of this patient's current condition?

	Answer	Image
A	Aortic valve regurgitation	
B	Reversible pulmonary hypertension	
C	Intrapulmonary shunting	
D	Right ventricular hypertrophy	
E	Asymmetric septal hypertrophy	

Hint

Congenital heart defects associated with Down syndrome include atrioventricular, ventricular, and atrial septal defects.

Correct Answer

A - Aortic valve regurgitation

Explanation Why

[Aortic valve regurgitation](#) is most commonly seen in [connective tissue disorders](#) such as [Marfan syndrome](#). Backflow of blood from the aorta into the [left ventricle](#) would result in [left ventricular hypertrophy](#) and increased left atrial pressure rather than increased right atrial pressure. In addition, [aortic valve regurgitation](#) presents with fatigue but not with [clubbing](#) and [cyanosis](#). This patient's history of [Down syndrome](#) makes a different mechanism more likely.

B - Reversible pulmonary hypertension

Explanation Why

[Pulmonary hypertension](#) is a manifestation of [acyanotic heart defects](#), which are the result of a left-to-right shunt. The increased volume of the shunted blood causes a remodeling of the pulmonary vasculature, leading to [pulmonary hypertension](#). This patient most likely has [pulmonary hypertension](#) but this is an irreversible process, not reversible.

C - Intrapulmonary shunting

Explanation Why

Intrapulmonary shunting occurs in situations where parts of the [lungs](#) are perfused but not ventilated, resulting in a [ventilation-perfusion mismatch](#). Some common conditions that can cause intrapulmonary shunt include pulmonary [AVM](#), [hepatopulmonary syndrome](#), severe cases of [pulmonary edema](#), and lobar [atelectasis](#). This patient's history of [Down syndrome](#) makes an underlying congenital [heart](#) defect a more likely cause of his condition, rather than a pulmonary condition.

D - Right ventricular hypertrophy

Explanation Why

Central [cyanosis](#), weakness, and [clubbing](#) in a patient with a history of [Down syndrome](#) are highly suggestive of [Eisenmenger syndrome](#) due to a congenital [heart](#) defect, most likely an endocardial or [atrioventricular cushion](#) defect. Both of these defects initially cause a left-to-right blood shunt (i.e., [acyanotic heart defect](#)) due to higher pressures within the left [heart](#). Over time, the increased blood flow within the right [heart](#) and the pulmonary vasculature results in permanent remodeling of pulmonary vessels and [right ventricular hypertrophy](#). The [right ventricular](#) pressure eventually becomes higher (normal 5–10 mmHg) than the [left ventricular](#) pressure, causing a reversal of the shunt and [cyanosis](#).

E - Asymmetric septal hypertrophy

Explanation Why

Asymmetric septal [hypertrophy](#) is seen in [hypertrophic obstructive cardiomyopathy \(HOCM\)](#), a hereditary condition that can manifest with fatigue and an increased atrial pressure. However, [HOCM](#) would not manifest with central [cyanosis](#), [digital clubbing](#), and increased right atrial pressure (the left side is more commonly involved). This patient's history of [Down syndrome](#) makes a different etiology more likely.

Question # 27

A 73-year-old man with coronary artery disease and hypertension is brought to the emergency department by ambulance 90 minutes after the acute onset of substernal chest pain and dyspnea. He has smoked 2 packs of cigarettes daily for 52 years. Shortly after arriving at the hospital, he loses consciousness and is pulseless. Despite attempts at cardiopulmonary resuscitation, he dies. Examination of the heart at autopsy shows complete occlusion of the left anterior descending artery with a red thrombus overlying a necrotic plaque. Which of the following pathophysiologic mechanisms is most likely responsible for this patient's coronary occlusion?

	Answer	Image
A	Stasis of blood flow	
B	Type III collagen deposition	
C	Influx of lipids into the endothelium	
D	Proliferation of smooth muscle cells	

	Answer	Image
E	Secretion of matrix metalloproteinases	The diagram illustrates the progression of atherosclerosis in a blood vessel wall over time, categorized into Asymptomatic, Decades, Symptomatic, and Minutes. The vessel wall layers shown are the endothelium, intima, and media. Asymptomatic: Normal endothelium: The vessel wall is healthy. Endothelial dysfunction: Caused by chronic stress (e.g., due to hypertension, inflammation). Decades: Infiltration of monocytes and LDL resulting in accumulation of macrophages and oxLDL: LDL enters the intima, is oxidized to oxLDL, and attracts monocytes which become macrophages. Foam cell formation resulting in fatty streaks: Macrophages ingest oxLDL, becoming foam cells. Proliferation and migration of SMCs and deposition of ECM: Smooth muscle cells (SMCs) migrate from the media to the intima and proliferate, depositing extracellular matrix (ECM). Symptomatic: Fibrous plaque formation: The plaque becomes more complex with a fibrous cap. Plaque destabilization: The fibrous cap weakens. Plaque rupture (e.g., in context of increased matrix degradation): The cap ruptures, releasing contents into the bloodstream. Thrombus formation: Platelets aggregate to form a thrombus on the ruptured plaque. Legend: - LDL (green circle) - Monocyte (purple circle) - Macrophage (yellow star) - oxLDL (green star) - Foam cell (orange cell) - Platelets (red dots) - Lymphocytes (blue circle) - Extracellular matrix (ECM) (orange mesh) - Smooth muscle cell (SMC) (pink cell) - Cholesterol crystals (blue crystals) - Cellular debris (brown dots) - Fibrous cap (pink layer) - Necrotic core (yellow area) - SMC apoptosis, breakdown of the stabilizing ECM (pink cell with arrows pointing away)
F	Release of platelet-derived growth factor	
G	Migration of macrophages to the tunica intima	

Hint

The cause of death in this patient is an acute myocardial infarction due to acute plaque rupture and thrombosis, as evidenced by the autopsy finding of coronary occlusion by a red thrombus (composed of platelets and fibrin). Plaque rupture occurs when the protective cap covering the plaque loses its integrity and thrombotic material causes vessel obstruction.

Correct Answer

A - Stasis of blood flow

Explanation Why

Stasis of blood flow is part of the tripartite pathophysiology of [deep venous thrombosis](#). The other two contributing factors are [endothelial](#) damage and relative [hypercoagulability](#) ([Virchow triad](#)). However, the hemodynamics of the [coronary arteries](#) is characterized by high-speed flow, even in the case of mural thrombus. [Endothelial](#) damage and relative [hypercoagulability](#) play a more prominent role than stasis in the pathophysiology of [atheromas](#) and this patient's [myocardial infarction](#).

B - Type III collagen deposition

Explanation Why

Deposition of [type III collagen](#) by [fibroblasts](#) increases the thickness of the fibrous cap and stabilizes the [atheroma](#). This is seen at the site of previous [atheroma](#) after plaque rupture in patients who survive [myocardial infarctions](#). A different process is responsible for acute [atheroma](#) rupture.

C - Influx of lipids into the endothelium

Explanation Why

The influx of lipids into the [endothelium](#) is responsible for [atheroma](#) formation. Because lipids are not normally found in the [endothelium](#), their presence sets off a potent [immune response](#), which is what ultimately causes the nascent [atheroma](#) to develop and enlarge. However, the presence of lipids does not play a role in the acute rupture of the [atheroma](#).


D - Proliferation of smooth muscle cells

Explanation Why

The [proliferation](#) of [smooth muscle cells](#) is responsible for enlargement of the [atheroma](#). [Smooth muscle hyperplasia](#) is thought to occur secondary to signaling via endothelin and [platelet-derived growth factor](#). However, this process does not contribute to the acute rupture of the [atheroma](#).

E - Secretion of matrix metalloproteinases

Image



Explanation Why

The secretion of [matrix metalloproteinases](#) is thought to be responsible for breaking down the fibrous cap of an [atheroma](#). [Matrix metalloproteinases](#), which are secreted by inflammatory cells in the [atheroma](#) (e.g., [macrophages](#)), weaken the fibrous cap by breaking down the [extracellular matrix](#). Subsequently, minor stress can rupture the fibrous cap and expose the [atheroma](#)'s highly

thrombogenic lipid core. It is this event that then causes the acute, total obstruction of the vessel and subsequent [myocardial infarction](#).

F - Release of platelet-derived growth factor

Explanation Why

[Platelet-derived growth factor](#) induces [smooth muscle hyperplasia](#) and [fibroblast](#) growth in the [atheroma](#). Although this is considered a crucial pathophysiologic mechanism in the growth of an [atheroma](#), [platelet-derived growth factor](#) plays no direct role in acute plaque rupture.

G - Migration of macrophages to the tunica intima

Explanation Why

Migration of [macrophages](#) to the [tunica intima](#) is responsible for the formation of the [atheroma](#), but it is not responsible for rupture of the [atheroma](#).

Question # 28

A 19-year-old woman comes to the physician for a routine health maintenance examination. She appears well. Her vital signs are within normal limits. Cardiac auscultation shows a mid-systolic click and a grade 3/6, late-systolic, crescendo murmur that is best heard at the cardiac apex in the left lateral recumbent position. After the patient stands up suddenly, the click is heard during early systole and the intensity of the murmur increases. Which of the following is the most likely underlying cause of this patient's examination findings?

	Answer	Image
A	Myosin heavy chain defect	
B	Aortic root dilation	The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The ECG shows a normal sinus rhythm. The PCG shows a normal S1 and S2. The S2 is split into S1 and S2, with the S2 being significantly louder than the S1. The PCG shows a normal S1 and S2. The S2 is split into S1 and S2, with the S2 being significantly louder than the S1. The PCG shows a normal S1 and S2. The S2 is split into S1 and S2, with the S2 being significantly louder than the S1.
C	Dermatan sulfate deposition	The image displays two cardiac waveforms: an Electrocardiogram (ECG) in blue and a Phonocardiogram (PCG) in orange. The ECG shows a normal sinus rhythm. The PCG shows a normal S1 and S2. The S2 is split into S1 and S2, with the S2 being significantly louder than the S1. The PCG shows a normal S1 and S2. The S2 is split into S1 and S2, with the S2 being significantly louder than the S1. The PCG shows a normal S1 and S2. The S2 is split into S1 and S2, with the S2 being significantly louder than the S1.

	Answer	Image
D	Chordae tendinae thickening	The image displays two cardiac monitoring traces. The top trace is an Electrocardiogram (ECG) in blue, showing a regular rhythm with alternating shaded regions for Diastole (light blue) and Systole (light grey). The bottom trace is a Phonocardiogram (PCG) in orange, showing heart sounds. The PCG is aligned with the ECG, with S1 (first heart sound) occurring at the start of the Diastole and S2 (second heart sound) at the start of the Systole. An arrow labeled 'OS' points to the S2 sound, which is notably soft and muffled, a characteristic finding in chordae tendinae thickening.
E	Congenital valvular fusion	
F	Congenital interventricular communication	
G	Dystrophic valvular calcification	The image displays two cardiac monitoring traces. The top trace is an Electrocardiogram (ECG) in blue, showing a regular rhythm with alternating shaded regions for Diastole (light blue) and Systole (light grey). The bottom trace is a Phonocardiogram (PCG) in orange, showing heart sounds. The PCG is aligned with the ECG, with S1 (first heart sound) occurring at the start of the Diastole and S2 (second heart sound) at the start of the Systole. An arrow labeled 'OS' points to the S2 sound, which is notably soft and muffled, a characteristic finding in dystrophic valvular calcification.

Hint

The presence of a mid-systolic click followed by a crescendo murmur at the mitral area (cardiac apex) is suggestive of mitral valve prolapse (MVP). Maneuvers that decrease the left ventricular size (e.g., sudden standing) increase the intensity of the murmur and cause the click to occur earlier in systole.

Correct Answer


A - Myosin heavy chain defect

Explanation Why

A [myosin heavy chain](#) defect is associated with familial [hypertrophic cardiomyopathy](#), which is frequently asymptomatic and classically manifests with a [heart murmur](#) that can be intensified by standing up. However, [HOCM](#) usually presents with a [mitral regurgitation systolic murmur](#) and/or a [systolic ejection murmur](#) (crescendo-decrescendo), but not with a mid-systolic click and crescendo [systolic murmur](#).

B - Aortic root dilation

Image




Explanation Why

[Aortic root dilation](#) is a characteristic feature of [Marfan syndrome](#) and can lead to [aortic regurgitation](#). However, AR would manifest with a [diastolic murmur](#) with a decrescendo best heard at the left sternal border, not with a mid-systolic click and crescendo [systolic murmur](#) heard in the [mitral area](#). Moreover, the [murmur](#) in AR intensifies when the patient leans forward and decreases instead of increases by standing.

C - Dermatan sulfate deposition

Image



Explanation But

Maneuvers that increase [afterload](#) (e.g., handgrip, squatting) are another way to help differentiate [MVP](#) from other causes of mitral insufficiency. In [MVP](#), these maneuvers cause the click to move later in [systole](#) and become softer due to an increase in [left ventricular](#) size.


Explanation Why

[Dermatan sulfate](#) deposition in the [mitral valve](#) leaflets, which is characteristic of myxomatous

degeneration of the [mitral valve](#), is the most likely underlying cause of this patient's [mitral valve prolapse](#). MVP is most commonly idiopathic but can also be associated with [connective tissue disorders](#) (e.g., [Marfan syndrome](#), [Ehlers-Danlos syndrome](#), [fragile X](#) syndrome) or occur as a complication of [infective endocarditis](#), [rheumatic heart disease](#), and [myocardial infarction](#). The mid-[systolic](#) click heard on auscultation is caused by sudden tensing of the chordae tendinae, which occurs earlier and more forcefully with decreased [left ventricular](#) size.

D - Chordae tendinae thickening

Image



Explanation Why

Chordae tendinae thickening can be seen in [mitral valve stenosis](#) due to large vegetations, [rheumatic fever](#), or autoimmune inflammatory processes (e.g., [SLE](#)). However, [mitral stenosis](#) would present with a delayed [diastolic murmur](#) with a decrescendo, not with a mid-[systolic](#) click and crescendo [systolic murmur](#). MVP, which this patient likely has based on her [murmur](#), is associated with thinning rather than thickening of the chordae tendinae.

E - Congenital valvular fusion

Explanation Why

Congenital valvular fusion is the underlying cause of [bicuspid aortic valve](#), which is usually asymptomatic but can predispose to both [aortic stenosis](#) (AS) and [aortic regurgitation](#). A calcified [bicuspid aortic valve](#) manifests with an aortic [systolic ejection click](#) followed by a late [systolic](#) crescendo-decrescendo [murmur](#) best heard over the 2nd right [intercostal space](#), not with a mid-[systolic](#) click and crescendo [systolic murmur](#) heard best in the [mitral area](#). Furthermore, the [murmur](#) in AS decreases instead of increases by sudden standing. [Aortic regurgitation](#), in turn, presents with a [murmur](#) during [diastole](#) (not [systole](#)).

F - Congenital interventricular communication

Explanation Why

Congenital interventricular communication causes a [ventricular septal defect](#), which is usually asymptomatic (if the defect is small). However, [VSD](#) manifests with a harsh [holosystolic murmur](#) over the left lower sternal border and a mid-[diastolic murmur](#) over the cardiac apex, not with a mid-[systolic](#) click and crescendo [systolic murmur](#) heard in the [mitral area](#). In addition, the intensity of the [murmur](#) in [VSD](#) would be decreased or remain unchanged by standing up instead of increasing.

G - Dystrophic valvular calcification

Image



Explanation Why

[Dystrophic](#) valvular calcification is associated with calcific [aortic stenosis](#), which can be initially asymptomatic. However, [aortic stenosis](#) would manifest with a harsh crescendo-decrescendo [systolic ejection murmur](#) parasternal over the 2nd right [intercostal space](#), not with a mid-[systolic](#) click and crescendo [systolic murmur](#) heard in the [mitral area](#). Also, the [murmur](#) in [aortic stenosis](#) is best heard when the patient leans forward and would be decreased instead of increased by standing. Moreover, patients with [aortic stenosis](#) are usually much older (≥ 75 years).

Question # 29

An otherwise healthy 65-year-old man comes to the physician for a follow-up visit for elevated blood pressure. Three weeks ago, his blood pressure was 160/80 mmHg. Subsequent home blood pressure measurements at days 5, 10, and 15 found: 165/75 mm Hg, 162/82 mm Hg, and 170/80 mmHg, respectively. He had a cold that was treated with over-the-counter medication 4 weeks ago. Pulse is 72/min and blood pressure is 165/79 mm Hg. Physical examination shows no abnormalities. Laboratory studies, including thyroid function studies, serum electrolytes, and serum creatinine, are within normal limits. Which of the following is the most likely underlying cause of this patient's elevated blood pressure?

	Answer	Image
A	Reduced renal artery blood flow	
B	Decrease in arterial compliance	
C	Increase in aldosterone production	
D	Increase in left ventricular end-diastolic volume	
E	Decrease in baroreceptor sensitivity	
F	Medication-induced vasoconstriction	

Hint

This elderly patient has a wide pulse pressure with elevated systolic blood pressure readings and normal diastolic pressures.

Correct Answer

A - Reduced renal artery blood flow

Explanation Why

A decrease in [renal artery](#) blood flow would activate the [renin-angiotensin system](#) and result in [renovascular hypertension](#). Important causes of reduced [renal artery](#) blood flow include [renal artery stenosis](#), external compression of the [renal arteries](#) (e.g., by a [tumor](#) or cyst), [atherosclerosis](#) of the [renal arteries](#), and [fibromuscular dysplasia](#). While [atherosclerosis](#) of the [renal arteries](#) should be considered given the patient's age, his normal [diastolic](#) pressures make another underlying cause more likely. Furthermore, only bilateral [renal artery](#) disease resulting in a significant decrease in [GFR](#) would manifest with [hypertension](#); this patient's normal [creatinine](#) makes such a decrease unlikely.

B - Decrease in arterial compliance

Explanation Why

A decrease in arterial compliance secondary to age-related reduction in arterial elasticity and increased stiffness is common in individuals > 60 years old. This decrease results in [isolated systolic hypertension](#), which is characterized by a wide [pulse pressure](#), as seen in this patient. Treatment with an anti-hypertensive agent, such as a [thiazide diuretic](#) or [calcium channel blocker](#), should be initiated to decrease the risk of a cardiovascular event (e.g., [MI](#), [stroke](#), or renal dysfunction).

C - Increase in aldosterone production

Explanation Why

[Primary hyperaldosteronism](#) is the most common cause of [secondary hypertension](#) and classically manifests with [resistant hypertension](#) and [hypokalemia](#). The blood pressure seen in [primary hyperaldosteronism](#) is often markedly elevated and causes increases in both [systolic](#) and [diastolic](#) pressures. This patient's [isolated systolic hypertension](#) and normal lab values make this diagnosis unlikely.

D - Increase in left ventricular end-diastolic volume

Explanation Why

Elevated LV end-diastolic volume, as seen in [aortic regurgitation](#) (AR), increases [stroke volume](#) and can also widen [pulse pressure](#). However, AR alone would not cause this patient's [hypertension](#), and a high-pitched, early [diastolic murmur](#) would be expected.

E - Decrease in baroreceptor sensitivity

Explanation Why

A decrease in [baroreceptor](#) sensitivity is the underlying mechanism of [orthostatic hypotension](#). Unlike this patient's asymptomatic [hypertension](#), it is characterized by [autonomic dysfunction](#) and insufficient counter-regulation after postural changes that leads to [dizziness](#) and [syncope](#) after standing up.

F - Medication-induced vasoconstriction

Explanation Why

Most over-the-counter nasal decongestants contain [sympathomimetics](#) (e.g., [phenylephrine](#), [pseudoephedrine](#)), which induce both local and systemic vasoconstriction. While these medications can cause [hypertension](#), the time course of this patient's presentation and his isolated [systolic](#) pressures make a medication-induced etiology unlikely.

Question # 30

A 2-year-old girl is brought to the physician by her mother for a well-child examination. Cardiac auscultation is shown. When she clenches her fist forcefully for a sustained time, the intensity of the murmur increases. Which of the following is the most likely cause of this patient's auscultation findings?

[Video](#)

	Answer	Image
A	Prolapse of the mitral valve	
B	Fusion of the right and left coronary leaflets	
C	Defect in the atrial septum	
D	Asymmetric hypertrophy of the left ventricle	
E	Defect in the ventricular septum	
F	Failure of the ductus arteriosus to close	

Hint

The handgrip maneuver increases afterload by increasing total peripheral resistance.

Correct Answer

A - Prolapse of the mitral valve

Explanation Why

[Mitral valve prolapse](#) leads to a [systolic murmur](#) that decreases in intensity with maneuvers that increase [afterload](#) (e.g., handgrip), which is seen here. However, this [murmur](#) is heard best at the cardiac apex (5th [intercostal space](#) in the [midclavicular line](#)). In addition, this [murmur](#) has a late [systolic](#) crescendo character, and it is preceded by a click due to tensing of the chordae tendinae.

B - Fusion of the right and left coronary leaflets

Explanation Why

Fusion of the right and left coronary leaflets of the [aortic valve](#) is the pathologic process that underlies a [bicuspid aortic valve](#), which can lead to a [systolic murmur](#) caused by [aortic stenosis](#) (AS). A [murmur](#) caused by AS is heard best at the right upper sternal border. In addition, this [murmur](#) has a crescendo-decrescendo character, and it decreases in intensity with maneuvers that increase [left ventricular afterload](#) (e.g., handgrip).

C - Defect in the atrial septum

Explanation Why

An [atrial septal defect](#) can cause a [systolic murmur](#) because of increased flow across the [pulmonary valve](#), but the [murmur](#) is heard best at the left upper sternal border. In addition, this [murmur](#) has a midsystolic crescendo-decrescendo character, and it is accompanied by a wide, fixed [split second heart sound](#).


D - Asymmetric hypertrophy of the left ventricle

Explanation Why

Asymmetric [left ventricular hypertrophy](#) is seen in [hypertrophic cardiomyopathy](#), which can cause a [murmur](#) as a result of [mitral regurgitation \(MR\)](#) or left ventricular outflow tract (LVOT) obstruction. A [murmur](#) due to [MR](#) is [holosystolic](#), and it increases in intensity with maneuvers that increase [left ventricular afterload](#) (e.g., handgrip), but this [murmur](#) is heard best at the cardiac apex (5th [intercostal space](#) in the [midclavicular line](#)). A [murmur](#) due to LVOT obstruction is heard best at the left lower sternal border, but this [murmur](#) has a midsystolic, crescendo-decrescendo character, and it decreases in intensity with maneuvers that increase [left ventricular afterload](#).

E - Defect in the ventricular septum

Image



Explanation Why

A [ventricular septal defect \(VSD\)](#), which causes a [left-to-right shunt](#), leads to a harsh [holosystolic murmur](#) because of [turbulent blood flow](#) across the defect. The [murmur](#) is heard best at the left lower sternal border ([tricuspid area](#)) and becomes louder with maneuvers that increase [left ventricular afterload](#) (e.g., handgrip) because of increased left-to-right shunting. The [murmur](#) is typically louder with smaller defects as blood rushes through a smaller space. Eventually, a [VSD](#) can lead to [pulmonary hypertension](#) and a reversal in blood flow ([Eisenmenger syndrome](#)), in which case the [murmur](#) may disappear or no longer become louder with increased LV [afterload](#) because of a countering [right-to-left shunting](#).

F - Failure of the ductus arteriosus to close

Explanation Why

The failure of the [ductus arteriosus](#) to close ([patent ductus arteriosus](#)) leads to a [murmur](#) heard best at the left infraclavicular region. This [murmur](#) occurs during both [diastole](#) and [systole](#) (i.e., a [continuous murmur](#)), unlike the harsh [holosystolic murmur](#) seen here.

Question # 31

A 28-year-old woman, gravida 1, para 0, at 32 weeks' gestation is admitted to the hospital for the management of elevated blood pressures. On admission, her pulse is 81/min, and blood pressure is 165/89 mm Hg. Treatment with an intravenous drug is initiated. Two days after admission, she has a headache and palpitations. Her pulse is 116/min and regular, and blood pressure is 124/80 mm Hg. Physical examination shows pitting edema of both lower extremities that was not present on admission. This patient most likely was given a drug that predominantly acts by which of the following mechanisms?

	Answer	Image
A	Inhibition of β_1 , β_2 , and α_1 receptors	
B	Activation of α_2 adrenergic receptors	
C	Inhibition of angiotensin II production	
D	Inhibition of sodium reabsorption	
E	Direct dilation of the arterioles	
F	Inhibition of nondihydropyridine calcium channels	

Hint

This woman has likely been treated with intravenous hydralazine, a first-line treatment for severe hypertension during pregnancy (i.e., systolic blood pressure ≥ 160 mm Hg and/or diastolic blood pressure ≥ 110 mm Hg).

Correct Answer

A - Inhibition of β_1 , β_2 , and α_1 receptors

Explanation Why

[Labetalol](#) is a nonselective [\$\beta\$ -blocker](#) with additional α -blocking action used as a first-line treatment for severe [hypertension](#) during [pregnancy](#). Its action on [\$\beta\$ receptors](#) decreases [heart rate](#) and cardiac contractility. Its action on [\$\alpha\$ receptors](#) decreases [systemic vascular resistance](#). Common adverse effects of [labetalol](#) therapy include [orthostatic hypotension](#), [dizziness](#), and fatigue. [Tachycardia](#) and [edema](#), as seen in this case, are not typical.

B - Activation of α_2 adrenergic receptors

Explanation Why

[Methyldopa](#) is a central α_2 adrenergic agonist used as a first-line treatment for severe [hypertension](#) during [pregnancy](#). It decreases [sympathetic](#) outflow to the [heart](#), [kidneys](#), and peripheral vasculature to reduce blood pressure. Peripheral [edema](#) is a common adverse effect of [methyldopa](#) therapy. However, other adverse effects include [orthostatic hypotension](#), sedation, and [bradycardia](#) (not [tachycardia](#)), which are not seen in this case.

C - Inhibition of angiotensin II production

Explanation Why

ACE inhibitors inhibit the conversion of [angiotensin I](#) to [angiotensin II](#) and are commonly used to treat [hypertension](#); however, their use is contraindicated during [pregnancy](#) due to their [teratogenic](#) effects on fetal [renal development](#). Common adverse effects of ACE inhibitor therapy include [cough](#), [angioedema](#), and [hyperkalemia](#).

D - Inhibition of sodium reabsorption

Explanation Why

[Thiazide diuretics](#) inhibit the reabsorption of sodium in the [distal convoluted tubule](#) (which increases the excretion of sodium, water, potassium, and hydrogen ions) and are commonly used to treat [hypertension](#). Although not typically initiated during [pregnancy](#), patients treated with [thiazide diuretics](#) prior to [pregnancy](#) can continue to use them throughout the gestational period. Common adverse effects of [thiazide diuretic](#) therapy include [hypokalemia](#), [hyponatremia](#), [hyperglycemia](#), [hyperlipidemia](#), [hyperuricemia](#), and [hypercalcemia](#), but not peripheral [edema](#).

E - Direct dilation of the arterioles

Explanation But

[Nifedipine](#), a [dihydropyridine calcium channel blocker](#) also used to treat [hypertension](#) during [pregnancy](#), has a similar adverse effect profile.

Explanation Why

[Hydralazine](#) is a direct peripheral [vasodilator](#) that preferentially affects arteriolar [smooth muscle](#) and increases the intracellular concentration of [cGMP](#), which reduces [afterload](#) and blood pressure. Common adverse effects of [hydralazine](#) therapy include [headache](#) (due to direct [vasodilation](#)), [tachycardia](#) (due to reflex [sympathetic](#) activation), and peripheral [edema](#) (due to increased [hydrostatic pressure](#), retention of sodium and water, and fluid extravasation). To reduce these adverse effects, direct arteriolar [vasodilators](#) are often coadministered with a [β-blocker](#) and/or a [diuretic](#).

F - Inhibition of nondihydropyridine calcium channels

Explanation But

[Nifedipine](#) is used to treat [hypertension](#) during [pregnancy](#) and causes [headache](#), [tachycardia](#), and peripheral [edema](#) as seen in this case. However, [nifedipine](#) is a [dihydropyridine calcium channel](#)

[blocker](#), not a [nondihydropyridine calcium channel blocker](#).

Explanation Why

[Verapamil](#) is a [nondihydropyridine calcium channel blocker](#) sometimes used to treat [supraventricular tachycardia](#) during [pregnancy](#). It is not a preferred agent for the management of [hypertension](#) during [pregnancy](#). Common adverse effects of [verapamil](#) therapy include [edema](#), [bradycardia](#) (not [tachycardia](#)), [atrioventricular block](#), and [constipation](#).

Question # 32

An investigator is studying the physiological response during congestive heart failure exacerbations in patients with systolic heart failure. A hormone released by ventricular cardiomyocytes in response to increased wall stress is isolated from a patient's blood sample. The intracellular mechanism by which this hormone acts is most similar to the effect of which of the following substances?

	Answer	Image
A	Thrombopoietin	
B	Nitric oxide	
C	Human chorionic gonadotropin	
D	Aldosterone	
E	Angiotensin II	
F	Platelet-derived growth factor	

Hint

B-type natriuretic peptide (BNP) is a peptide hormone released by ventricular cardiac myocytes in response to increased wall stress. It promotes the excretion of sodium and water from the kidneys.

Correct Answer


A - Thrombopoietin

Explanation Why

[Thrombopoietin](#) acts via the [JAK-STAT](#) signaling pathway. [BNP](#) has a different intracellular signaling mechanism.

B - Nitric oxide

Image



Explanation But

Certain drugs exploit the intracellular effects of [cGMP](#) for therapeutic benefit (e.g., [nesiritide](#) for [heart failure](#) and [PDE-5 inhibitors](#) for [erectile dysfunction](#) and [pulmonary hypertension](#)).

Explanation Why

[ANP](#), [BNP](#), and [nitric oxide](#) ([endothelium](#)-derived relaxation factor) act by increasing the activity of [guanylate cyclase](#) and thereby producing [cGMP](#), which activates protein kinase G (PKG). In the [distal](#) collecting tubule and [collecting duct](#), PKG [phosphorylates](#) Na^+ channels and decreases their activity. This effect is partially responsible for the [natriuretic](#) effects of [ANP](#) and [BNP](#). In [smooth muscles](#) (e.g., [tunica media](#), muscularis externa), [cGMP](#) activates [myosin light chain phosphatase](#), which causes relaxation. This process explains NO's generalized [vasodilatory](#) effect and the [vasodilatory](#) effects of [ANP](#) and [BNP](#) on the [afferent arteriole](#). However, for unknown reasons, [ANP](#) and [BNP](#) cause vasoconstriction in the [efferent arteriole](#). The net effect of [efferent arteriole](#) constriction and [afferent arteriole](#) dilation is increased [GFR](#), which also causes [natriuresis](#).

C - Human chorionic gonadotropin

Explanation Why

[Human chorionic gonadotropin](#) acts via the [cyclic adenosine monophosphate](#) ([cAMP](#)) signaling pathway. [BNP](#) acts via a different cyclic [nucleotide](#).

D - Aldosterone

Explanation Why

[Aldosterone](#) acts by binding to a [nuclear receptor](#), thereby increasing the [translation](#) of [epithelial sodium channels](#) ([ENaC](#)) in the [collecting duct](#). [BNP](#) does act on the [collecting duct](#) and decreases the activity of [ENaC](#), but it does so via a different signaling mechanism.

E - Angiotensin II

Explanation Why

[Angiotensin II](#) acts via the [inositol triphosphate](#) signaling pathway. [BNP](#) has a different intracellular

signaling mechanism.

F - Platelet-derived growth factor

Explanation Why

[Platelet-derived growth factor](#) acts via the mitogen-activated protein [kinases](#). [BNP](#) has a different intracellular signaling mechanism.

Question # 33

An investigator is comparing the risk of adverse effects among various antiarrhythmic medications. One of the drugs being studied primarily acts by blocking the outward flow of K^+ during myocyte repolarization. Further investigation shows that the use of this drug is associated with a lower rate of ventricular tachycardia, ventricular fibrillation, and torsade de pointes when compared to similar drugs. Which of the following drugs is most likely being studied?

	Answer	Image
A	Esmolol	
B	Sotalol	
C	Mexiletine	
D	Verapamil	
E	Amiodarone	

Hint

This medication is a class III antiarrhythmic that increases effective refractory periods (ERP) and QT intervals. It is associated with thyroid abnormalities.

Correct Answer

A - Esmolol

Explanation Why

[Esmolol](#) is a cardioselective β_1 -blocker with [class II antiarrhythmic drug](#) properties. It is used in the acute [treatment of atrial fibrillation](#), [atrial flutter](#), and [supraventricular tachycardia](#). It inhibits β -[adrenergic](#) activation of [adenylate cyclase](#) and can delay conduction velocity. Unlike the drug being studied, [esmolol](#) has no effect on potassium channels.

B - Sotalol

Explanation Why

[Sotalol](#) is a [nonselective beta blocker](#) with potent class II and [class III antiarrhythmic](#) properties. It is used in the treatment of [supraventricular arrhythmias](#) and [ventricular arrhythmias](#). Like the drug being investigated, [sotalol](#) can also block potassium channels during [myocyte repolarization](#), prolonging the [action potential](#) duration and [effective refractory period](#). Unlike the drug being studied, [sotalol](#) has high arrhythmogenic potential and carries a high risk of inducing [ventricular arrhythmias](#), such as [torsades de pointes](#).

C - Mexiletine

Explanation Why

[Mexiletine](#) is a [class IB antiarrhythmic drug](#) that primarily acts by blocking sodium channels. It is typically used to treat [ventricular arrhythmias](#) (especially following [myocardial infarction](#)) and [digitalis](#)-induced [cardiac arrhythmias](#). Unlike the drug being investigated, [mexiletine](#) has a minimal effect on the [action potential](#) duration and has no effect on potassium channel blockade.

D - Verapamil

Explanation Why

[Verapamil](#) is a [nondihydropyridine calcium channel blocker](#) with [class IV antiarrhythmic drug](#) properties. [Class IV antiarrhythmic drugs](#) prolong the [repolarization](#) of the [AV node](#) and can increase effective refractory periods. They are used in the treatment of [paroxysmal supraventricular tachycardia](#), [atrial fibrillation](#), [atrial flutter](#), and [multifocal atrial tachycardia](#). Unlike the drug under investigation, [verapamil](#) has no effect on potassium channels.

E - Amiodarone

Explanation But

[Adverse events](#) associated with [amiodarone](#) include [thyroid](#) and [liver](#) dysfunction, pulmonary [fibrosis](#), [peripheral neuropathy](#), corneal deposits, and [photosensitivity](#).

Explanation Why

[Amiodarone](#) (a [class III antiarrhythmic drug](#)) acts primarily by blocking outward potassium channels during [myocyte repolarization](#), in addition to blocking sodium and calcium channels. [Class III antiarrhythmic](#) drugs prolong the [action potential](#) duration (APD) and [effective refractory period](#) (ERP), with no delay in conduction velocity. Like other class III agents, [amiodarone](#) can cause [QT prolongation](#), although it carries a much lower risk of [arrhythmia](#) and [torsades de pointes](#) than other drugs of the same class.

Question # 34

A 34-year-old woman, gravida 2, para 2, is admitted to the hospital because of shortness of breath and fatigue 2 weeks after delivery of a full-term female newborn. She has no history of major medical illness. Cardiac examination on admission shows an S3 gallop and a grade 2/6 holosystolic murmur heard best at the apex. Treatment is initiated with intravenous furosemide and captopril. Her symptoms resolve, and 3 weeks later, cardiac examination shows no murmur. Which of the following is the most likely explanation for the initial auscultation findings?

	Answer	Image
A	Mitral annular dilatation	
B	Myxomatous mitral valve degeneration	
C	Perivalvular abscess	
D	Mitral valve leaflet fibrosis	
E	Papillary muscle rupture	
F	Mitral annular calcification	


Hint

The presentation of shortness of breath, fatigue, and a murmur of mitral regurgitation in the early peripartum period suggests peripartum cardiomyopathy.

Correct Answer

A - Mitral annular dilatation

Image



Explanation Why

[Peripartum cardiomyopathy](#) is a type of [dilated cardiomyopathy](#). Patients with [dilated cardiomyopathy](#) can develop [mitral regurgitation](#) caused by [left ventricular failure](#) and/or dilation of the [mitral valve](#) annulus. [Dilated cardiomyopathy](#) caused by [peripartum cardiomyopathy](#) is typically reversible. [Heart failure](#) (e.g., [dyspnea](#), [S3 gallop](#)) and [mitral regurgitation](#) can be treated with [diuretics](#).

B - Myxomatous mitral valve degeneration

Explanation Why

Myxomatous degeneration of the [mitral valve](#), which occurs in [connective tissue diseases](#) (e.g., [Marfan syndrome](#)), can lead to [mitral regurgitation](#); but such a valvular defect is irreversible without surgery.

C - Perivalvular abscess

Explanation Why

Perivalvular abscesses occur in individuals with [infective endocarditis](#) and can cause severe [mitral regurgitation](#), which is seen here. However, valvular defects due to [infective endocarditis](#) are usually irreversible and require treatment with high-dose [antibiotic therapy](#) as well as surgical repair.

D - Mitral valve leaflet fibrosis

Explanation Why

[Fibrosis](#) of the [mitral valve](#) leaflets, which is seen in [rheumatic heart disease](#) (RHD), can cause [mitral regurgitation](#) or [mitral stenosis](#); but valvular defects due to RHD are irreversible and require surgical repair.

E - Papillary muscle rupture

Explanation Why

[Papillary muscle rupture](#), which typically occurs following [myocardial infarction](#), can lead to [mitral](#)

[regurgitation](#) but is irreversible without surgery.

F - Mitral annular calcification

Explanation Why

Mitral annular calcification can lead to [mitral regurgitation](#), but this condition typically occurs in the older population. Furthermore, mitral annular calcification is irreversible without surgery.

Question # 35

A 42-year-old man is admitted to the hospital for pain and swelling in his right foot. His temperature is 39.7°C (103.5°F), pulse is 116/min, respirations are 23/min, and blood pressure is 69/39 mmHg. A drug is administered via a peripheral intravenous line that works primarily by increasing inositol trisphosphate concentrations in arteriolar smooth muscle cells. Eight hours later, the patient has pain at the right antecubital fossa. Examination shows the skin around the intravenous line site to be pale and cool to touch. After discontinuing the infusion, which of the following is the most appropriate pharmacotherapy to prevent further tissue injury in this patient?

	Answer	Image
A	Procaine	
B	Phentolamine	
C	Tamsulosin	
D	Conivaptan	
E	Heparin	
F	Midodrine	

Hint

This patient was given a vasopressor (most likely norepinephrine) for his hypotension. Vasopressors are not usually administered via peripheral IV lines because of the risk of extravasation and local tissue necrosis.

Correct Answer

A - Procaine

Explanation Why

[Procaine](#) is an [ester-type local anesthetic](#) that can be used for minor surgical procedures and [spinal anesthesia](#). It has not been proven effective in managing extravasation caused by vasopressor agents.

B - Phentolamine

Explanation Why

The [ischemic](#) tissue damage in this patient is caused by persistent vasoconstriction due to activation of [\$\alpha_1\$ receptors](#) by the extravasated pressor agent, which increases arteriolar [smooth muscle IP₃](#). As a short-acting, nonselective α -antagonist, phentolamine counteracts the [\$\alpha_1\$ receptor](#)-mediated vasoconstriction, improving blood flow to the damaged area and preventing further tissue loss. [Terbutaline](#) and topical [nitroglycerin](#) ointment may also be considered in cases of pressor extravasation.

C - Tamsulosin

Explanation Why

[Tamsulosin](#) is an α_1 -antagonist that is most commonly used for the treatment of BPH. [Tamsulosin](#) acts selectively on α_{1A} receptors in the [prostate gland](#) and has only minimal effect on peripheral vasomotor tone.

D - Conivaptan

Explanation Why

[Conivaptan](#) acts as an [ADH](#) inhibitor and is used in the treatment of [SIADH](#). It has not been proven effective in managing extravasation caused by vasopressor agents.

E - Heparin

Explanation Why

[Heparin](#) is most commonly used for the treatment of thrombosis or embolic disease, which could also manifest with a cool, pale extremity. However, signs of [ischemia](#) restricted to the [antecubital fossa](#), as seen in this patient, are more suggestive of [ischemia](#) due to vasopressor extravasation, in which case [heparin](#) would not be effective.



F - Midodrine

Explanation Why

[Midodrine](#) is an $\alpha 1$ agonist that promotes vasoconstriction and is used for autonomic insufficiency and [orthostatic hypotension](#). It would only worsen the symptoms of vasopressor extravasation in this patient.

Question # 36

A 65-year-old man with hypertension and type 2 diabetes mellitus is brought to the emergency department 20 minutes after the onset of severe anterior chest pain and shortness of breath. He has smoked one pack of cigarettes daily for 30 years. He appears distressed. His pulse is 116/min, respirations are 22/min, and blood pressure is 156/88 mm Hg. Pulse oximetry on room air shows an oxygen saturation of 98%. A grade 3/6, high-pitched, blowing, diastolic murmur is heard over the right upper sternal border. Which of the following is the most likely cause of this patient's symptoms?

	Answer	Image
A	Tear in the tunica intima	
B	Fibrofatty plaque in the aortic wall	
C	Obstruction of the pulmonary arteries	

	Answer	Image
D	Rupture of a bulla in the lung	
E	Perforation of the esophageal wall	
F	Infarction of the ventricular wall	


Hint

Dyspnea with a high-pitched, blowing, diastolic murmur at the right upper sternal border suggests aortic regurgitation.

Correct Answer

A - Tear in the tunica intima

Image



Explanation But

Nonmodifiable [risk factors](#) for [aortic dissection](#) include advanced age, male sex, and certain genetic conditions that weaken the [tunica media](#) (e.g., [Marfan syndrome](#), vascular [Ehlers-Danlos syndrome](#), [Turner syndrome](#)).


Explanation Why

An [aortic dissection](#) is typically caused by a tear in the [tunica intima](#) and subsequent longitudinal extension of the dissecting [hematoma](#). [Hypertension](#), which increases the risk of [endothelial](#) injury, is the most common predisposing factor for [aortic dissection](#). Most [aortic dissections](#) begin in the ascending portion of the [thoracic aorta](#) because this is the region of maximum hydraulic stress. Patients present with sudden and severe [pain](#) in the chest or interscapular region. The [pain](#) can migrate as the dissection progresses. [Proximal](#) extension of the dissection into the [aortic valve](#) ring can cause acute [aortic regurgitation](#) (AR), as seen here. Other features of [aortic dissection](#) include [syncope](#), diminished [distal](#) pulses, and asymmetrical blood pressure and [pulse](#) readings between

limbs. Unlike the [murmur](#) of AR from [aortic valve](#) disease, which is prominent along the left sternal border, the [murmur](#) of AR from [aortic dissection](#) is prominent along the right sternal border.

B - Fibrofatty plaque in the aortic wall

Image



Explanation Why

The formation of a fibrofatty plaque in the aortic wall is characteristic of [atherosclerosis](#), which typically involves the abdominal portion of the [descending aorta](#). [Atherosclerosis](#) is implicated in the formation of abdominal [aortic aneurysms](#); [atheroma](#) ulceration has been reported in some patients with abdominal [aortic dissections](#) and descending thoracic [aortic dissections](#). Although both [aortic dissection](#) and [atherosclerosis](#) share similar [risk factors](#) (e.g., smoking, [hypertension](#), advanced age, and male sex), [atherosclerosis](#) is not a major [risk factor](#) for most [aortic dissections](#), which tend to occur in the [ascending aorta](#).

C - Obstruction of the pulmonary arteries

Explanation Why

Obstruction of the [pulmonary arteries](#) is seen in [pulmonary embolism](#), which can manifest with acute-onset [dyspnea](#), [chest pain](#), [tachypnea](#), and [tachycardia](#). However, [pulmonary embolism](#) would not explain the [aortic regurgitation](#) seen here.

D - Rupture of a bulla in the lung

Explanation Why

Rupture of a [bulla](#) in the [lung](#) causes [pneumothorax](#), which can manifest with [dyspnea](#), [chest pain](#), [tachypnea](#), and [tachycardia](#). However, [pneumothorax](#) would not explain the [aortic regurgitation](#) seen here. Instead, chest auscultation would reveal decreased breath sounds caused by [pneumothorax](#).

E - Perforation of the esophageal wall

Explanation Why

[Esophageal rupture](#) can manifest with [chest pain](#), [tachycardia](#), [tachypnea](#), and [dyspnea](#) (due to [mediastinitis](#) and/or [pleural effusion](#)). However, [esophageal rupture](#) would not cause a [murmur](#). Instead, chest auscultation might reveal a crunching sound over the precordium that is synchronous with each heartbeat ([Hamman sign](#)) due to a [pneumomediastinum](#) and/or decreased breath sounds due to a [pleural effusion](#). Furthermore, [esophageal rupture](#) usually has a distinct precipitating cause (e.g., retching or forceful vomiting), which is not seen here.

F - Infarction of the ventricular wall

Explanation Why

[Infarction](#) of the ventricular wall ([myocardial infarction](#)) can cause acute-onset [chest pain](#), [dyspnea](#), [tachypnea](#), [tachycardia](#), and a [murmur](#) due to [mitral regurgitation](#) that is caused by [papillary muscle rupture](#). However, the [murmur](#) due to [mitral regurgitation](#) is [systolic](#) rather than [diastolic](#) and heard best at the cardiac apex, not the right upper sternal border. Moreover, [papillary muscle rupture](#) typically occurs 2–7 days after [infarction](#), not at the outset.

Question # 37

A 79-year-old woman with type 2 diabetes mellitus and hypertension undergoes ^{99m}Tc cardiac scintigraphy for the evaluation of a 3-month history of retrosternal chest tightness on exertion. The patient's symptoms are reproduced following the administration of dipyridamole. A repeat ECG shows new ST depression and T wave inversion in leads V₅ and V₆. Which of the following is the most likely underlying mechanism of this patient's signs and symptoms during the procedure?

	Answer	Image
A	Vasospasm of the coronary arteries	
B	Transient atrioventricular nodal blockade	
C	Reduced left ventricular preload	
D	Dilation of coronary vasculature	
E	Ruptured cholesterol plaque within a coronary vessel	
F	Increased myocardial oxygen demand	

Hint

This patient underwent dipyridamole stress testing for evaluation of suspected coronary artery disease. The results of her pharmacologic stress test are consistent with decreased blood flow in the anterolateral region. In addition to being a platelet phosphodiesterase inhibitor, dipyridamole prevents the breakdown of adenosine by inhibiting adenosine deaminase.

Correct Answer

A - Vasospasm of the coronary arteries

Explanation Why

Vasospasm of [coronary arteries](#), which is the cause of [Prinzmetal angina](#), would lead to recurrent episodes of chest tightness and the [ECG](#) changes seen in this patient. However, patients with [Prinzmetal angina](#) typically develop symptoms at rest, unlike this patient who reported chest tightness on exertion. The onset of [chest pain](#) after the administration of [dipyridamole](#), which induces [vasodilation](#) rather than vasospasm, suggests a different cause for this patient's symptoms.

B - Transient atrioventricular nodal blockade

Explanation Why

A 2nd or 3rd degree [AV block](#) can occasionally cause chest tightness and [dipyridamole](#) has caused an [AV block](#) in rare cases. However, an [AV block](#) is characterized by [bradycardia](#), a prolonged [PR interval](#), dropped [QRS complexes](#), or atrioventricular dissociation. The [ECG](#) findings in this patient are not consistent with an [AV block](#).

C - Reduced left ventricular preload

Explanation Why

[Left ventricular preload](#) is decreased by venous [vasodilators](#) (e.g., [nitroglycerin](#)) and by [hypovolemia](#). [Dipyridamole](#) can decrease [left ventricular afterload](#) by inducing [vasodilation](#) of peripheral [arteries](#) but it does not significantly alter [left ventricular preload](#).

D - Dilation of coronary vasculature

Explanation But

In the case of a [cardiac exercise stress test](#) or a dobutamine stress test, cardiac [ischemia](#) is reproduced by increasing [heart rate](#) and contractility, which, in turn, increases myocardial oxygen demand. The stress test should be terminated as soon as the patient experiences [chest pain](#) and/or [ST elevations](#).

Explanation Why

[Dipyridamole](#) increases adenosine levels, which, in turn, causes coronary [vasodilation](#) via the activation of adenosine A2A receptors. [Coronary steal syndrome](#) is [myocardial ischemia](#) caused by redistribution of blood flow. The [rigidity](#) of severely stenotic [arteries](#) allows only a low coronary flow reserve. When [coronary arteries](#) dilate, blood is, therefore, diverted to healthier vessels and collaterals. This leads to tissue [ischemia distal](#) to the stenosis in the [coronary artery](#), which manifests as chest tightness and new [ECG](#) changes. This phenomenon is the underlying principle of the adenosine or [dipyridamole pharmacological stress test](#).

E - Ruptured cholesterol plaque within a coronary vessel

Explanation Why

Thrombotic occlusion of a coronary vessel following the rupture of a [cholesterol](#) plaque is the process that underlies [acute coronary syndrome \(ACS\)](#), which would cause chest tightness and the [ECG](#) changes seen here. While there is a risk for [ACS](#) with every stress test, it is low with a [dipyridamole](#) stress test because [dipyridamole](#) inhibits [platelet aggregation](#). A different mechanism is more likely to have provoked this patient's chest tightness.

F - Increased myocardial oxygen demand

Explanation Why


Myocardial oxygen demand (MVO₂) is increased during the dobutamine stress test and [cardiac](#)

[exercise stress test](#). [Dipyridamole](#) alters coronary perfusion and affects [myocardial](#) oxygen supply but it does not significantly alter MVO₂.

Question # 38

A 32-year-old woman comes to the emergency department because of a 12-hour history of a severe headache. She does not smoke or use illicit drugs. Her blood pressure at admission is 180/125 mm Hg. Physical examination shows a bruit in the epigastric region. Fundoscopy shows bilateral optic disc swelling. Which of the following investigations is most likely to confirm the diagnosis?

	Answer	Image
A	Serum IGF-I level	
B	Urinary catecholamine metabolites	
C	Echocardiography	
D	Serum 17-hydroxyprogesterone level	
E	Serum PTH level	
F	High-dose dexamethasone suppression test	
G	Oral sodium loading test	

	Answer	Image
H	CT angiography	 A CT angiography image of the renal vasculature. The image shows a cross-section of the kidneys and the renal arteries. A prominent, rounded, and dilated segment of the renal artery is visible, indicating a renal artery aneurysm. The surrounding renal parenchyma and other vessels appear normal.

Hint

In a young patient with a hypertensive emergency (blood pressure of 185/125 mm Hg, papilledema, severe headache) and an epigastric or flank bruit on examination, fibromuscular dysplasia (FMD) is the most likely diagnosis.

Correct Answer

A - Serum IGF-I level

Explanation Why

Serum [IGF-I](#) levels are used in the evaluation of [acromegaly](#). Although [acromegaly](#) can cause [secondary hypertension](#), it is also associated with other clinical features, such as a large [skull](#), coarse facies, and widened hands, fingers, and feet. In addition, [acromegaly](#) would not explain this patient's abdominal [bruit](#).

B - Urinary catecholamine metabolites

Explanation Why

Urinary [catecholamine](#) metabolites confirm the diagnosis of [pheochromocytoma](#). Although [pheochromocytoma](#) can manifest with episodic [hypertension](#) and [headaches](#), it would not explain this patient's abdominal [bruit](#).

C - Echocardiography

Explanation Why

[Echocardiography](#) with Doppler is used e.g., to confirm [coarctation of the aorta](#). Although [coarctation of the aorta](#) can lead to [hypertension](#), other symptoms are also typically present, such as [hypotension](#) and [claudication](#) of the lower extremities, or a [continuous murmur](#) below the left clavicle and between the shoulder blades. An abdominal [bruit](#), however, would not be expected. [Echocardiography](#) has no role in diagnosing [fibromuscular dysplasia](#).

D - Serum 17-hydroxyprogesterone level

Explanation Why

Serum [17-hydroxyprogesterone](#) levels are used to screen for [congenital adrenal hyperplasia \(CAH\)](#). Although [congenital adrenal hyperplasia](#) can cause [secondary hypertension](#), it is often associated with [precocious puberty](#) and electrolyte abnormalities and would not explain this patient's abdominal [bruit](#). In addition, all [newborns](#) in the US are screened for [CAH](#); therefore, a delayed diagnosis at the age of 32 years would be unlikely.

E - Serum PTH level

Explanation Why

Serum [PTH](#) levels are used in the diagnostic evaluation of [hyperparathyroidism](#). [Primary hyperparathyroidism](#) can lead to [hypertension](#) due to progressive calcification of the [arteries](#) and [heart](#) muscle. However, this condition is typically associated with other findings, such as [nephrolithiasis](#), [constipation](#), [gastric ulcers](#), psychological symptoms (e.g., depression, fatigue), and/or bone, muscle, and [joint pain](#). In addition, [primary hyperparathyroidism](#) would not explain this patient's abdominal [bruit](#).

F - High-dose dexamethasone suppression test

Explanation Why

A high-dose [dexamethasone suppression test](#) is used to confirm [Cushing syndrome](#). While [Cushing syndrome](#) is a common cause of [secondary hypertension](#), it is usually accompanied by other symptoms, such as [obesity](#), [hirsutism](#), and easily bruisable [skin](#) with [stretch marks](#). In addition, [Cushing syndrome](#) would not explain this patient's abdominal [bruit](#).


G - Oral sodium loading test

Explanation Why

[Oral sodium loading test](#) is used to confirm [primary hyperaldosteronism](#). Although [primary hyperaldosteronism](#) is a common cause of [secondary hypertension](#) in young people, symptoms of [hypokalemia](#), such as fatigue or muscle weakness, are also typically present. In addition, [primary hyperaldosteronism](#) would not explain this patient's abdominal [bruit](#).

H - CT angiography

Image



Explanation Why

CT angiography is a noninvasive imaging method that has a high [accuracy](#) in confirming the diagnosis of [fibromuscular dysplasia \(FMD\)](#), which most commonly affects the renal and carotid [arteries](#). [Renal artery stenosis](#) due to [FMD](#) leads to [hypertension](#), as seen here. Imaging typically


shows a so-called “string of beads” appearance of the [renal arteries](#) due to segmental [hyperplasia](#) of the [tunica media](#). Other imaging methods that can be used to diagnose [FMD](#) are [duplex ultrasonography](#) and magnetic resonance angiography.

Question # 39

A 43-year-old man from Chile comes to the physician because of a 1-day history of upper back pain and difficulty swallowing. He has had pain in his shoulder and knee joints over the past 10 years. He is 190 cm (6 ft 3 in) tall and weighs 70.3 kg (155 lb); BMI is 19.4 kg/m². His blood pressure is 142/86 mm Hg in the right arm and 130/70 mg Hg in the left arm. Physical examination shows a depression in the sternum and a grade 3/6 diastolic murmur at the right upper sternal border. A CT scan of the chest with contrast is shown. Which of the following is the most likely underlying cause of this patient's condition?



	Answer	Image
A	Protozoal infection	
B	Autoimmune valve damage	
C	Granulomatous inflammation	

	Answer	Image
D	Systemic fibroblast proliferation	
E	Congenital aortic narrowing	
F	Cystic medial degeneration	 Marfan syndrome Epidemiology Prevalence: 1-5/10,000 Etiology Mutation of fibrillin-1 gene (FBN1) on chromosome 15, autosomal dominant inheritance Complications Primarily cardiovascular (e.g., aortic dissections) Life expectancy Can be normal, given timely diagnosis and appropriate management (cardiological check-ups) Key features shown in the diagram: - Ectopia lentis, severe myopia - Heart valve defects - Pectus deformity (pectus carinatum or pectus excavatum) - Aortic disease (e.g., aortic dissection, aortic aneurysm) - Arachnodactyly, joint hypermobility - Hyperextensible skin - Tall stature - Berry aneurysms - High-arched palate - Spinal deformities (scoliosis, hyperkyphosis) - Long extremities - Pes planus, hindfoot valgus
G	Atheroma formation	

Hint

This axial CT shows an ascending aortic aneurysm, which is often associated with an aortic regurgitation murmur (diastolic murmur at the right upper sternal border) caused by dilation of the aortic valve annulus. Dissection of the weakened aneurysmal wall can cause an acute increase in the size of an aneurysm, manifesting with back pain, discordant blood pressure readings in the upper limbs, and difficulty swallowing (dysphagia aortica due to esophageal compression).

Correct Answer

A - Protozoal infection

Explanation Why

Infection with the protozoan [Trypanosoma cruzi](#) causes [Chagas disease](#), which is [endemic](#) to South America. This disease can cause [dysphagia](#) (due to [achalasia](#)) as well as [aortic regurgitation](#) (due to [dilated cardiomyopathy](#)), which are seen here. However, [Chagas disease](#) would not cause [pectus excavatum](#), discordant blood pressure readings, or dilation of the aorta.

B - Autoimmune valve damage

Explanation Why

Autoimmune valve damage is the cause of [rheumatic heart disease](#) (RHD), which, in the US, is most commonly seen in the immigrant population. RHD can cause a [diastolic murmur](#) (due to [mitral stenosis](#)), and, rarely, difficulty [swallowing](#) (dysphagia megalatriensis due to [left atrial enlargement](#)). [Rheumatic fever](#), which is the pathologic precursor to RHD, would also cause [pain](#) in large [joints](#) such as the [knee](#). However, the [joint pain](#) would subside after the acute episode of [rheumatic fever](#), and the [diastolic murmur](#) of [mitral stenosis](#) would be heard best at the cardiac apex. This patient's [murmur](#) is caused by [aortic regurgitation](#), which is not common in RHD. Most importantly, RHD would not cause dilation of the [ascending aorta](#).

C - Granulomatous inflammation

Explanation Why

[Granulomatous inflammation](#) is seen in [giant cell arteritis](#) ([GCA](#)), which predisposes to the formation of ascending [aortic aneurysms](#) and can present with back and shoulder [pain](#) (due to concurrent [polymyalgia rheumatica](#)). However, [aortic aneurysm](#) usually occurs late in the course of [GCA](#) and the disease virtually never affects patients < 50 years.

[Granulomatous inflammation](#) is also seen in [Takayasu arteritis](#), which can lead to chronic arthralgia

and discordant blood pressure readings between the upper limbs (due to [aortic arch](#) narrowing). [Takayasu arteritis](#) can, occasionally, lead to [aneurysmal](#) dilation of the [ascending aorta](#) with [aortic regurgitation](#) and [dysphagia](#), which is seen here. However, the condition typically occurs in Asian women and would usually be associated with other features, such as cutaneous nodules, [fever](#), or night sweats.

D - Systemic fibroblast proliferation

Explanation Why

Systemic [fibroblast proliferation](#) is a pathologic process that occurs in patients with [systemic sclerosis \(SSc\)](#), which can cause [dysphagia](#) (due to [esophageal dysmotility](#)), chronic arthralgia, and vascular disease. [Dysphagia](#) due to [SSc](#) usually has a subacute or chronic onset. In addition, arthralgia caused by [SSc](#) typically affects the small [joints](#) of the hands and feet, and the vasculopathy associated with [SSc](#) manifests as [Raynaud phenomenon](#), [telangiectasia](#), or [thromboembolic phenomena](#). Furthermore, SS would not cause [pectus excavatum](#), discordant blood pressure readings, [aortic regurgitation](#), or dilation of the [ascending aorta](#).


E - Congenital aortic narrowing

Explanation Why

Congenital narrowing of the [aortic arch](#) ([aortic coarctation](#)) can cause discordant blood pressure readings in the limbs. Additionally, a [bicuspid aortic valve](#), which is commonly associated with [aortic coarctation](#), can lead to [aortic regurgitation](#) ([diastolic murmur](#) at the right upper sternal border), which is seen here. However, in the case of [aortic coarctation](#), the discordance is usually between the blood pressure measurements of the upper and lower extremities because the coarctation is typically located just [distal](#) to the [left subclavian artery](#). Only in very rare instances would [aortic coarctation](#) cause disparate blood pressure measurements in the arms. Furthermore, this condition would not be expected to cause significant dilation of the [ascending aorta](#).

F - Cystic medial degeneration

Image



Explanation Why

This patient's tall stature and [pectus excavatum](#) suggest [Marfan syndrome](#). Individuals with [Marfan syndrome](#) have hypermobile [joints](#), which predisposes them to chronic arthralgia as a result of recurrent sprains, [bursitis](#), and/or early-onset [osteoarthritis](#). [Cystic medial degeneration](#), which occurs with [aging](#) and in [connective tissue disorders](#) such as [Marfan syndrome](#) and [Ehlers-Danlos syndrome](#), leads to the weakening of the aortic wall and [aneurysmal](#) dilation, which are seen here.

G - Atheroma formation

Explanation Why

Weakening of the vessel wall due to [atheroma](#) formation ([atherosclerosis](#)) is the most common cause of an [aortic aneurysm](#), and the [marfanoid features](#) in this patient (e.g., tall stature, [pectus excavatum](#))

might be due to [homocystinuria](#), which is a [risk factor](#) for early-onset [atherosclerosis](#). However, [aneurysms](#) due to [atherosclerosis](#) typically occur in the thoracic segment and the abdominal segment of the [descending aorta](#). The ascending [aortic aneurysm](#) in this patient suggests a different pathologic process.

Question # 40

During a clinical study evaluating the effects of exercise on muscle perfusion, 15 healthy individuals perform a 20-minute treadmill run at submaximal effort. Before and after the treadmill session, perfusion of the quadriceps muscle is evaluated with contrast-enhanced magnetic resonance imaging. The study shows a significant increase in muscle blood flow per unit of tissue mass. Which of the following local changes is most likely involved in the observed change in perfusion?

	Answer	Image
A	Increase in thromboxane A ₂	
B	Decrease in lactic acid	
C	Decrease in prostacyclin	
D	Increase in endothelin	
E	Increase in adenosine	
F	Decrease in potassium	

Hint

During exercise, the autoregulation of skeletal muscle blood flow leads to hyperemia secondary to accumulation of local metabolites.

Correct Answer

A - Increase in thromboxane A₂

Explanation Why

[Thromboxane A₂](#) (TXA₂) is a potent vasoconstrictor secreted by [platelets](#) and [endothelial](#) cells. An increase in TXA₂ would decrease rather than increase local blood flow.

B - Decrease in lactic acid

Explanation Why

[Lactic acid](#) accumulates in [skeletal muscle](#) during intense exercise as a result of [anaerobic](#) respiration. A decrease in [lactic acid](#) even during submaximal exercise testing is therefore unlikely. Moreover, since [lactic acid](#) has a [vasodilatory](#) effect, a decrease in [lactic acid](#) would not increase local blood flow.

C - Decrease in prostacyclin

Explanation Why

[Prostacyclin](#) (PGI₂), which is a [vasodilator](#), is released by vascular [endothelium](#) during exercise. An increase, not a decrease, of this metabolite leads to a rise in local blood flow.

D - Increase in endothelin

Explanation Why

Endothelin is a very potent vasoconstrictor secreted by [endothelial](#) cells. Thus, an increase in endothelin would decrease rather than increase local blood flow.

E - Increase in adenosine

Explanation Why

[Skeletal muscle](#) contraction during exercise requires hydrolyzation of [ATP](#) to [ADP](#). [ADP](#) is [hydrolyzed](#) to [AMP](#), which is then dephosphorylated to adenosine by a [5'-nucleotidase](#) bound to the [sarcolemma](#). Adenosine diffuses into [interstitial](#) tissue and binds to adenosine A_{2A} receptors on arteriolar [smooth muscle cells](#), which causes local [vasodilation](#). The other locally produced [vasodilatory](#) metabolites are [lactate](#), H⁺, K⁺, and CO₂. These metabolites can increase muscle blood flow by a factor of 20–30 during exercise.

F - Decrease in potassium

Explanation Why

During exercise, increased [depolarization](#) of [skeletal muscle](#) leads to an increase, not decrease, of extracellular potassium, which, in turn, induces vasodilatation by causing [hyperpolarization](#) and relaxation of vascular [smooth muscle](#). A decrease in potassium would not increase local blood flow.

Question # 1

A 51-year-old woman with a history of paroxysmal atrial fibrillation comes to the physician for a follow-up visit. She feels well and wants to discuss pausing her only current medication, flecainide. Her pulse is 75/min and regular, blood pressure is 125/75 mm Hg. Physical examination shows no abnormalities. An ECG shows a PR interval of 180 ms, QRS time of 120 ms, and corrected QT interval of 440 ms. Which of the following ECG changes is most likely to be seen on cardiac stress testing in this patient?

	Answer	Image
A	Prolonged QTc interval	
B	False-positive ST-segment depression	
C	Shortened PR interval	
D	Prolonged QRS complex	
E	Decreased maximal heart rate	

Hint

Flecainide is a class IC antiarrhythmic used in patients with paroxysmal supraventricular arrhythmias and no other structural heart disease. One of its cardinal features is use-dependence.

Correct Answer

A - Prolonged QTc interval

Explanation But

[Class IC antiarrhythmic drugs](#) such as [flecainide](#) may prolong the uncorrected [QT interval](#), which may predispose to [ventricular tachyarrhythmias](#) (e.g., torsade de pointes), especially in patients with structural heart disease, electrolyte disturbances, or [coronary artery disease](#).

Explanation Why

Prolongation of the [QTc](#) interval is a possible [ECG](#) change seen on [cardiac stress testing](#) in patients taking class IA and [class III antiarrhythmic drugs](#), which block potassium channels, resulting in a prolonged [repolarization](#) phase of the [cardiac action potential](#). [Flecainide](#) has minimal effects on the ventricular [repolarization](#) but can result in shortening (rather than prolongation) of the [QTc](#) interval on stress testing.

B - False-positive ST-segment depression

Explanation Why

False-positive [ST-segment depression](#) is a possible [ECG](#) change seen on [cardiac stress testing](#) in patients taking [digoxin](#). [Flecainide](#) has only very rarely been associated with false-positive [ST-segment depression](#).

C - Shortened PR interval

Explanation Why

Class I antiarrhythmic drugs act by blocking the sodium channels in [cardiomyocytes](#), which results in prolongation of the [PR interval](#), not shortening. In patients taking [flecainide](#), first- and [second-degree AV block](#) may be seen on [cardiac stress testing](#).

D - Prolonged QRS complex

Explanation Why

Class I antiarrhythmic drugs such as [flecainide](#) block the fast sodium channels responsible for phase 0 of the [myocardial action potential](#) (rapid upstroke and [depolarization](#)). The effect is an [action potential](#) with decreased slope and amplitude, which prolongs the spread of [depolarization](#) to neighboring cells, slowing conduction through the [myocardium](#). Since phase 0 for ventricular [myocytes](#) corresponds to the [QRS complex](#) on [ECG](#), [flecainide](#) prolongs the [QRS complex](#). This particular effect of class I antiarrhythmic drugs, especially of [class IC antiarrhythmic drugs](#), is use/state-dependent, i.e., it increases with the [heart rate](#) (e.g., during [cardiac stress testing](#)), as more sodium channels are recruited for cardiac contraction.

E - Decreased maximal heart rate

Explanation But

[Flecainide](#) may also have a limiting effect on the maximal [heart rate](#), but a different [ECG](#) change is much more common.

Explanation Why

Decreased maximal [heart rate](#) is an [ECG](#) change typically seen on [cardiac stress testing](#) in patients taking [beta blockers](#). [Beta blockers](#) inhibit the influence of the [sympathetic nervous system](#) on cardiac electrical activity, causing negative [chronotropic](#), [inotropic](#), and [dromotropic](#) effects that result in decreased [heart rate](#), decreased cardiac contractility, and decreased conduction (especially at the [AV node](#)), respectively.

Question # 2

An investigator is studying cardiovascular changes during exercise. He is following a 25-year-old female volunteer who is training for a marathon. Her training consists of running on a treadmill for increasing amounts of time. During her training sessions, laboratory studies show increased concentrations of adenosine and lactate in the active muscles. Which of the following is most likely a direct effect of these metabolites?

	Answer	Image
A	Decreased stroke volume	
B	Decreased capillary pressure	
C	Decreased systemic vascular resistance	
D	Increased venous return	
E	Increased mixed venous oxygen saturation	
F	Increased mean arterial pressure	

Hint

Adenosine and lactate are metabolites produced by skeletal muscle cells in response to hypoxia (e.g., during exercise). These metabolites cause local vasodilation to increase blood flow to tissues.

Correct Answer

A - Decreased stroke volume

Explanation Why

[Stroke volume](#) increases during exercise, in which the body compensates for increased muscle oxygen demand with increased [cardiac output](#). [Cardiac output](#) is the product of [stroke volume](#) and [heart rate](#), which are both likely increased in this volunteer as a result of [sympathetic](#) stimulation during her training session.

B - Decreased capillary pressure

Explanation Why

[Capillary](#) pressure increases during exercise because of increased blood flow in [vasodilated](#) larger vessels.

C - Decreased systemic vascular resistance

Explanation Why

[Systemic vascular resistance](#) is a measure of blood resistance through blood vessels. According to [Poiseuille equation](#), resistance through a tube is directly proportional to the length of the tube and the viscosity of the fluid, and inversely proportional to the [radius](#) to the 4th power. Adapting this equation for a blood vessel, it becomes apparent that [vasodilating](#) metabolites will directly decrease [systemic vascular resistance](#) by increasing the arteriolar [radius](#), resulting in a significant increase in blood supply to the exercised [skeletal muscle](#). Although [sympathetic](#) stimulation causes vasoconstricting effects during exercise, [vasodilating](#) metabolites help maintain [mean arterial pressure](#).

D - Increased venous return

Explanation Why

Increased [venous return](#) is typical during exercise, as muscle contractions compress systemic [veins](#) and thereby facilitate blood flow back to the [heart](#). However, this effect is not directly due to [vasodilating](#) metabolites.

E - Increased mixed venous oxygen saturation

Explanation Why

[Mixed venous oxygen saturation](#) is an indicator of whether [oxygen delivery](#) sufficiently meets the body's demands. During exercise, muscles have increased oxygen demands, resulting in increased oxygen extraction from the blood. With time, the [heart](#) compensates for this increased oxygen extraction with increased [cardiac output](#). As a result, there may be an initially decreased [mixed venous oxygen saturation](#) that normalizes during exercise; an increased [mixed venous oxygen saturation](#) would not be expected.


F - Increased mean arterial pressure

Explanation Why



[Mean arterial pressure \(MAP\)](#) during exercise should stay relatively constant but may slightly increase as [cardiac output](#) increases to compensate for increased blood flow through muscles. Increases in [MAP](#) are slight and not the direct result of [vasodilating](#) metabolites.

Question # 3

A 76-year-old man is admitted to the hospital for evaluation of sudden-onset chest pain. Three days after admission, he develops severe shortness of breath. Despite appropriate care, the patient dies. The heart at autopsy is shown. Which of the following most likely contributed to this patient's cause of death?



	Answer	Image
A	Occlusion of the posterior descending artery	

	Answer	Image
B	Exudative inflammation of the pericardium	
C	Pseudoaneurysmatic dilation of the left ventricle	
D	Rupture of the interventricular septum	
E	Bacterial infection of the mitral valve endocardium	


Hint

The image shows a rupture in the posteromedial papillary muscle. Papillary muscle rupture is life-threatening and can occur 2–7 days after myocardial infarction.

Correct Answer

A - Occlusion of the posterior descending artery

Image



Explanation But


Compared to the posteromedial [papillary muscle](#), the anterolateral [papillary muscle](#) is much less likely to rupture as a complication of [myocardial infarction](#) because it receives a dual blood supply from the [left anterior descending artery](#) and the [left circumflex artery](#).

Explanation Why

The [posterior descending artery \(PDA\)](#) arises from the [right coronary artery \(RCA\)](#), supplies the [posterior](#) aspect of the [heart](#), and provides the only blood supply to the posteromedial [papillary muscle](#). Occlusion of the [PDA](#) or the [RCA](#) causing [myocardial infarction](#), can result in [ischemic rupture](#) of the posteromedial [papillary muscle](#) as seen in this patient. Clinical manifestations of [papillary muscle rupture](#) include sudden-onset [hypotension](#), a [holosystolic murmur](#) (due to [mitral regurgitation](#)), and symptoms of [pulmonary edema](#) (e.g., [dyspnea](#), [cough](#)). Severe [mitral regurgitation](#) can cause acute [heart failure](#) and death.

B - Exudative inflammation of the pericardium

Image



Explanation Why

Postinfarction fibrinous pericarditis, which is characterized by exudative [inflammation](#) of the [pericardium](#), can cause [dyspnea](#) 1–3 days after [myocardial infarction](#). However, this complication is typically self-limited and thus unlikely to cause death in the acute postinfarction period. In addition, the gross appearance of the [heart](#) on autopsy of patients with this complication shows fibrinous exudate on the [pericardium](#), not a [papillary muscle rupture](#) as seen in this patient.

C - Pseudoaneurysmatic dilation of the left ventricle

Explanation Why

A ventricular pseudoaneurysm can form 3–14 days after [myocardial infarction](#) and cause [dyspnea](#). If left untreated, these [pseudoaneurysms](#) have a high risk of rupture and can cause [sudden cardiac](#)

[death](#). However, the gross appearance of the [heart](#) on autopsy of patients with these complications shows a bulging and lax [pericardium](#) that overlies the site of a ventricular wall rupture, not a [papillary muscle rupture](#) as seen in this patient.


D - Rupture of the interventricular septum

Explanation Why

The interventricular septum can rupture 3–5 days after a [myocardial infarction](#) due to weakening by the inflammatory response to [necrotic cardiomyocytes](#). This complication can result in death due to decreased [cardiac output](#) and severe [hypotension](#). However, the gross appearance of the [heart](#) on autopsy of patients with this complication shows a [ventricular septal defect](#), not a [papillary muscle rupture](#) as seen in this patient.

E - Bacterial infection of the mitral valve endocardium

Image




Explanation Why


[Infective endocarditis](#) can cause [dyspnea](#) and life-threatening cardiac decompensation as a result of severe valvular disease, [pulmonary edema](#), and/or [cardiac arrhythmias](#), but it wouldn't typically manifest with sudden-onset [chest pain](#). In addition, the gross appearance of the [heart](#) on autopsy of patients with this condition shows thickened [heart valves](#) with vegetations, ulcerations, and/or perforation, not a [papillary muscle rupture](#) as seen in this patient.

Question # 4

A 5-year-old boy is brought to the physician by his mother because she has noticed that he sometimes has difficulty breathing when he is playing outside. He is also easily fatigued by exertion. He has been otherwise healthy and appears well. Cardiac examination shows a regular rate and rhythm with wide splitting of S2. The lungs are clear to auscultation. This patient is most likely to have a systolic murmur best auscultated at which of the following labeled locations?



	Answer	Image
A	A	
B	B	
C	C	

	Answer	Image
D	D	 I - Aortic valve II - Pulmonary valve III - Tricuspid valve IV - Mitral valve
E	E	
F	F	
G	G	
H	H	

Hint

This patient's widely split S2 suggests delayed right ventricle emptying (e.g., from pulmonary valve stenosis).

Correct Answer

A - A

Explanation Why

Point A corresponds to the right carotid [artery](#). Auscultatory findings here can include [bruits](#) (from stenosis or external compression) or radiation from the [systolic](#) crescendo-decrescendo [murmur](#) of [aortic stenosis](#). Neither of these lesions would be likely to account for easy fatigability or a wide-[split S2](#) in a pediatric patient.

B - B

Explanation Why

Point B corresponds to the left carotid [artery](#). Auscultatory findings here can include [bruits](#) (from stenosis or external compression) or radiation from the [systolic](#) crescendo-decrescendo [murmur](#) of [aortic stenosis](#). Neither of these lesions would be likely to account for fatigability or a wide-[split S2](#) in a pediatric patient.


C - C

Explanation Why

Point C, which corresponds to the second right [intercostal space](#) at the right sternal border, is the best auscultation site for [aortic stenosis](#), which is associated with a [systolic](#) crescendo-decrescendo [murmur](#). Although anomalies of the [aortic valve](#) (e.g., [bicuspid aortic valve](#)) are among the most common congenital [heart](#) defects, clinically apparent [aortic stenosis](#) would be unlikely in a 5-year-old patient. Moreover, in [aortic stenosis](#), a [paradoxical split](#) of S2 would be expected on [cardiac auscultation](#) rather than a wide [splitting of S2](#).

D - D

Image



Explanation But

Wide [splitting of S2](#) can also be a sign of [right bundle branch block](#), [pulmonary hypertension](#), and [right ventricular](#) volume overload (e.g., due to an [atrial septal defect](#)).

Explanation Why

Point D, which corresponds to the second left [intercostal space](#) at the parasternal border, is the best auscultation site for [pulmonary stenosis](#). This 5-year-old boy presents with episodic [dyspnea](#), easy fatigability, and a widely [split S2](#), which should raise concern for a congenital valvular issue. Clear [lung auscultation](#) indicates an anomaly that is likely right-sided. [Pulmonary stenosis](#) causes an outflow obstruction from the [right ventricle](#), which will cause delayed closing of the pulmonic valve ([split S2](#)) and a [systolic ejection murmur](#). Flow [murmurs](#) and the [murmur](#) of ASD can also be auscultated at this site.

E - E

Explanation Why

Point E corresponds to the right [axilla](#). Unless the patient has dextrocardia, a cardiac [murmur](#) would not be expected at this site.

F - F

Explanation Why

Point F corresponds to the right 5th [intercostal space](#) at the parasternal line. Unless the patient has dextrocardia, a cardiac [murmur](#) would not be expected at this site.

G - G

Explanation Why

Point G, which corresponds to the fifth left [intercostal space](#) at the [midclavicular line](#), is the best auscultation site for [murmurs](#) originating from the [mitral valve](#). A [diastolic murmur](#) would be expected in [mitral stenosis](#), a [systolic murmur](#) in [mitral valve prolapse](#), and a [holosystolic murmur](#) in [mitral regurgitation](#). Severe [mitral regurgitation](#) can cause wide [splitting of S2](#), which is seen in this patient. However, severe [mitral regurgitation](#) would classically manifest with signs of [left-sided heart failure](#) (e.g., [pulmonary edema](#), [dyspnea](#), [cough](#)), which this patient does not have.

H - H

Explanation Why

Point H corresponds to the left [axilla](#). The high-pitched, blowing, [holosystolic murmur](#) of [mitral regurgitation](#) can be auscultated at this site. Severe [mitral regurgitation](#) can cause wide [splitting of S2](#), which is seen in this patient. However, severe [mitral regurgitation](#) would classically manifest with signs of [left-sided heart failure](#) (e.g., [pulmonary edema](#), [dyspnea](#), [cough](#)), which this patient does not have.

Question # 5

A 52-year-old man comes to the physician because of a 3-day history of intermittent chest tightness that worsens with exercise. He has chronic atrial fibrillation treated with a drug that prolongs the QT interval. During cardiac stress testing, an ECG shows progressive shortening of the QT interval as the heart rate increases. Which of the following drugs is this patient most likely taking?

	Answer	Image
A	Diltiazem	
B	Lidocaine	
C	Flecainide	
D	Dofetilide	
E	Quinidine	
F	Carvedilol	

Hint

This patient is taking a drug that exhibits reverse use-dependence, a phenomenon in which the effects of an antiarrhythmic (e.g., QT interval prolongation for this drug) decrease as the heart rate increases.

Correct Answer

A - Diltiazem

Explanation Why

[Diltiazem](#) is a [class IV antiarrhythmic](#) that is used for rate control in patients with [atrial fibrillation](#). [Class IV antiarrhythmic drugs](#) inhibit slow calcium channels which mediate cardiac pacemaker [depolarization](#). This inhibition decreases the ability of [cardiac pacemaker cells](#) to reach the [threshold potential](#) and decreases conduction velocity. As a result, the duration of the [effective refractory period](#) increases, which results in [PR interval](#) prolongation on [ECG](#). [Class IV antiarrhythmics](#) do not affect the [QT interval](#).

B - Lidocaine

Explanation Why

[Lidocaine](#) is a [class IB antiarrhythmic](#) that is used to treat [ventricular arrhythmias](#); it is not used to treat [atrial fibrillation](#). [Class IB antiarrhythmics](#) block sodium channels in cardiac [myocytes](#) and reduce the duration of the [action potential](#). They do not affect the [QT interval](#). Moreover, while [class IB antiarrhythmics](#) exhibit use-dependence in very rare cases, they do not exhibit [reverse use-dependence](#).

C - Flecainide

Explanation Why

[Flecainide](#) is a [class IC antiarrhythmic](#) that is used to convert [atrial fibrillation](#) to [sinus rhythm](#). [Class IC antiarrhythmics](#) block sodium channels in cardiac [myocytes](#) and prolong the duration of the [effective refractory period](#) in the [atrioventricular node](#). Although this class of [antiarrhythmics](#) has little effect on ventricular [repolarization](#), the [QT interval](#) is minimally prolonged due to the increase in the QRS duration. Moreover, [flecainide](#) exhibits use-dependence; in rare cases, there is a progressive prolonging of the [QT interval](#) on [cardiac stress testing](#) (i.e., as the [heart rate](#) increases) rather than the shortening of the [QT interval](#) seen in this patient.

D - Dofetilide

Explanation Why

[Dofetilide](#) is a [class III antiarrhythmic](#) that is used to maintain normal [sinus rhythm](#) in patients with chronic [atrial fibrillation](#). [Class III antiarrhythmics](#) inhibit potassium channels which are responsible for [repolarization](#) of the cardiac membrane. Delayed [repolarization](#) prolongs the duration of the [cardiac action potential](#) and the [effective refractory period](#), which results in [QT interval prolongation](#) on [ECG](#). All [class III antiarrhythmics](#), with the exception of [amiodarone](#), exhibit [reverse use-dependence](#), as seen during this patient's [cardiac stress testing](#). Accordingly, these drugs have decreased antiarrhythmic effects at faster [heart rates](#) and are associated with the development of [torsades de pointes](#) at slower [heart rates](#).

E - Quinidine

Explanation Why

[Quinidine](#) is a [class IA antiarrhythmic](#) that is used to maintain [sinus rhythm](#) in patients with [atrial fibrillation](#). [Class IA antiarrhythmics](#) decrease sodium influx during [depolarization](#) and potassium efflux during [repolarization](#). This activity prolongs the duration of the [cardiac action potential](#) and the [effective refractory period](#), which results in [QT interval prolongation](#) on [ECG](#). However, [class IA antiarrhythmics](#) exhibit use-dependence and thus cause progressive prolongation, not shortening, of the [QT interval](#) as the [heart rate](#) increases.

F - Carvedilol

Explanation Why

[Carvedilol](#) is a [β-blocker](#) and [class II antiarrhythmic](#) that is used for rate control in patients with [atrial fibrillation](#). In the cardiac [myocyte](#), [β-adrenergic](#) activation of [adenylate cyclase](#) results in the production of [cAMP](#). This [second messenger](#) facilitates the release of calcium from the [sarcoplasmic reticulum](#), which leads to cardiac pacemaker [depolarization](#). [Class II antiarrhythmics](#) decrease the conduction velocity of [cardiac pacemaker cells](#), which results in [PR interval](#) prolongation on [ECG](#).

They do not affect the [QT interval](#).

Question # 6

A 62-year-old man comes to the physician for a 1-month history of fever, malaise, and skin rash. He has had a 5-kg (11-lb) weight loss during this period. He does not smoke, drink alcohol, or use illicit drugs. He appears pale. His temperature is 39.1°C (102.3°F), pulse is 110/min, and blood pressure is 140/85 mm Hg. Physical examination shows nontender, erythematous macules on the palms and soles. A photograph of one of his fingernails is shown. Microscopic examination of the nail lesion is most likely to show which of the following?



	Answer	Image
A	IgE immune complexes	
B	Aschoff granulomas	
C	Arteriovenous malformations	

	Answer	Image
D	Microemboli	
E	Non-caseating granulomas	

Hint

This patient's symptoms of fever, malaise, and weight loss along with his exam findings of Janeway lesions (nontender macules on palms and soles) and splinter hemorrhages (photograph) make infectious endocarditis the most likely diagnosis.

Correct Answer

A - IgE immune complexes

Explanation Why

There is growing evidence to suggest that deposition of [immune complexes](#) within the [capillary](#) beds is a cause of [vasculitis](#) leading to [splinter hemorrhages](#), [roth spots](#), and/or glomerulonephritis. However, the offending [immune complexes](#) in [endocarditis](#) are predominantly formed by [IgA](#) and/or [IgG](#), not [IgE](#).

B - Aschoff granulomas

Explanation Why

Aschoff [granulomas](#) ([Aschoff bodies](#)) form due to [inflammation](#) caused by [rheumatic fever](#), which can also manifest with [fever](#), malaise, and [skin](#) findings (subcutaneous nodules and [erythema marginatum](#)). However, [Aschoff bodies](#) would be found in the patient's [myocardium](#) rather than his fingernails. Moreover, [rheumatic fever](#) mostly occurs within the first two decades of life and would not be expected in this 62-year-old patient. This patient also does not satisfy [Jones criteria](#).

C - Arteriovenous malformations

Explanation Why

[Arteriovenous malformations](#) ([AVMs](#)) occur as congenital connections between [arteries](#) and [veins](#). These defects are associated with [ischemia](#) and hemorrhages. Unlike this patient's presentation, [AVMs](#) do not cause systemic signs such as [fever](#), malaise, and weight loss, and are not associated with forming [Janeway lesions](#).

D - Microemboli

Image



Explanation Why

Microemboli trapped within the [capillary](#) networks of [nail](#) beds can cause [splinter hemorrhages](#) like the ones seen in this patient. These microemboli break off from infected left-[heart](#) cardiac valves and enter [systemic circulation](#), where they become trapped in [capillary](#) beds throughout the body. In addition, there is growing evidence to suggest that [vasculitis](#) due to deposition of [immune complexes](#) within the [capillary](#) beds contributes to the formation of [splinter hemorrhages](#), [roth spots](#), and glomerulonephritis, all of which can occur in patients with [infectious endocarditis](#). [Janeway lesions](#) are caused by microabscesses and neutrophilic [capillary](#) infiltration resulting from [septic microemboli](#).

E - Non-caseating granulomas

Explanation Why

The presence of [non-caseating granulomas](#) within the lesion would signify the occurrence of [granulomatous inflammation](#). A granulomatous disease associated with both systemic ([fever](#), malaise, weight loss) and dermatological ([splinter hemorrhages](#), [blistering](#) rashes) manifestations similar to those seen in this patient is [granulomatosis with polyangiitis](#). However, this patient does not exhibit the most common features of [granulomatosis with polyangiitis](#) (i.e., chronic [rhinitis/sinusitis](#); lower respiratory tract symptoms such as [cough](#), [dyspnea](#), and [hemoptysis](#); and [RPGN](#)).

Question # 7

A 38-year-old woman, gravida 2, para 1, at 24 weeks' gestation comes to the physician for a routine prenatal evaluation. She has no history of major medical illness and takes no medications. Fetal ultrasonography shows a cardiac defect resulting from abnormal development of the endocardial cushions. This defect is most likely to result in which of the following?

	Answer	Image
A	Tetralogy of Fallot	
B	Atrioventricular septal defect	
C	Sinus venosus defect	
D	Transposition of the great vessels	
E	Dextrocardia	
F	Patent foramen ovale	

Hint

Down syndrome is associated with endocardial cushion defects.

Correct Answer

A - Tetralogy of Fallot

Explanation Why

[Tetralogy of Fallot](#) is characterized by [right ventricular outflow tract obstruction](#) due to [pulmonary valve stenosis](#), [right ventricular hypertrophy](#), a [ventricular septal defect](#), and an overriding aorta above the [ventricular septal defect](#). The underlying pathology is thought to be a malpositioned infundibular septum. Abnormal development of the [endocardial cushions](#) does not cause [tetralogy of Fallot](#).

B - Atrioventricular septal defect

Explanation Why

In normal cardiac development, the superior and inferior [endocardial cushions](#) fuse to form the atrioventricular canal and parts of the [atrioventricular valves](#). Abnormal development of the [endocardial cushions](#) causes [atrioventricular septal defects \(AVSDs\)](#), the most common of which is an [ostium secundum atrial septal defect](#). In mild cases of [AVSD](#), patients can have septal defects with separate [atria](#) and ventricles. In severe cases, patients can have a common atrioventricular canal (i.e., combined [atrial septal defect](#) and [ventricular septal defect](#)) with a single [atrioventricular valve](#).

C - Sinus venosus defect

Explanation Why

[Sinus venosus](#) defects result from abnormal resorption of the [sinus venosus](#) during cardiac development. This causes malpositioning of the [superior vena cava](#) or [inferior vena cava](#) in relation to the [right atrium](#) and is associated with the development of abnormal connections between the [pulmonary veins](#) and the [right atrium](#). Abnormal development of the [endocardial cushions](#) does not cause [sinus venosus](#) defects.

D - Transposition of the great vessels

Explanation Why

[Transposition of the great vessels](#) results from failure of the [aortopulmonary septum](#) to undergo a spiral transformation. The aorta and [pulmonary trunk](#) are attached to the right and left ventricles, respectively, prior to spiral transformation of the septum. Following transformation, the aorta is attached to the [left ventricle](#), and the [pulmonary trunk](#) is attached to the [right ventricle](#). If this process fails, the [pulmonary circulation](#) and [systemic circulation](#) function as completely separate parallel circuits, and oxygenated blood cannot pass into the [systemic circulation](#). To sustain life, oxygenated and deoxygenated blood must mix through associated cardiac defects (e.g., [atrial septal defects](#), [ventricular septal defects](#), [patent ductus arteriosus](#)). Abnormal development of the [endocardial cushions](#) does not cause [transposition of the great vessels](#).

E - Dextrocardia

Explanation Why

Dextrocardia is an abnormal orientation of the [heart](#) in which the apex is located to the right aspect of the [mediastinum](#) rather than the left. It is often associated with [situs inversus](#) and is caused by a defect in [dynein](#) (e.g., from [Kartagener syndrome](#)). Abnormal development of the [endocardial cushions](#) does not cause dextrocardia.


F - Patent foramen ovale

Explanation Why


A [patent foramen ovale](#) results from failure of fusion of the [septum primum](#) and [septum secundum](#), which typically occurs after [birth](#). Abnormal development of the [endocardial cushions](#) does not cause a [patent foramen ovale](#).




Question # 8



A 58-year-old man comes to the physician because of a 5-day history of progressively worsening shortness of breath and fatigue. He has smoked 1 pack of cigarettes daily for 30 years. His pulse is 96/min, respirations are 26/min, and blood pressure is 100/60 mm Hg. An x-ray of the chest is shown. Which of the following is the most likely cause of this patient's findings?



	Answer	Image
A	Left ventricular failure	

	Answer	Image
B	Tricuspid regurgitation	
C	Spontaneous pneumothorax	
D	Acute respiratory distress syndrome	

	Answer	Image
E	Pulmonary embolism	 A frontal chest X-ray showing a wedge-shaped consolidation in the right lower lung field, which is a classic sign of pulmonary embolism. The consolidation is located in the lower lung field, and the heart size appears normal. The right and left sides are labeled 'R' and 'L' respectively.
F	Pulmonary emphysema	 A frontal chest X-ray showing hyperinflation of the lungs, flattened diaphragms, and increased retrosternal space, which are characteristic findings of pulmonary emphysema. The heart size appears normal.
G	Interstitial pneumonia	 Two chest X-rays showing bilateral interstitial infiltrates, suggestive of interstitial pneumonia. Image A is a frontal view, and Image B is a lateral view. The infiltrates are visible as increased lung markings throughout both lung fields. The lateral view (B) shows the infiltrates extending to the periphery of the lung. The right and left sides are labeled 'R' and 'L' respectively, and 'Upright' is noted on the lateral view.

	Answer	Image
H	Bronchial asthma	 A frontal chest X-ray showing hyperinflation of the lungs, which is a characteristic finding in bronchial asthma. The diaphragms are flattened, and the lung fields appear more radiolucent than normal.
I	Pulmonary fibrosis	 A frontal chest X-ray showing bilateral interstitial markings and reticular opacities, which are characteristic findings in pulmonary fibrosis. The markings are most prominent in the lower lung zones.

Hint

This patient's x-ray shows butterfly-shaped perihilar opacification, Kerley B lines, cardiomegaly, and blunting of the costophrenic angle.

Correct Answer

A - Left ventricular failure

Explanation But


Noncardiogenic pulmonary edema due to [ARDS](#) or fluid overload (e.g., aggressive [fluid therapy](#), [oliguric renal failure](#)) can manifest with radiological findings similar to the ones seen here. However, this patient's [cardiomegaly](#) suggests a chronic underlying [heart](#) condition and a cardiac cause for the [pulmonary edema](#).

Explanation Why

This patient's acute onset [dyspnea](#) and [x-ray](#) findings suggest cardiogenic pulmonary edema due to [left ventricular failure](#). Individuals with [left ventricular failure](#) initially develop [interstitial edema](#), which results in smooth reticular opacification and [Kerley B lines](#) due to accentuation of the vascular markings. With the further increase in [pulmonary capillary wedge pressure](#), individuals can develop features of [alveolar edema](#) (e.g., consolidation with an [air bronchogram](#)) and [pleural effusion](#) (e.g., blunting of the [costophrenic angle](#)). In [pulmonary edema](#), the opacification can often assume a perihilar distribution (a so-called “batwing” or “butterfly-shaped” pattern), which is seen here, because the periphery of the [lung](#) has better [lymphatic drainage](#) than the central regions.

B - Tricuspid regurgitation

Image



Explanation Why

[Tricuspid regurgitation](#) is a rare valvular condition. It typically manifests with clinical features of [right heart failure](#), such as [jugular venous distention](#), pulsatile [hepatomegaly](#), and pedal [edema](#), none of which are seen here. Moreover, the chest [x-ray](#) findings for [tricuspid regurgitation](#) typically include right atrial and [right ventricular](#) enlargement, unlike the [left ventricular](#) enlargement seen on this patient's [x-ray](#).

C - Spontaneous pneumothorax

Image



Explanation Why

[Spontaneous pneumothorax](#) causes acute onset [dyspnea](#), and a chest x-ray would show a [radiolucent](#) region without [lung](#) markings (air collection). This patient's chest [x-ray](#) findings are inconsistent with a [pneumothorax](#).

D - Acute respiratory distress syndrome

Image




Explanation Why

[Acute respiratory distress syndrome](#) can cause [dyspnea](#), and a chest x-ray would show butterfly-shaped perihilar opacification, which is seen here. However, this patient's [cardiomegaly](#) suggests a chronic underlying cardiovascular condition and, thus, a cardiac cause for the [pulmonary edema](#).

E - Pulmonary embolism

Image




Explanation Why

[Pulmonary embolism](#) can cause acute onset [dyspnea](#) as well as blunting of the [costophrenic angle](#) due to a [pleural effusion](#) caused by [pulmonary infarction](#). [Kerley B lines](#) and butterfly-shaped perihilar opacities would not, however, be seen in [pulmonary embolism](#). The chest [x-ray](#) findings associated with [pulmonary embolism](#) are relatively nonsensitive and nonspecific and include wedge-shaped pulmonary opacification due to [pulmonary infarction](#), a regional decrease in pulmonary vascular markings, and enlargement of the [pulmonary artery](#), none of which are seen here.

F - Pulmonary emphysema

Image




Explanation Why

Smoking is an important [risk factor](#) for pulmonary [emphysema](#), which causes [dyspnea](#) and fatigue. However, the symptoms of [emphysema](#) typically evolve over months and years rather than a few days. Moreover, the characteristic [x-ray](#) findings for [emphysema](#) are increased [lung](#) lucency and flattening of the diaphragm due to hyperinflation. This patient's [x-ray](#) findings are inconsistent with pulmonary [emphysema](#).

G - Interstitial pneumonia

Image




Explanation Why

Certain infectious causes of [interstitial pneumonia](#) (e.g., [Mycoplasma pneumoniae](#), [Pneumocystis pneumonia](#)) lead to acute onset [dyspnea](#), fatigue, and perihilar [interstitial](#) infiltrates. However, blunting of the [costophrenic angle](#) (due to [pleural effusion](#)) is uncommon in both [Mycoplasma](#) and [Pneumocystis pneumonia](#), and features such as [cardiomegaly](#) and [Kerley B lines](#) would not be expected in [interstitial pneumonia](#).

H - Bronchial asthma

Image




Explanation Why

An acute exacerbation of [bronchial asthma](#) can cause acute onset [dyspnea](#), and the [x-ray](#) would be normal or show evidence of hyperinflation (e.g., increased [lung](#) lucency, flattening of the diaphragm). This patient's [x-ray](#) findings are inconsistent with [bronchial asthma](#).

I - Pulmonary fibrosis

Image



Explanation Why

Smoking is an important [risk factor](#) for pulmonary [fibrosis](#), which can cause [dyspnea](#) and reticular opacification. However, the symptoms of pulmonary [fibrosis](#) typically evolve over months and years rather than a few days. In addition, the reticular opacification of pulmonary [fibrosis](#) manifests with [honeycombing](#) of the [lung](#), and it typically affects the base of the [lungs](#), unlike the butterfly-shaped perihilar reticular opacities seen here. Moreover, the [x-ray](#) findings of [Kerley B lines](#) and blunting of the [costophrenic angle](#) are inconsistent with pulmonary [fibrosis](#).

Question # 9

A 32-year-old woman comes to the physician for a screening health examination that is required for scuba diving certification. The physician asks her to perform a breathing technique: following deep inspiration, she is instructed to forcefully exhale against a closed airway and contract her abdominal muscles while different cardiovascular parameters are evaluated. Which of the following effects is most likely after 10 seconds in this position?

	Answer	Image
A	Increased venous return to right atrium	
B	Increased venous return to left atrium	
C	Decreased systemic vascular resistance	
D	Decreased left ventricular stroke volume	
E	Decreased intra-abdominal pressure	
F	Decreased pulse rate	

Hint

This patient is performing the Valsalva maneuver.

Correct Answer

A - Increased venous return to right atrium

Explanation Why

The [Valsalva maneuver](#) increases intrathoracic pressure, as the respiratory muscles contract against a closed [glottis](#). [Venous return](#) varies inversely with intrathoracic pressure. It would therefore be decreased in this patient, not increased.

B - Increased venous return to left atrium

Explanation Why

While there is a transient increase in [venous return](#) immediately on initiation of phase 1 (as the blood is squeezed out of the [lungs](#) and into the [left atrium](#)), [venous return](#) quickly decreases in the remainder of phase 1 and 2 of the [Valsalva maneuver](#) due to the increased intrathoracic pressure. In this patient, who has maintained the [Valsalva maneuver](#) for 10 seconds, [venous return](#) to the [left atrium](#) would rather be decreased.

C - Decreased systemic vascular resistance

Explanation Why

[Systemic vascular resistance](#) increases during the [Valsalva maneuver](#), especially during phase 2. The reduced [cardiac output](#) seen in phase 2 stimulates [carotid sinus baroreceptors](#), which reflexively increases the [heart rate](#) and promotes peripheral vasoconstriction.

D - Decreased left ventricular stroke volume

Explanation Why

A sustained [Valsalva maneuver](#) reduces the [left ventricular stroke volume](#). When the [Valsalva maneuver](#) is initiated, four phases are typically seen. Phase 1 (onset of straining) and phase 2 (continuation of straining) describe the changes that occur during active expiration against a closed [glottis](#), while phase 3 (release of straining) and phase 4 (recovery phase) follow the release of pressure. The increased intrathoracic pressure seen in phase 2 reduces total [venous return](#), and increases [systemic vascular resistance](#), both of which reduce [stroke volume](#). Once pressure is released, i.e., in phases 3 and 4, the sudden release of pressure would increase [stroke volume](#).

E - Decreased intra-abdominal pressure

Explanation Why

Because the abdominal and respiratory muscles contract forcefully during the [Valsalva maneuver](#), both the intra-abdominal and intrathoracic pressure would increase, not decrease.


F - Decreased pulse rate

Explanation Why

During the first seconds (phase 1) of the [Valsalva maneuver](#), increased intrathoracic pressure causes a shift of blood from the [pulmonary circulation](#) into the left [heart](#). The resulting transient increase in [cardiac output](#) and blood pressure stimulates aortic [baroreceptors](#), which decreases [pulse](#) rate during the initial phase. Continuation of the [Valsalva maneuver](#) (phase 2), as described here, results in a decreased arterial blood pressure (due to reduced [venous return](#)). The drop in blood pressure induces [reflex tachycardia](#), which is reflected in an increased (not decreased) [pulse](#) rate during phase 2.

Question # 10

A 32-year-old woman who recently emigrated to the USA from Japan comes to the physician because of a 3-month history of night sweats, malaise, and joint pain. During this time, she has also had a 6-kg (13-lb) weight loss. Physical examination shows weak brachial and radial pulses. There are tender subcutaneous nodules on both legs. Carotid bruits are heard on auscultation bilaterally. Laboratory studies show an erythrocyte sedimentation rate of 96 mm/h. A CT scan of the chest shows thickening and narrowing of the aortic arch. Microscopic examination of the aortic arch is most likely to show which of the following findings?

	Answer	Image
A	Fibrinoid necrosis of the intima and media	
B	Granulomatous inflammation of the media	
C	Subendothelial hyaline deposition	
D	Subendothelial immune complex deposition	

	Answer	Image
E	Calcification of the media	


Hint

This patient's findings of weak peripheral pulses, carotid bruits, elevated ESR, and thickening of the aortic arch should raise concern for Takayasu arteritis. This type of large vessel vasculitis predominantly occurs in women under the age of 40 years (especially those with Asian ancestry).

Correct Answer

A - Fibrinoid necrosis of the intima and media

Image



Explanation Why

Fibrinoid necrosis of the tunica intima and tunica media are characteristic of polyarteritis nodosa (PAN), which can manifest with fever, night sweats, malaise, arthralgia, and erythema nodosum. However, PAN involves small- to medium-sized arteries (e.g., renal arteries, vasa nervorum, arteries supplying the intestines, and cutaneous vessels) rather than large vessels, which are affected in this patient.

B - Granulomatous inflammation of the media

Image



Explanation But

Patients with [Takayasu arteritis](#) should be treated with [corticosteroids](#).

Explanation Why

The hallmark of [Takayasu arteritis](#) (TA) is [granulomatous inflammation](#) of the arterial wall affecting the large [arteries](#) (typically the aorta and its branches). TA is more common among women with Asian ancestry, and its peak [incidence](#) is between 15 and 45 years of age. This condition manifests with systemic features that include [fever](#), malaise, and arthralgia. Other features of TA include decreased brachial and radial pulses ([pulseless disease](#)), [syncope](#), [carotid bruit](#), [angina pectoris](#), impaired vision, and [Raynaud phenomenon](#), all of which are due to narrowing of the large [arteries](#). [Erythema nodosum](#) and [urticaria](#) may also be seen in TA. Microscopic examination of the diseased vessels would show [granulomatous inflammation](#) of the vessel walls, with inflammatory cells infiltrating the [tunica media](#) and [tunica adventitia](#).

C - Subendothelial hyaline deposition

Explanation Why

Subendothelial [hyaline](#) deposition is a form of [arteriosclerosis](#) commonly seen in individuals with chronic [essential hypertension](#) and/or [diabetes](#). It can also be part of the normal [aging](#) process. However, thickening of the [aortic arch](#), [erythema nodosum](#), and the systemic symptoms of night sweats, malaise, [joint pain](#), and weight loss are not associated with [hyaline](#) deposits. Furthermore, this patient is 32 years of age and has none of the [risk factors](#) for [hyaline](#) deposition.

D - Subendothelial immune complex deposition

Explanation Why

[Immune complex](#) deposition is characteristic of [Henoch-Schonlein purpura](#) (HSP), which is a [small vessel vasculitis](#) that can manifest with malaise, arthralgia, and cutaneous lesions. However, the cutaneous lesions of HSP are nontender palpable [purpura](#) rather than tender nodules ([erythema nodosum](#)). Moreover, decreased peripheral pulses and involvement of large vessels (e.g., aorta, [brachial arteries](#)) would not be expected. Instead, HSP manifests with other features, such as [nephritic syndrome](#) (due to [IgA nephropathy](#)) and abdominal [pain](#).

E - Calcification of the media

Explanation Why

[Mönckeberg arteriosclerosis](#) is characterized by calcification of the [tunica media](#), and it is associated with [diabetes mellitus](#). This condition usually affects the smaller vessels in the extremities and is not associated with stenosis. Furthermore, thickening of the [aortic arch](#), [erythema nodosum](#), and the systemic symptoms of night sweats, malaise, [joint pain](#), and weight loss are not associated with [Mönckeberg arteriosclerosis](#).

Question # 11

A 38-year-old man comes to the physician because of a 3-week history of a painful rash affecting his left foot. For the past 2 years, he has had recurrent episodes of color changes in his fingers when exposed to the cold; his fingers first turn white and then progress to blue and red before spontaneously resolving. He has smoked two packs of cigarettes daily for 20 years. His blood pressure is 115/78 mm Hg. Physical examination shows multiple tender, dark purple nodules on the lateral surface of the left foot with surrounding erythema that follow the course of the lateral marginal vein. There are dry ulcers on the tip of his right index finger and on the distal aspect of his right hallux. Serum lipid studies show no abnormalities. Biopsy of the dorsalis pedis artery will most likely show which of the following findings?

	Answer	Image
A	Granulomatous inflammation with narrowing of the vessel lumen	
B	Segmental thrombosing inflammation with sparing of the internal elastic lamina	
C	Calcification of the tunica media with foam cells and fibrous cap formation	
D	Intraluminal fibrin clot predominantly composed of red blood cells	
E	Transmural inflammation with fibrinoid necrosis of the vessel wall	

Hint

This patient's signs of digit ulceration, superficial thrombophlebitis (painful nodules), and Raynaud phenomenon (vasospasm of the fingers in response to cold exposure), together with his significant smoking history, are suggestive of thromboangiitis obliterans (TAO).

Correct Answer

A - Granulomatous inflammation with narrowing of the vessel lumen

Explanation Why

[Granulomatous inflammation](#) with narrowing of the vessel lumen is a distinguishing feature of [Takayasu arteritis](#). This medium-to-large vessel arteritis is associated with Raynaud phenomenon, which is seen in this patient. However, [Takayasu arteritis](#) most commonly affects middle-aged women of Asian descent and manifests with [fevers](#), polyarthralgias, visual impairments, and [hypertension](#), none of which are seen in this patient.

B -

Segmental thrombosing inflammation with sparing of the internal elastic lamina

Image



Explanation But

Smoking is the single most important [risk factor](#) for TAO. The exact mechanism by which smoking causes TAO is unknown, but direct [endothelial](#) cell toxicity from the components of tobacco is presumed to play a role. Complete cessation of smoking is, therefore, the single most important therapeutic measure to prevent [gangrene](#) and [amputation](#).

Explanation Why

Segmental thrombosing [inflammation](#) of small and medium-sized vessels that spares the [internal elastic lamina](#) is the characteristic histologic finding in [thromboangiitis obliterans](#). TAO most commonly occurs in male smokers < 45 years of age and manifests with [migratory thrombophlebitis](#), [Raynaud phenomenon](#), and digital [ischemia](#) with [intermittent claudication](#). Eventually, [critical limb ischemia](#) may develop, which can result in [rest pain](#) of the [distal](#) extremities, peripheral pulselessness, and/or [gangrene](#) with autoamputation of digits.

C - Calcification of the tunica media with foam cells and fibrous cap formation

Explanation Why

Calcification of the [tunica media](#) with [foam cell](#) infiltration and fibrous cap formation is a distinguishing feature of [atherosclerosis](#). [Atherosclerosis](#) is most commonly seen in elderly men with a significant smoking history. In addition, luminal narrowing due to formation of [atherosclerotic plaques](#) is the pathophysiology underlying [peripheral artery disease](#), which can cause digit ulcers, as seen in this patient. However, [atherosclerosis](#) would not explain this patient's [Raynaud phenomenon](#) and [superficial thrombophlebitis](#). Moreover, [atherosclerosis](#) is strongly associated with [arterial hypertension](#) and [dyslipidemia](#), none of which are seen here.

D - Intraluminal fibrin clot predominantly composed of red blood cells

Explanation Why

Intraluminal [fibrin](#) clots that are predominantly composed of [red blood cells](#) are known as [red thrombi](#), which are commonly seen in [deep venous thrombosis](#). Patients with [DVT](#) often report unilateral leg [erythema](#) and swelling with notable tenderness that is worse upon palpation. [DVT](#) does not explain this patient's [Raynaud phenomenon](#), digital ulcers, or the waxing and waning timecourse of his symptoms.


E - Transmural inflammation with fibrinoid necrosis of the vessel wall

Explanation Why

Transmural [inflammation](#) with [fibrinoid necrosis](#) is a distinguishing histologic feature of [polyarteritis nodosa](#). [Skin](#) ulcerations can occur due to obstruction of medium-sized [arteries](#) (e.g., digital [arteries](#)), ultimately causing [ischemia](#) in the fingertips. However, patients with [PAN](#) also present with [hypertension](#) (due to renal involvement), [polyneuropathy](#), and signs of gastrointestinal distress (e.g., nausea, vomiting), none of which are seen in this patient. Moreover, [PAN](#) is not commonly associated with [Raynaud phenomenon](#).

Question # 12

A 38-year-old woman comes to the physician because of a 1-month history of progressively worsening dyspnea, cough, and hoarseness of voice. Her pulse is 92/min and irregularly irregular, respirations are 20/min, and blood pressure is 110/75 mm Hg. Cardiac examination shows a rumbling mid-diastolic murmur that is best heard at the apex in the left lateral decubitus position. Which of the following is the most likely underlying cause of this patient's condition?

	Answer	Image
A	Annular calcification	
B	Myxomatous degeneration	
C	Hematogenous spread of bacteria	
D	Antibody cross-reactivity	
E	Sarcomeric gene mutation	
F	Congenital valvular defect	

Hint

The rumbling mid-diastolic murmur at the apex is characteristic of mitral stenosis.

Correct Answer

A - Annular calcification

Explanation Why

Mitral annular calcification, when severe, can cause [mitral stenosis](#) as a result of impaired [diastolic](#) dilation and restriction of valve motion. However, this condition typically affects older adults (> 50 years of age) and is usually asymptomatic; it is not a likely cause of this patient's condition.

B - Myxomatous degeneration

Explanation Why

Myxomatous degeneration with pathological weakening of valvular [connective tissue](#) results in [mitral valve prolapse \(MVP\)](#), which can manifest with features of [heart failure](#) (e.g., [dyspnea](#), [cough](#)) and a [murmur](#) that is best heard at the [mitral area](#) (cardiac apex). [Left atrial enlargement](#) due to chronic [MVP](#) can cause [atrial fibrillation](#) and hoarseness of voice due to [recurrent laryngeal nerve](#) compression. However, unlike this patient's [diastolic murmur](#), [MVP](#) is characterized by a late [systolic](#) crescendo [murmur](#) with a mid-[systolic](#) click due to sudden tensing of the chordae tendinae.


C - Hematogenous spread of bacteria

Explanation Why

[Infective endocarditis](#) can cause a [murmur](#) at the cardiac apex due to [mitral valve](#) involvement, and it can also cause clinical features of [heart failure](#) (e.g., [dyspnea](#), [cough](#)). However, the [murmur](#) is caused by [mitral regurgitation](#) and would, therefore, be [systolic](#) rather than [diastolic](#); [infective endocarditis](#) is very rarely associated with [mitral stenosis](#). Moreover, this condition is not associated with hoarseness of voice.

D - Antibody cross-reactivity

Image



Explanation Why

[Rheumatic heart disease](#) is by far the most common cause of [mitral stenosis](#). Following infection by [group A streptococci](#), patients can develop [rheumatic fever](#), in which the [antibodies](#) to the [streptococcal M protein](#) cross-react with similar [proteins](#) on the [heart valves](#). This triggers an inflammatory process that results in valvular [fibrosis](#), scarring, and narrowing. Symptoms of [mitral stenosis](#) include fatigue, exertional dyspnea, [cough](#), and [atrial fibrillation](#) (irregularly irregular [pulse](#)). Hoarseness can develop in severe cases because of [left atrial enlargement](#) and compression of the [recurrent laryngeal nerve](#) (Ortner syndrome).

E - Sarcomeric gene mutation

Explanation Why

[Sarcomeric gene mutations](#) underlie [hypertrophic cardiomyopathy \(HCM\)](#), which can manifest in young adults with features of [heart failure](#) (e.g., [dyspnea](#), [cough](#)) and a [murmur](#) (due to left ventricular outflow tract obstruction). [HCM](#) is also a [risk factor](#) for the development of [atrial fibrillation](#). However, the [murmur](#) associated with [HCM](#) is a [systolic](#) ejection [murmur](#) (crescendo-decrescendo [murmur](#)) that is best heard at the left sternal border. Moreover, [HCM](#) is not associated with hoarseness of voice.

F - Congenital valvular defect

Explanation Why

Congenital valvular defects only very rarely cause [mitral stenosis](#), accounting for approx. 1% of all cases; it is not a likely cause of this patient's condition.

Question # 13

A 67-year-old woman comes to the physician because of fever, chills, myalgias, and joint pain 1 month after undergoing aortic prosthetic valve replacement due to high-grade aortic stenosis. She does not drink alcohol or use illicit drugs. Her temperature is 39.3°C (102.8°F). She appears weak and lethargic. Physical examination shows crackles at both lung bases and a grade 2/6, blowing diastolic murmur over the right sternal border. Laboratory studies show leukocytosis and an elevated erythrocyte sedimentation rate. The causal organism is most likely to have which of the following characteristics?

	Answer	Image
A	Alpha hemolytic, optochin-sensitive diplococci	
B	Novobiocin-sensitive, coagulase-negative cocci	
C	Catalase-negative cocci that grows in 6.5% saline	
D	Beta hemolytic, bacitracin-sensitive cocci	
E	Alpha hemolytic, optochin-resistant cocci	

Hint

In combination with a recent history of aortic valve replacement, this patient's symptoms suggest prosthetic valve endocarditis, the most common cause of which would be *Staphylococcus epidermis* infection.

Correct Answer

A - Alpha hemolytic, optochin-sensitive diplococci

Explanation Why

S. pneumoniae is an [alpha hemolytic](#), optochin-sensitive diplococcus that may, in extremely rare cases, cause [endocarditis](#). This patient more likely has [prosthetic valve endocarditis](#) from *S. epidermidis*, which is a [gamma hemolytic](#) coccus. Furthermore, [pneumococcal pneumonia](#) typically causes localized [lung](#) consolidation, which manifests in [lung crackles](#) in distinct regions, rather than the bibasilar [crackles](#) heard in this patient.

B - Novobiocin-sensitive, coagulase-negative cocci

Explanation But

Staphylococcus aureus is another common causal pathogen of early [prosthetic valve endocarditis](#) (i.e., within 60 days after implantation).

Explanation Why

S. epidermidis is a novobiocin sensitive, coagulase-negative, catalase-positive, [gram-positive coccus](#) with the capability of [biofilm](#) formation. As *S. epidermidis* is part of the normal [skin](#) flora, contamination of prosthetic devices (e.g., [heart valves](#) and [hip](#) implants) may occur during surgery.

C - Catalase-negative cocci that grows in 6.5% saline

Explanation Why

Enterococcus is a catalase-negative coccus that grows in 6.5% saline. It can cause [infective endocarditis](#), typically following genitourinary or gastrointestinal procedures. However, *Enterococcus* is only implicated in 8% of cases of [endocarditis](#) following valve replacement surgery. *S. epidermidis*, which is the most likely cause of [endocarditis](#) in this patient, is catalase-positive and

does not grow in hypertonic saline.

D - Beta hemolytic, bacitracin-sensitive cocci

Explanation Why

Streptococcus pyogenes is a [beta hemolytic](#), [bacitracin](#) sensitive bacterium that can cause [acute rheumatic fever](#) (ARF). ARF can manifest with [fevers](#), myalgias, arthralgia, [diastolic murmurs](#) at the right sternal border due to [aortic regurgitation](#), basilar [lung crackles](#) due to cardiogenic pulmonary edema, increased [ESR](#), and [leukocytosis](#). However, ARF is more common in children and [adolescents](#). It rarely occurs in patients over the age of 45 years.

E - Alpha hemolytic, optochin-resistant cocci

Explanation Why

Streptococcus bovis and *Streptococcus mitis* are [viridans streptococci](#) ([alpha hemolytic](#), optochin-resistant [cocci](#)) that may cause [infective endocarditis](#) (IE). However, IE due to [viridans streptococci](#) usually occurs following dental or gastrointestinal procedures. The most likely cause of IE in patients with recent [heart valve](#) transplants is, however, *S. epidermidis* (a [γ-hemolytic](#) coccus).

Question # 14

A 73-year-old man dies 4 months after being diagnosed with advanced adenocarcinoma of the colon. Examination of the heart at autopsy shows vegetations lining the mitral valve margins. The vegetations are loosely attached and can be easily scraped off. Microscopic examination shows the vegetations to be composed of interwoven fibrin strands with mononuclear cells. The mitral valve endothelium is intact. Which of the following is the most likely underlying cause of these autopsy findings?

	Answer	Image
A	Procoagulant release	
B	Dystrophic calcifications	
C	Antibody cross-reactivity	
D	Bacterial colonization	
E	Metastatic infiltration	

Hint

Valvular vegetations interwoven with fibrin strands and mononuclear cells are commonly seen in patients with mucin-producing tumors (e.g., adenocarcinoma of the colon or pancreas) or SLE.

Correct Answer

A - Procoagulant release

Explanation But

[Nonbacterial thrombotic endocarditis](#) is associated with [malignancy](#) and certain autoimmune conditions (e.g., [rheumatoid arthritis](#), [SLE](#)), and most commonly affects the aortic and mitral valves. Patients are usually asymptomatic until there is evidence of [thromboembolic phenomena](#).

Explanation Why

Some [adenocarcinomas](#) (e.g., [colon](#), [pancreas](#)) excrete [procoagulant](#) factors (e.g., [tissue factor](#), cancer [procoagulant](#)), resulting in sterile vegetations ([nonbacterial thrombotic endocarditis](#)) composed predominantly of [platelets](#). These [white thrombi](#) are easily dislodged from [heart valves](#), placing patients at high risk of developing tissue infarctions (e.g., [strokes](#), [myocardial infarctions](#)).

B - Dystrophic calcifications

Explanation Why

[Dystrophic](#) calcifications occur in previously damaged tissues with chronic [inflammation](#). The patient's advanced age would predispose him to such [dystrophic calcification](#) as a result of age-related degeneration of the valves. However, there was no damage found on the patient's [endothelium](#) and the vegetations were easily scraped off rather than being firmly attached and showed no hard calcium deposits.

C - Antibody cross-reactivity

Explanation Why

[Antibody cross-reactivity](#) to highly antigenic [M protein \(virulence factor\)](#) of [Streptococcus pyogenes](#) occurs in [rheumatic fever](#). A late sequela of this disease is [rheumatic heart disease](#), in which

vegetations can form on [heart valve](#) surfaces, particularly on the [mitral valve](#). However, such vegetations are inflammatory in nature and do not scrape off easily. Moreover, the [endocardium](#) would show signs of damage with probable histological findings of [Aschoff bodies](#).

D - Bacterial colonization

Explanation Why

Bacterial [colonization](#) of the [heart valves](#) is seen in [infective endocarditis](#) and results in the formation of vegetations of varying size. However, the [septic](#) vegetations in [infective endocarditis](#) are classically firmly attached to the valvular [endothelium](#) and microscopic examination would show dense clusters of bacteria and, possibly, calcification. Moreover, this patient does not show marked [inflammation](#) and destruction of the valvular [endothelium](#), as would be expected in [acute bacterial endocarditis](#).

E - Metastatic infiltration

Explanation Why

[Metastatic](#) infiltration of [heart valves](#) is an infrequent manifestation of cancers, most commonly occurring in [melanoma](#), leukemia, [carcinoma of the lung](#) or [breast](#), and rarely in [colon cancer](#). However, [metastatic heart](#) masses would be infiltrative and destructive, and unlikely to be scraped off easily.

Question # 15

A 50-year-old man comes to the physician because of swelling of his legs for 2 months. Three months ago, he was diagnosed with hypertension and started on a new medication. His blood pressure is 145/95 mm Hg. Physical examination shows 2+ edema in both lower extremities. Laboratory studies are within the reference ranges. This patient was most likely treated with which of the following drugs?

	Answer	Image
A	Losartan	
B	Lisinopril	
C	Spironolactone	
D	Hydrochlorothiazide	
E	Amlodipine	
F	Propranolol	

Hint

The patient was started on an antihypertensive medication and now presents with peripheral edema. This side effect is due to vasodilation, which increases the hydrostatic pressure within the blood vessels, causing an efflux of plasma into the interstitial space.

Correct Answer

A - Losartan

Explanation Why

[Angiotensin II receptor blockers](#) (e.g., [losartan](#)) cause decreased activity of [angiotensin II](#) on the [AT₁ receptor](#), which leads to decreased systemic vasoconstriction as well as decreased renal reabsorption of water, sodium, and [bicarbonate](#). While [ARBs](#) are used in the treatment of [hypertension](#), the most common adverse effects are [angioedema](#) and [hyperkalemia](#), not peripheral [edema](#).

B - Lisinopril

Explanation Why

ACE inhibitors (e.g., [lisinopril](#)) inhibit the conversion of [angiotensin I](#) to [angiotensin II](#) by [ACE](#), thereby decreasing systemic vasoconstriction and renal reabsorption of water and sodium. While ACE inhibitors are used in the treatment of [hypertension](#), common adverse effects include [cough](#), [angioedema](#), and [hyperkalemia](#), but not peripheral [edema](#).

C - Spironolactone

Explanation Why

[Aldosterone receptor antagonists](#) (e.g., [spironolactone](#)) block [aldosterone](#)-mediated sodium and water reabsorption in the [distal convoluted tubule](#). While [aldosterone receptor antagonists](#) are used in the treatment of [hypertension](#), common adverse effects include [hyperkalemia](#) and antiandrogen effects (e.g., [gynecomastia](#)), but not peripheral [edema](#).

D - Hydrochlorothiazide

Explanation Why

[Thiazide diuretics](#) (e.g., [hydrochlorothiazide](#)) inhibit the reabsorption of sodium in the [distal convoluted tubule](#) (which increases the excretion of sodium, water, potassium, and hydrogen ions). While [thiazide diuretics](#) are used in the treatment of [hypertension](#), common adverse effects include [hypokalemia](#), [hyponatremia](#), [hyperglycemia](#), [hyperlipidemia](#), [hyperuricemia](#), and [hypercalcemia](#), but not peripheral [edema](#).

E - Amlodipine

Explanation Why

[Dihydropyridine calcium channel blockers](#) such as [amlodipine](#) inhibit voltage-dependent slow calcium channels. Blockage of these channels results in a decrease in the intracellular calcium that is needed for [electromechanical coupling](#) and muscle contraction. This causes vascular [smooth muscle](#) relaxation, which leads to systemic [vasodilation](#) and a decrease in blood pressure. Other common side effects of [CCBs](#) include [headaches](#), [dizziness](#), facial flushing, and [reflex tachycardia](#).

F - Propranolol

Explanation Why

[Nonselective beta blockers](#) (e.g., [propranolol](#)) inhibit both [beta-1 receptors](#) and [beta-2 receptors](#). While [beta blockers](#) are frequently used in the treatment of [hypertension](#), common adverse effects of [nonselective beta blockers](#) include [bradycardia](#), [orthostatic hypotension](#), bronchoconstriction, and [erectile dysfunction](#). [Propranolol](#) would be unlikely to cause peripheral [edema](#) because it decreases the [vasodilatory](#) effect of peripheral [beta-2 receptors](#).

Question # 16

A 59-year-old man is brought to the emergency department by his wife because of fever, chills, night sweats, and generalized fatigue for 2 weeks. His temperature is 39.1°C (102.4°F). He appears ill. Physical examination shows a grade 3/6 mid-diastolic murmur at the left sternal border, and crackles at both lung bases. A transesophageal echocardiography shows a 12 mm vegetation on the aortic valve. Blood cultures show gram-positive, catalase-negative, gamma-hemolytic cocci in chains that are unable to grow in a 6.5% NaCl medium. Which of the following is the most likely predisposing factor for this patient's current condition?

	Answer	Image
A	Periodontal disease	
B	Colon cancer	
C	Congenital heart defect	
D	Valve replacement	
E	Intravenous drug use	
F	Urinary tract infection	

Hint

The patient's blood cultures and echocardiography findings are consistent with aortic regurgitation (diastolic murmur at the left sternal border) due to *Streptococcus gallolyticus* endocarditis.

Correct Answer

A - Periodontal disease

Explanation Why

Periodontal disease is a [risk factor](#) for IE caused by [viridans group streptococci](#) (e.g., *S. mutans*). However, [Streptococci](#) of the viridans group are [alpha-hemolytic](#) organisms, unlike the [gamma-hemolytic](#) type seen in this patient.

B - Colon cancer

Explanation Why

S. gallolyticus can be found in the normal bowel flora of healthy humans. There is a well-established link between [S. gallolyticus bacteremia](#) (particularly in [endocarditis](#)) and [colorectal cancer](#). However, it remains unclear whether *S. gallolyticus* is part of the etiology of colorectal malignancies or a consequence of the disease. Animal studies suggest that this pathogen may play a role in the pathogenesis of colorectal malignancies. A colonoscopy is indicated in all patients with [S. gallolyticus](#) on blood culture.

C - Congenital heart defect

Explanation Why

Congenital [heart](#) defects (CHDs) are associated with an increased risk of IE. The risks vary based on the underlying defect (e.g., [VSD](#), [tetralogy of Fallot](#), and [PDA](#) pose high risks; ASD and [transposition of the great vessels](#) pose low risks) and the most common etiologic agents are [viridans group streptococci](#) and [Staphylococcus](#) spp. This patient has no evidence of [CHD](#) on his [echocardiogram](#), and [S. gallolyticus](#) is not a typical organism of [CHD](#)-associated IE.

D - Valve replacement

Explanation Why

A history of valve replacement is a [risk factor](#) for IE. IE associated with artificial [heart valves](#) is typically due to [S. epidermidis](#), [S. aureus](#), or gram-negative rods if symptoms occur within 60 days after surgery, or due to [S. aureus](#) or [enterococci](#) if symptoms occur > 60 days after valve replacement. [S. gallolyticus](#), however, is not a typical organism of [prosthetic valve endocarditis](#).

E - Intravenous drug use

Explanation Why

Intravenous drug use is a major [risk factor](#) for [infective endocarditis](#) (IE). However, whereas this patient shows left-sided vegetations seen, IE associated with IV drug use usually affects the right-sided (tricuspid and pulmonary) valves. Also, the common responsible bacteria for [right-sided IE](#) are [Staphylococcus aureus](#) and *Pseudomonas* spp., rather than the [S. gallolyticus](#) seen in this patient.

F - Urinary tract infection

Explanation Why

[Urinary tract infections](#) can cause [bacteremia](#) and subsequent IE in urosepsis, which would manifest with [fever](#), chills, night sweats, and generalized fatigue as well as a [heart murmur](#) and valve vegetations. IE in the setting of urosepsis would typically be associated with gram-negative organisms such as [E. coli](#), *Klebsiella*, and *Enterobacter*, but not [S. gallolyticus](#).

Question # 17

A 41-year-old man with a history of hypertension and hyperlipidemia is brought to the emergency department by his wife for difficulty breathing after choking on food at dinner. He is unconscious and pulseless on arrival. Despite appropriate life-saving measures, he dies. Examination of the heart shows a necrotic, pale yellow plaque in the left circumflex artery. Similar lesions are most likely to be found in which of the following locations?

	Answer	Image
A	Subclavian artery	
B	Superficial temporal artery	
C	Renal artery	
D	Abdominal aorta	
E	Thoracic aorta	
F	Internal carotid artery	

	Answer	Image
G	Pulmonary artery	

Hint

The coronary lesion described on this patient's autopsy is an atherosclerotic plaque. These plaques typically appear in vessels with high levels of endothelial strain.

Correct Answer

A - Subclavian artery

Explanation Why

Occlusion of the [subclavian arteries](#) can occur through thromboses, compression by surrounding structures (e.g., [Pancoast tumor](#)), and certain forms of [vasculitis](#) (e.g., [Takayasu arteritis](#)). While [atherosclerosis](#) can also affect these [arteries](#), they are a less common site for the development of plaques.

B - Superficial temporal artery

Explanation Why

Occlusion of the [superficial temporal artery](#) is classically caused by [giant cell arteritis](#). However, this [artery](#) is less commonly affected by [atherosclerosis](#).


C - Renal artery

Explanation Why

Occlusion of the [renal arteries](#) can occur through [fibromuscular dysplasia](#) and certain forms of [vasculitis](#) (e.g., [polyarteritis nodosa](#)). While [atherosclerosis](#) can affect the [renal arteries](#), they are a less common site for the development of plaques.

D - Abdominal aorta

Image



Explanation Why

The [abdominal aorta](#) is the most common site for the development of [atherosclerotic plaques](#), followed by the coronary, popliteal, and [internal carotid arteries](#). Turbulent, high-pressure flow in these vessels places chronic strain on the [endothelium](#). The [abdominal aorta](#) lacks [vasa vasorum](#) below the level of the [renal arteries](#) and is therefore particularly susceptible to [endothelial](#) injury.

E - Thoracic aorta

Explanation Why

Occlusion of the [thoracic aorta](#) can occur through [aneurysm](#) and/or vessel dissection. While [atherosclerosis](#) can affect this region, the [thoracic aorta](#) is not the most common area for the development of [atherosclerotic plaques](#).

F - Internal carotid artery

Explanation Why

Occlusion of the [internal carotid arteries](#) can occur through vessel dissection or stenosis. While these [arteries](#) can be affected by [atherosclerosis](#), they are less commonly affected than other regions.

G - Pulmonary artery

Explanation Why

Occlusion of the [pulmonary arteries](#) can occur due to emboli (e.g., thrombotic, air, or fat) and certain forms of [vasculitis](#) (e.g., [Takayasu arteritis](#)). However, these [arteries](#) are less commonly affected by [atherosclerosis](#).

Question # 18

A 61-year-old man with hypertension and hyperlipidemia comes to the physician for a 4-month history of recurrent episodes of retrosternal chest pain, shortness of breath, dizziness, and nausea. The episodes usually start after physical activity and subside within minutes of resting. He has smoked one pack of cigarettes daily for 40 years. He is 176 cm (5 ft 9 in) tall and weighs 95 kg (209 lb); BMI is 30 kg/m². His blood pressure is 160/100 mm Hg. Coronary angiography shows an atherosclerotic lesion with stenosis of the left anterior descending artery. Compared to normal healthy coronary arteries, increased levels of platelet-derived growth factor (PDGF) are found in this lesion. Which of the following is the most likely effect of this factor?

	Answer	Image
A	Increased expression of vascular cell-adhesion molecules	
B	Accumulation of foam cells to form fatty streaks	
C	Calcification of the atherosclerotic plaque core	

	Answer	Image
D	Intimal migration of smooth muscle cells	The diagram illustrates the progression of atherosclerosis in a blood vessel over time, categorized into three phases: Asymptomatic, Decades, and Symptomatic, with a final 'Minutes' phase. The vessel wall is shown in cross-section, with the lumen on the left and the vessel wall on the right. The process begins with 'Normal endothelium' and 'Chronic stress (e.g. due to hypertension, inflammation)'. This leads to 'Endothelial dysfunction', followed by 'Infiltration of monocytes and LDL resulting in accumulation of macrophages and oxLDL'. This stage leads to 'Foam cell formation resulting in fatty streaks'. The next stage is 'Proliferation and migration of SMCs and deposition of ECM', which leads to 'Fibrous plaque formation'. This stage leads to 'Plaque destabilization', which leads to 'Plaque rupture (e.g., in context of increased matrix degradation)'. The final stage is 'Thrombus formation'. The diagram includes a legend with the following items: LDL (green circle), Monocyte (purple circle), Macrophage (yellow circle), oxLDL (green circle with 'x'), Foam cell (orange circle), Platelets (red dots), Lymphocytes (blue dots), Cholesterol crystals (blue dots), Smooth muscle cell (SMC) (pink circle), Cellular debris (brown dots), Extracellular matrix (ECM) (yellow dots), PDGF (pink dots), Fibrous cap (pink layer), Necrotic core (yellow layer), and SMC apoptosis, breakdown of the stabilizing ECM (pink layer with arrows).
E	Ingestion of cholesterol by mature monocytes	
F	Invasion of T-cells through the disrupted endothelium	
G	Adherence of platelets to injured subendothelium	
H	Inflammatory cytokine release from the endothelium	

Hint

This patient's stenosis on angiography restricts coronary blood flow and is most likely the cause of his recurrent episodes of angina pectoris (retrosternal chest pain, dyspnea, dizziness, and nausea).

Correct Answer

A - Increased expression of vascular cell-adhesion molecules

Explanation Why

Increased expression of vascular cell-adhesion molecules is an essential process in the pathogenesis of [atherosclerosis](#). In response to [inflammation](#), [endothelial](#) cells increase the expression of [ICAM-1](#) and begin expressing other cell-adhesion molecules such as vascular cell adhesion molecule 1 ([VCAM-1](#)). The increased expression of [VCAM-1](#) leads to adhesion and subsequent migration of [leukocytes](#). However, [PDGF](#) does not increase the expression of vascular cell-adhesion molecules.

B - Accumulation of foam cells to form fatty streaks

Explanation Why

Accumulation of [foam cells](#) in the [intima](#), forming [fatty streaks](#) together with intracellular and extracellular lipid deposits, is one of the first steps in the [pathogenesis of atherosclerosis](#). [PDGF](#) release does not mediate the accumulation of [foam cells](#) to form [fatty streaks](#).


C - Calcification of the atherosclerotic plaque core

Explanation Why

Calcification of the [atherosclerotic plaque](#) core is one of the final steps in the pathogenesis of [atherosclerosis](#). As the [atherosclerotic plaque](#) grows, it causes stenosis of the vessel lumen, which subsequently reduces blood flow and results in [ischemia](#) (manifesting as [angina pectoris](#) in this patient). However, [PDGF](#) does not mediate the calcification of the [atherosclerotic plaque](#) core.

D - Intimal migration of smooth muscle cells

Image



Explanation Why

[Platelet-derived growth factor](#) is a growth factor that binds to [tyrosine kinase receptors](#) and stimulates the transition of cells from the [G₁ phase](#) to the [S phase](#). [PDGF](#) is released at atherosclerotic lesion sites by [platelets](#) and [macrophages](#) and stimulates [proliferation](#) and migration of [smooth muscles](#) cells in the [tunica intima](#), an important step in [atherosclerosis](#) that occurs in response to [endothelial](#) injury. [PDGF](#) also mediates the differentiation of [fibroblasts](#) into [myofibroblasts](#), which is necessary for fibrous cap formation.

E - Ingestion of cholesterol by mature monocytes

Explanation Why

During [atherosclerosis](#), [monocytes](#) migrate into the [intima](#) to mature into [macrophages](#), which ingest

[cholesterol](#) from oxidized [LDL](#) and then turn into [foam cells](#). Although [macrophages](#) can secrete [PDGF](#), ingestion of [cholesterol](#) by mature [monocytes](#) is not mediated by [PDGF](#).

F - Invasion of T-cells through the disrupted endothelium

Explanation Why

Invasion of [T-cells](#) through the disrupted [endothelium](#) is the result of increased permeability of the damaged [endothelium](#) during [atherosclerosis](#). Damaged [endothelial](#) cells express an increased number of cell-adhesion molecules, which mediate the adherence of [leukocytes](#). However, [PDGF](#) does not mediate the invasion of [T-cells](#) through the disrupted [endothelium](#).

G - Adherence of platelets to injured subendothelium

Explanation Why

[Platelet](#) adherence to damaged subendothelium in [atherosclerosis](#) is triggered by the increased permeability of [endothelium](#) due to the deleterious effects of long-standing [hypertension](#) or [diabetes](#). [PDGF](#) does not mediate [platelet](#) adherence.


H - Inflammatory cytokine release from the endothelium

Explanation Why

Inflammatory [cytokine](#) release from the [endothelium](#) (e.g., [IL-1](#) and [TNF- \$\alpha\$](#)) enhances the expression of cell-adhesion molecules on [endothelial](#) cells and promotes the migration of [leukocytes](#) in [atherosclerosis](#). [PDGF](#) does not mediate [cytokine](#) release from the [endothelium](#).

Question # 19

A 58-year-old man is brought to the emergency department by his wife 30 minutes after the sudden onset of severe retrosternal chest pain radiating to his back. He has a history of hyperlipidemia, hypertension, and type 2 diabetes mellitus. He has smoked one-half pack of cigarettes daily for 20 years. Medications include aspirin, captopril, atorvastatin, and metformin. His pulse is 80/min and blood pressure is 160/60 mm Hg. A CT scan of the chest is shown. Which of the following is the strongest predisposing factor for this patient's current condition?



	Answer	Image
A	Hyperlipidemia	
B	Diabetes mellitus	
C	Age	

	Answer	Image
D	Genetic collagen disorder	
E	Hypertension	Arterial hypertension Epidemiology Most common risk factor for cardiovascular disease Prevalence (adults): ~65% in individuals aged > 60 years Etiology 1. Primary (essential) hypertension (~90%), idiopathic 2. Secondary hypertension (~10%): Endocrine hypertension, e.g., due to: Hyperaldosteronism Hyperthyroidism Pheochromocytoma Renal hypertension, e.g., due to: Renal artery stenosis Glomerulonephritis Coarctation of the aorta Obstructive sleep apnea Medications (e.g., glucocorticoids, sympathomimetic drugs) Diagnostic criteria BP > 130/80 mmHg Subtypes Hypertensive crisis (i.e., systolic BP > 180 mmHg or diastolic BP > 120 mmHg) Isolated systolic hypertension (i.e., systolic BP > 140 mmHg and diastolic BP < 90 mmHg) White coat hypertension
F	History of smoking	

Hint

The patient's presentation (acute retrosternal chest pain radiating to the back, wide pulse pressure) and CT findings (double aortic lumen) suggest aortic dissection.

Correct Answer

A - Hyperlipidemia

Explanation Why

[Hyperlipidemia](#) can result in the formation of plaque in the [tunica intima](#) of arterial walls. This plaque can contribute to the development of [atherosclerosis](#), which increases the risk of [aortic dissection](#). However, [hyperlipidemia](#) is not the strongest predisposing factor for this condition.

B - Diabetes mellitus

Explanation Why

[Diabetes mellitus](#) is a [risk factor](#) for [atherosclerosis](#), which increases the risk of [aortic dissection](#). However, select studies have indicated that [diabetes mellitus](#) may have protective effects against [aortic aneurysm](#) formation and [aortic dissection](#). Therefore, [diabetes mellitus](#) is not the strongest predisposing factor for this patient's [aortic dissection](#).

C - Age

Explanation Why

The peak [incidence](#) of [aortic dissection](#) is 60–80 years of age. Therefore, this patient's age (58 years) is not the strongest predisposing factor for his [aortic dissection](#).


D - Genetic collagen disorder

Explanation Why

Genetic [collagen](#) disorders (e.g. [Ehlers-Danlos](#) syndrome) are a common cause of [aortic dissection](#) in patients ≤ 40 years of age. These disorders are very rare in older patients with [aortic dissection](#). In addition, these disorders more commonly involve the [ascending aorta](#) rather than the descending, which is involved in this patient. Moreover, this patient has no [family history](#), personal history, or features (e.g., long fingers, hyperextensible [joints](#), or hyperelastic [skin](#)) that would raise concern for these conditions.

E - Hypertension

Image



Explanation Why

[Aortic dissection](#) most commonly occurs secondary to [hypertension](#); more than 70% of patients with

[aortic dissection](#) have [elevated blood pressure](#), and [hypertension](#) accounts for more than 50% of the population at risk of [aortic dissection](#). Uncontrolled [hypertension](#) leads to propagation of [aortic dissection](#) and increases the risk of rupture.

F - History of smoking

Explanation But

Smoking is the strongest modifiable predisposing factor for abdominal [aortic aneurysms](#).

Explanation Why

Smoking is a [risk factor for atherosclerosis](#), which is associated with [aortic dissection](#), and an independent [risk factor](#) for [aortic dissection](#). Cessation of smoking is an important preventive measure for [aortic dissection](#) but smoking is not the strongest predisposing factor for this condition.

Question # 20

A 52-year-old man comes to the physician for a routine health maintenance examination. He has not seen a physician for 10 years. He works as a telemarketer and does not exercise. Cardiac examination shows a dull, low-pitched sound during late diastole that is best heard at the apex. The sound is loudest in the left lateral decubitus position and during end-expiration. Which of the following is the most likely cause of this finding?

	Answer	Image
A	Concentric left ventricular hypertrophy	
B	Congenital atrial septal defect	
C	Dilation of both ventricles	
D	Fusion of mitral valve leaflets	
E	Right bundle branch block	
F	Aortic root dilatation	

Hint


What is described here is a left-sided fourth heart sound (S4).

Correct Answer

A - Concentric left ventricular hypertrophy

Image

- ① Aortic valve
- ② Pulmonary valve
- ③ Tricuspid valve
- ④ Mitral valve
- ⑤ Erb's point



Explanation Why

The [fourth heart sound](#) is the result of filling a noncompliant ventricle through atrial contraction in late [diastole](#). Positioning the patient in the left-[lateral](#) decubitus position increases filling of the [left atrium](#) and can thus amplify the [heart sound](#) on auscultation. Concentric [left ventricular hypertrophy](#) decreases ventricle compliance, which is classically seen in advanced age. In this middle-aged, physically inactive patient, the most likely antecedent cause for [left ventricular hypertrophy](#) is hypertensive [heart](#) disease secondary to persistently increased [afterload](#).

B - Congenital atrial septal defect

Explanation Why

Large [atrial septal defects](#) can lead to volume overload of the [right ventricle](#) (due to [left-to-right shunting](#)) and [pulmonary hypertension](#). [Pulmonary hypertension](#) can result in [right ventricular dysfunction](#) and a right-sided [fourth heart sound](#). However, significant defects would be unlikely to remain asymptomatic until the age of 52. Moreover, this patient does not have the wide, fixed [splitting of S2](#), which is a characteristic finding of [atrial septal defects](#) on auscultation.

C - Dilation of both ventricles

Explanation Why

Dilation of both ventricles occurs in [dilated cardiomyopathy](#) and chronic [heart failure](#). However, this patient has no symptoms of these conditions (e.g., [nocturia](#), fatigue, [tachyarrhythmia](#)) and auscultation revealed a [fourth heart sound](#) rather than a [third heart sound](#) ([S3 gallop](#)), which would be expected in ventricular dilation.

D - Fusion of mitral valve leaflets

Explanation Why

Fusion of the [mitral valve](#) leaflets leads to [mitral stenosis](#), which is often asymptomatic. Upon [cardiac auscultation](#), [mitral stenosis](#) can produce a low-pitched [diastolic](#) rumbling [murmur](#), the duration of which depends on the severity of the stenosis. Other auscultatory findings may include a loud S1 and S2 and an [opening snap](#). However, the [fourth heart sound](#) detected in this patient would not be expected.

E - Right bundle branch block

Explanation Why

[Right bundle branch block](#) most commonly occurs as a consequence of [right ventricular](#) distension, [myocardial infarction](#), [myocarditis](#), [hypertension](#), or congenital [heart](#) disease. It is mostly asymptomatic but may manifest with a wide, fixed-splitting of S2 due to asynchronous ventricular contraction. [RBBB](#) does not cause a [fourth heart sound](#).


F - Aortic root dilatation

Explanation Why

Dilatation of the aortic root occurs in patients with [connective tissue disorders](#) (e.g., [Marfan syndrome](#)), a [bicuspid aortic valve](#), [hypertension](#), [atherosclerosis](#), [aortic dissection](#), or [aortic stenosis](#). Significant dilatation of the aortic annulus can lead to [aortic regurgitation](#), which may be asymptomatic in chronic cases similar to the disorder affecting this patient. However, auscultation would reveal a decrescendo [diastolic murmur](#), unlike the [fourth heart sound](#) detected in this patient.

Question # 21

An investigator is studying the physiological factors involved in regulating coronary vessel blood flow during high metabolic activity in healthy human volunteers. In addition to the direct effects of coronary venous oxygen and carbon dioxide concentrations, the effects of adenosine, ATP-dependent potassium channels, and adrenergic activity on coronary vasodilation are examined. The volunteers undergo cardiac catheterization to measure left ventricular, left atrial, and aortic pressures. An image of the pressure tracings in one healthy volunteer is shown. Blood flow through the coronary arteries is most likely to peak at which of the following labeled points of the diagram?



	Answer	Image
A	A	
B	B	
C	C	

	Answer	Image
D	D	
E	E	Pressure (mm Hg) Systole Contraction Ejection Relaxation Diastole Filling 120 80 40 0 Time Left ventricle Aorta Left atrium 1 2 3 4
F	F	

Hint

External compression of the coronary vessels by myocardial tissue results in paradoxical peak flow.

Correct Answer

A - A

Explanation Why

This point marks the onset of the [isovolumetric contraction](#) phase of [systole](#) and closure of the [mitral valve](#). The blood flow through the coronary vessels would peak at a different point in the [cardiac cycle](#).

B - B

Explanation Why

This point marks the onset of the [rapid ejection phase](#) of [systole](#) and the opening of the [aortic valve](#), identified by the sudden rise in aortic pressure. The blood flow through the coronary vessels would peak at a different point in the [cardiac cycle](#).

C - C

Explanation Why

This point marks peak [systolic](#) pressure in both the ventricle and the aorta. The blood flow through peripheral [arteries](#) would reach its peak at this point in the [cardiac cycle](#), but the blood flow through the coronary vessels would reach its [nadir](#).


D - D

Explanation Why

This point marks the onset of the [isovolumetric relaxation](#) phase of ventricular [diastole](#) and the closure of the [aortic valve](#). The blood flow through the coronary vessels would peak at a different point in the [cardiac cycle](#).

E - E

Image



Explanation But

During exercise, the shortened duration of [diastole](#) is a limiting factor for coronary blood flow.

Explanation Why

In most peripheral [arteries](#), blood flow peaks at the end of the [rapid ejection phase](#) of [systole](#) when the ventricular and aortic pressures are the highest (point C). However, [coronary arteries](#), which arise from the aortic root, are unique because they are embedded within the ventricular [myocardium](#) and are compressed when ventricular wall tension increases. Therefore, blood flow through the coronary vessels occurs predominantly during [diastole](#) when the ventricular pressure is lower than the aortic pressure. Thus, the point of peak coronary flow in the [cardiac cycle](#) occurs when the pressure differential between the aorta and the [left ventricle](#) is the greatest (point E).


F - F

Explanation Why

This point marks the slow [filling phase](#) of ventricular [diastole](#) and the onset of atrial contraction (“atrial kick”). The blood flow through the coronary vessels would peak at a different point in the [cardiac cycle](#).

Question # 22

A 72-year-old woman is admitted to the intensive care unit for shortness of breath and palpitations. A cardiac catheterization is performed and measurements of the left ventricular volume and pressure at different points in the cardiac cycle are obtained. The patient's pressure-volume loop (gray) is shown with a normal pressure-volume loop (black) for comparison. Which of the following is the most likely underlying cause of this patient's symptoms?



	Answer	Image
A	Mitral valve regurgitation	The image shows a pressure-volume loop graph with 'LV pressure [mmHg]' on the y-axis (0 to 200) and 'LV volume [mL]' on the x-axis (50 to 200). It compares two loops: a 'Normal' loop (solid line) and a 'Mitral regurgitation' loop (dashed line). The normal loop has a higher stroke volume, ending at approximately 150 mL. The mitral regurgitation loop is shifted to the left, ending at approximately 100 mL, which indicates a smaller stroke volume.

	Answer	Image
B	Increased systemic vascular resistance	
C	Increased ventricular wall stiffness	
D	Impaired left ventricular contractility	
E	Aortic stenosis	


Hint

This patient's pressure-volume loop shows increased left ventricular pressures for a given volume during the filling phase of diastole (i.e., a decreased ability to expand in volume in response to pressure), indicating diastolic dysfunction.

Correct Answer

A - Mitral valve regurgitation

Image



Explanation Why

In [mitral valve regurgitation \(MR\)](#), the [stroke volume](#) is divided between the output through the [aortic valve](#) and through the abnormal regurgitant [mitral valve](#), which results in reduced [afterload](#) ($\downarrow$ outflow resistance) and decreased end-[systolic](#) volume (top left of the pressure-volume loop). This patient's [pressure-volume loop](#) also shows a decreased [end-systolic volume](#) but, in [MR](#), there is an increase in left-ventricular end-[diastolic](#) volume (bottom right), which is not seen here. This ventricular volume overload results in increased [preload](#), which increases the [stroke volume](#) (apparent in a [pressure-volume loop](#) that is widened, not narrowed). Moreover, there is no true [isovolumetric contraction](#) phase in [MR](#).


B - Increased systemic vascular resistance

Explanation Why

Increased [systemic vascular resistance](#) (SVR) increases [afterload](#), resulting in a [pressure-volume loop](#) that is taller and narrower than normal. The increased height reflects the fact that the [left ventricle](#) must reach a higher pressure to overcome the aortic pressure (so that the [aortic valve](#) opens), and, consequently, an end-[systolic](#) pressure that is higher than normal. The [pressure-volume loop](#) is narrower because the increased [afterload](#) from high SVR leads to a decreased [stroke volume](#).

C - Increased ventricular wall stiffness

Image



Explanation But

Causes of [diastolic heart failure](#) include concentric [left ventricular hypertrophy](#) secondary to [hypertension](#) (most common cause), [restrictive cardiomyopathy](#) (most commonly due to


[amyloidosis](#)), and [hypertrophic cardiomyopathy](#).

Explanation Why

Increased ventricular wall stiffness decreases ventricular compliance, which is the underlying cause of [diastolic heart failure](#). Stiffening of the ventricles results in defective [diastolic](#) filling, while [ejection fraction](#) (i.e., [systolic](#) function) is (for the most part) preserved. The slope of the end-[diastolic](#) pressure-volume curve (the bottom line of the [pressure-volume loop](#)) is the inverse of the ventricular compliance, which is decreased in this patient. The narrowed pressure-volume loop shows a decreased [stroke volume](#), which is also classically seen in [diastolic heart failure](#). Patients typically present with signs and symptoms of heart failure (e.g., [shortness of breath](#), [jugular venous distention](#)) and an S4 [atrial gallop](#), which is caused by contraction of the [atria](#) against the noncompliant ventricles.

D - Impaired left ventricular contractility

Image




Explanation Why

Impaired [left ventricular contractility](#) is consistent with [systolic heart failure](#), in which case the [pressure-volume loop](#) is shifted to the right, not upwards and to the left. In [systolic heart failure](#), the

end-systolic volume (top left of the [pressure-volume loop](#)) is increased due to decreased contractility of the [left ventricle](#), which results in decreased [cardiac output](#). A compensatory increase in end-diastolic volume (bottom right) elevates [preload](#) and helps maintain [stroke volume](#) via the [Frank-Starling](#) mechanism. However, [ejection fraction](#) is still significantly reduced. The slope of the end-diastolic pressure-volume curve (the bottom line of the [pressure-volume loop](#)) is normal, as the ventricle can expand normally during [diastolic](#) filling.

E - Aortic stenosis

Image



Explanation Why

The [pressure-volume loop](#) in [aortic stenosis](#) (AS) is narrower and taller than normal, not shifted upwards and to the left. In AS, the increased height of the [pressure-volume loop](#) is due to [left ventricular](#) outflow obstruction, which increases the peak [left ventricular](#) pressure during [systole](#), and the [pressure-volume loop](#) is narrower because the increased [afterload](#) from [aortic stenosis](#) leads to decreased [stroke volume](#).

Question # 23

A 45-year-old man comes to the physician because of a 1-month history of fever and poor appetite. Five weeks ago, he underwent molar extraction for dental caries. His temperature is 38°C (100.4°F). Cardiac examination shows a grade 2/6 holosystolic murmur heard best at the apex. A blood culture shows gram-positive, catalase-negative cocci. Transesophageal echocardiography shows a small vegetation on the mitral valve with mild regurgitation. The causal organism most likely has which of the following characteristics?

	Answer	Image
A	Production of CAMP factor	
B	Replication in host macrophages	
C	Formation of germ tubes at body temperature	
D	Production of dextrans	
E	Conversion of fibrinogen to fibrin	

Hint

Together with echocardiography showing a small vegetation, protracted fever and loss of appetite after a dental extraction suggest subacute bacterial endocarditis, which is most commonly due to viridans group streptococci.

Correct Answer

A - Production of CAMP factor

Explanation Why

Production of [CAMP factor](#) is a unique property of *S. agalactiae*, a gram-positive, catalase-negative coccus. While *S. agalactiae* can also cause [subacute bacterial endocarditis](#), it is a very rare causative agent and is usually associated with [diabetes mellitus](#), [malignancy](#), or [immunosuppression](#), not with recent dental surgery.

B - Replication in host macrophages

Explanation Why

Replication in host [macrophages](#) is a strategy adopted by many pathogens, including [Coxiella burnetii](#), which can also cause [endocarditis](#) (e.g., due to untreated [Q fever](#)). However, *C. burnetii* is a gram-negative bacterium that is very difficult to culture, in contrast to this patient's gram-positive [bacterial culture](#) result. [Viridans group streptococci](#) do not replicate in host [macrophages](#).

C - Formation of germ tubes at body temperature

Explanation Why

Formation of germ tubes at body temperature is a characteristic feature of [Candida](#) spp., which can cause [endocarditis](#). Fungal [endocarditis](#) typically manifests with small vegetations. [Risk factors](#) for fungal [endocarditis](#) include recent surgery, IV drug use, and [immunosuppression](#), none of which are present in this patient. Furthermore, blood culture would reveal budding [yeast](#), not gram-positive, catalase-negative [cocci](#).

D - Production of dextrans

Explanation Why

Dental surgery is an important [risk factor](#) for [subacute endocarditis](#) with [viridans group streptococci](#), which are present in normal oral flora. [Viridans group streptococci](#) (especially *S. mutans* and *S. sanguis*) produce [dextrans](#), which facilitates binding to tooth [enamel](#), as well as to [fibrin-platelet](#) deposits on damaged [heart valves](#). The subsequent formation of a [fibrin](#) clot and ensuing [inflammation](#) results in the vegetation formation that is characteristic of [endocarditis](#).

E - Conversion of fibrinogen to fibrin

Explanation Why

[Coagulase](#), which converts [fibrinogen](#) to [fibrin](#), is an enzyme produced by [Staphylococcus aureus](#). [S. aureus](#) is also a gram-positive coccus and one of the most common causes of bacterial [endocarditis](#). However, [S. aureus](#) is also catalase-positive, whereas the organism seen here is catalase-negative. Furthermore, [S. aureus](#) is typically associated with more rapid symptom onset and the formation of relatively large vegetations.

Question # 24

A 32-year-old woman brought to the emergency department because of a 1-week history of palpitations and shortness of breath. She has congestive heart failure. Current medications include furosemide, lisinopril, and atenolol. Her pulse is 124/min and irregularly irregular, and blood pressure is 110/70 mm Hg. Examination shows coarse crackles over the lower lung fields bilaterally. Treatment with digoxin is started. Five days later, an ECG shows prolongation of the PR interval. Which of the following is the most likely explanation for the observed effect of this drug?

	Answer	Image
A	Inhibition of myocardial Na^+/K^+ ATPase	
B	Inhibition of AV node L-type Ca^{2+} channels	
C	Increase in vagal tone	
D	Activation of $\text{Na}^+/\text{Ca}^{2+}$ exchanger	
E	Decrease in intracellular cAMP	

Hint

The PR interval measures the time between the onset of atrial depolarization and the onset of ventricular depolarization and expresses conduction through the AV node.

Correct Answer

A - Inhibition of myocardial Na^+/K^+ ATPase

Explanation Why

The primary action of [digoxin](#), inhibition of [myocardial \$\text{Na}^+/\text{K}^+\$ ATPase](#), results in a higher intracellular Na^+ concentration and reduced efficacy of [\$\text{Na}^+/\text{Ca}^{2+}\$ exchangers](#), which leads to higher intracellular Ca^{2+} concentration and, subsequently, increased cardiac [inotropy](#). However, [digoxin](#)'s action on the [AV node](#) resulting in [PR interval](#) prolongation occurs through a different mechanism.

B - Inhibition of AV node L-type Ca^{2+} channels

Explanation Why

[Calcium channel blockers](#) act via the inhibition of [AV nodal L-type \$\text{Ca}^{2+}\$ channels](#) and can cause [PR interval](#) prolongation. However, [digoxin](#)'s action on the [AV node](#) resulting in [PR interval](#) prolongation occurs through a different mechanism.

C - Increase in vagal tone

Explanation But

Younger patients taking [cardiac glycosides](#) are at an increased risk of developing [bradyarrhythmia](#); older patients more often develop [ventricular tachyarrhythmias](#).

Explanation Why

[Digoxin](#) indirectly increases [vagal](#) tone via inhibition of neuronal [\$\text{Na}^+/\text{K}^+\$ ATPase](#), causing a negative [chronotropic](#) effect that leads to a [PR interval](#) prolongation as seen in this patient's [ECG](#). This indirect effect on [vagal](#) tone can lead to [bradyarrhythmia](#) and increase the risk of [AV block](#).

[Cardiac glycosides](#) have a [narrow therapeutic index](#) and require close monitoring of serum concentrations. [Hypokalemia](#) increases the risk of [digoxin toxicity](#) by enhancing its affinity for [Na⁺/K⁺ ATPase](#). Other [ECG](#) findings seen with [digoxin](#) use include [premature ventricular beats](#), [T-wave inversion](#) or flattening, downsloping ST-segment depression, and a decrease in the [QT interval](#).

D - Activation of Na⁺/Ca²⁺ exchanger

Explanation Why

[Digoxin](#) inhibits [Na⁺/K⁺ ATPase](#) in [cardiomyocytes](#), which then indirectly inhibit, rather than activate, the activity of [Na⁺/Ca²⁺ exchangers](#). This inhibitory effect leads to higher intracellular [Ca²⁺](#) concentrations and an increase in cardiac contractility. However, [digoxin](#)'s action on the [AV node](#) resulting in [PR interval](#) prolongation occurs through a different mechanism.

E - Decrease in intracellular cAMP

Explanation Why

[Beta blockers](#) inhibit the [sympathetic](#) activation of [β-adrenergic receptors](#) leading to [PR interval](#) prolongation, [bradycardia](#), and decreased cardiac contractility. However, [digoxin](#)'s action on the [AV node](#) resulting in [PR interval](#) prolongation occurs through a different mechanism.

Question # 25

A 67-year-old man comes to the emergency department because of retrosternal chest pressure and shortness of breath for 4 hours. The symptoms started while he was walking to work and have only minimally improved with rest. He has a history of type 2 diabetes mellitus. He has smoked one pack of cigarettes daily for 35 years. He appears uncomfortable. His pulse is 95/min. Serum studies show a normal troponin concentration. An ECG shows no abnormalities. Which of the following is the most likely underlying cause of this patient's symptoms?

	Answer	Image
A	Aortic valve thickening and calcification	
B	Coronary artery occlusion due to transient increase in vascular tone	
C	Stable atherosclerotic plaque with 85% coronary artery occlusion	
D	Infarction and rupture of a papillary muscle	
E	Disruption of an atherosclerotic plaque with a non-occlusive coronary artery thrombus	
F	Atherosclerotic plaque thrombus with complete coronary artery occlusion	

Hint

This patient with multiple cardiovascular risk factors (older age, male sex, history of diabetes mellitus and smoking) has unstable angina pectoris, a condition that can cause chest pain and dyspnea at rest without cardiac biomarker elevation.

Correct Answer

A - Aortic valve thickening and calcification

Explanation Why

[Aortic valve](#) thickening and calcification result in [aortic stenosis](#). This condition typically remains asymptomatic for years, but exertional [angina](#) and [dyspnea](#) can develop gradually as the [aortic valve](#) becomes increasingly stenotic. If the stenosis becomes severe, patients can experience symptoms at rest. [Aortic stenosis](#) is not likely to initially manifest with sudden-onset symptoms that do not improve with rest as seen in this patient.

B - Coronary artery occlusion due to transient increase in vascular tone

Explanation Why

[Coronary artery](#) occlusion due to a transient increase in vascular tone is the underlying pathology of [vasospastic angina](#), which causes episodic [angina](#) and [dyspnea](#) at rest. Though [vasospastic angina](#) does not cause increased serum concentrations of [cardiac biomarkers](#) such as [troponin](#), it is associated with reversible [ST elevations](#) on [ECG](#), which are not seen in this patient. Moreover, episodes of [angina](#) caused by this condition typically last for 5–15 minutes. This patient has had 4 hours of progressive symptoms.

C - Stable atherosclerotic plaque with 85% coronary artery occlusion

Explanation Why

A stable [atherosclerotic plaque](#) with $\geq 70\%$ occlusion of a [coronary artery](#) can cause [stable angina pectoris](#), which manifests with exertional [chest pain](#) and [dyspnea](#) that resolve with rest. This patient's symptoms only minimally improved with rest. Moreover, a diagnosis of [stable angina pectoris](#) requires a history of prior episodes of [anginal chest pain](#), and there is no indication that this patient has had similar episodes in the past.

D - Infarction and rupture of a papillary muscle

Explanation Why

[Papillary muscle rupture](#) is a life-threatening complication of [myocardial infarction](#) that most commonly develops 2–7 days after [infarction](#) and involves the [mitral valve](#), which leads to sudden-onset [hypotension](#) and severe [mitral regurgitation](#) that can manifest with [dyspnea](#). This complication would not develop in the acute phase of [infarction](#).

E -

Disruption of an atherosclerotic plaque with a non-occlusive coronary artery thrombus

Explanation Why

Disruption of an [atherosclerotic plaque](#) with a non-occlusive [coronary artery thrombus](#) can impair [coronary artery](#) perfusion and result in [myocardial ischemia](#) that causes [unstable angina pectoris](#). This condition includes new-onset [angina pectoris](#) and [angina pectoris](#) that increases in frequency or severity over time. Symptoms typically persist for ≥ 20 minutes and can occur at rest or with minimal exertion as seen in this case. [Unstable angina pectoris](#) does not cause [cardiac biomarker](#) elevation. [ECG](#) can be normal or show [ST depression](#) and/or [T-wave inversion](#).

F - Atherosclerotic plaque thrombus with complete coronary artery occlusion

Explanation Why

An [atherosclerotic plaque thrombus](#) with complete [coronary artery](#) occlusion results in [myocardial infarction](#) (e.g., [STEMI](#), [NSTEMI](#)), which manifests with [chest pain](#) and [dyspnea](#). However, [myocardial infarction](#) causes increased serum concentrations of [cardiac biomarkers](#) such as [troponin](#), which are not seen in this patient. Moreover, [myocardial infarction](#) is usually associated with characteristic [ECG](#) changes (e.g., [ST elevation](#), [ST depression](#)).

Question # 26

A group of investigators is studying a drug to treat refractory angina pectoris. This drug works by selectively inhibiting the late influx of sodium ions into cardiac myocytes. At high doses, the drug also partially inhibits the degradation of fatty acids. Which of the following is the most likely effect of this drug?

	Answer	Image
A	Increased oxygen efficiency	
B	Increased prolactin release	
C	Decreased uric acid excretion	
D	Decreased insulin release	
E	Decreased serum pH	
F	Increased parasympathetic tone	

Hint

The drug in question is ranolazine.

Correct Answer

A - Increased oxygen efficiency

Explanation Why

[Ranolazine](#) increases oxygen efficiency in [cardiomyocytes](#) through two mechanisms. First, the drug decreases [myocardial](#) tension during [diastole](#) by impairing late phase sodium influx into cardiac [myocytes](#), disrupting the sodium-calcium exchange mechanism and leading to reduced calcium flux and wall tension. Second, [ranolazine](#) causes [cardiomyocytes](#) to favor [glycolysis](#) over the [beta-oxidation](#) of [fatty acids](#) for energy generation. Because [glycolysis](#) is not dependent on oxygen availability and [beta-oxidation](#) is oxygen-intensive, this shift reduces the oxygen needs of cardiac tissue. Importantly, neither mechanism causes significant changes in the [heart rate](#) or blood pressure.

B - Increased prolactin release

Explanation Why

Increased [prolactin](#) release can arise as a side effect of non-dihydropyridine [calcium channel blockers](#) (e.g., [verapamil](#)). [Hyperprolactinemia](#) is thought to occur as a result of central inhibition of [dopamine](#) production, which ordinarily prevents the production of [prolactin](#). While [verapamil](#) can effectively treat [angina](#) by decreasing [heart rate](#) and contractility, it does not affect [fatty acid metabolism](#) or selectively inhibit the late influx of sodium ions in cardiac [myocytes](#).

C - Decreased uric acid excretion

Explanation Why

Decreased [uric acid](#) excretion can be a consequence of treatment with low-dose [aspirin](#) or [niacin](#). [Aspirin](#) is prescribed to patients with [angina](#) as [secondary prevention](#). However, [aspirin](#) tends to increase [mitochondrial](#) long-chain [fatty acid oxidation](#), whereas the drug in question reduces it. [Niacin](#) is a lipid-lowering drug that is seldom used as [secondary prevention](#) for patients with [angina](#) ([statins](#) are much more commonly used in this role). In addition, neither [aspirin](#) nor [niacin](#) directly treats [anginal pain](#), and neither selectively inhibits sodium influx into cardiac [myocytes](#).

D - Decreased insulin release

Explanation Why

Decreased [insulin](#) release occurs as a consequence of medications that block the [beta 2 adrenergic receptor](#). Drugs in this class, commonly known as [beta blockers](#) (e.g., [propranolol](#)), will also have an inhibitory effect on [lipolysis](#) as a result of [beta receptor](#) blockage on [adipocytes](#). [Beta blockers](#) are very commonly used to treat [angina](#). They do not, however, inhibit the late influx of sodium ions in [cardiomyocytes](#).

E - Decreased serum pH

Explanation Why

Decreased serum pH ([acidosis](#)) would be seen in instances of increased [fatty acid oxidation](#). [Fatty acid oxidation](#) produces ketoacids, which enter the peripheral circulation and cause the pH to decrease. The drug in this case acts to inhibit [fatty acid oxidation](#) and, ultimately, reduce the production of ketoacids. Therefore, this drug would increase rather than decrease serum pH.


F - Increased parasympathetic tone

Explanation Why

Increased [parasympathetic](#) tone from the [vagus nerve](#) results in a reduced [heart rate](#) and is seen with [digoxin](#) administration. This drug inhibits the efflux of sodium ions from [cardiomyocytes](#) through the sodium-potassium pump. However, [digoxin](#) does not selectively inhibit the late influx of sodium ions into cardiac [myocytes](#), and it has no effect on [fatty acid metabolism](#). [Digoxin](#) is used to treat [heart failure](#) and atrial [tachyarrhythmias](#) (e.g., [atrial fibrillation](#) and flutter) in patients who have [congestive heart failure](#), but it is not used to treat [angina](#).

Question # 27

A 67-year-old woman comes to the physician because of intermittent chest pain and dizziness on exertion for 6 months. Her pulse is 76/min and blood pressure is 125/82 mm Hg. Cardiac examination shows a grade 3/6, late-peaking, crescendo-decrescendo murmur heard best at the right upper sternal border. An echocardiogram confirms the diagnosis. Three months later, the patient returns to the physician with worsening shortness of breath for 2 weeks. An ECG is shown. Which of the following changes is most likely responsible for this patient's acute exacerbation of symptoms?



	Answer	Image
A	Impaired contractility of the left ventricle	
B	Impaired pulmonary artery outflow	
C	Increased myocardial perfusion pressure gradient	

	Answer	Image
D	Decreased left ventricular preload	
E	Decreased impulse conduction across the AV node	
F	Increased systemic vascular resistance	

Hint

This patient's murmur is most likely caused by aortic stenosis. Her ECG shows findings of atrial fibrillation, including tachycardia, irregularly irregular RR intervals, indiscernible P waves, and narrow QRS complexes.

Correct Answer

A - Impaired contractility of the left ventricle

Explanation Why

Impaired [left ventricular contractility](#) can occur as a result of [myocardial infarction](#) (due to impaired cardiac relaxation) or [left ventricular hypertrophy](#) (due to [interstitial fibrosis](#) of the [myocardium](#)). The [ECG](#) findings, however, do not show any signs of [ischemia](#) (e.g., [ST elevation](#), [T-wave inversion](#)), and [chest pain](#) associated with [myocardial infarction](#) is usually persistent, not intermittent. While [aortic stenosis](#) can cause [left ventricular hypertrophy](#), this patient's acute [dyspnea](#) is most likely a manifestation of [atrial fibrillation](#), which can result in uncoordinated contractions of the [atrium](#) but does not impair the contractility of the [left ventricle](#).

B - Impaired pulmonary artery outflow

Explanation Why

Conditions that impair [pulmonary artery](#) outflow include [pulmonary hypertension](#) (PH) and [pulmonic stenosis](#) (PS). Either of these conditions can cause [angina](#) and [dizziness](#) (due to [right ventricular ischemia](#) in PH and due to progressive [heart failure](#) in PS). However, neither of these conditions is associated with [atrial fibrillation](#) or a crescendo-decrescendo [murmur](#). On auscultation, PH causes a loud [second heart sound](#), and PS causes a [systolic ejection murmur](#) heard best at the left upper sternal border and [splitting of the second heart sound](#).

C - Increased myocardial perfusion pressure gradient

Explanation Why

[Coronary artery](#) blood flow predominantly occurs during [diastole](#) and is driven by the myocardial perfusion pressure gradient. [Atrial fibrillation](#) causes a decreased, not increased, myocardial perfusion pressure gradient as a result of uncoordinated atrial contraction (which impairs [diastolic](#) filling of the [left ventricle](#)) and [tachycardia](#) (which shortens the duration of [diastole](#)). In addition, this patient has [aortic stenosis](#), which often causes [left ventricular hypertrophy](#). [Left ventricular](#)

[hypertrophy](#) further compromises the myocardial perfusion pressure gradient by impeding [diastolic](#) filling.

D - Decreased left ventricular preload

Explanation Why

[Aortic stenosis](#) (AS) causes increased [left ventricular afterload](#) and [end-systolic volume](#), which results in an increase in filling pressures and concentric [left ventricular hypertrophy](#). Because [left ventricular hypertrophy](#) commonly results in [diastolic dysfunction](#), individuals with AS rely on the contraction of the [left atrium](#) for [ventricular filling](#). The increase in [diastolic](#) filling pressure can, in turn, cause left atrial dilatation, which in turn increases the risk of [atrial fibrillation](#) (AF). AF can decrease the [left ventricular preload](#) as a result of uncoordinated atrial contraction (which impairs [diastolic](#) filling of the [left ventricle](#)) and [tachycardia](#) (which shortens the duration of [diastole](#)). If AF develops in individuals with severe AS, it can exacerbate symptoms of [angina](#) and [dyspnea](#) due to [pulmonary edema](#), which is likely responsible for this patient's worsening [shortness of breath](#).

E - Decreased impulse conduction across the AV node

Explanation Why

Decreased impulse conduction across the [AV node](#) occurs in patients with nodal conduction abnormalities, which can cause [angina](#), [dizziness](#), and/or an irregular [pulse](#), depending on the degree of [AV block](#). However, these conditions typically cause [bradycardia](#), not [tachycardia](#), and are associated with discernable [P waves](#) on [ECG](#). Moreover, none of these conditions is associated with a crescendo-decrescendo [murmur](#).

F - Increased systemic vascular resistance

Explanation Why

Increased [systemic vascular resistance](#) (SVR) occurs as part of the compensatory response to maintain tissue perfusion in patients with [aortic stenosis](#). A stenotic [aortic valve](#) reduces [cardiac output](#) and leads to hypoperfusion, which results in [angina](#) and [dizziness](#). An increase in SVR is not,

however, responsible for the acute exacerbation of symptoms, which is most likely related to [atrial fibrillation](#).

Question # 28

A 42-year-old man with systolic heart failure secondary to amyloidosis undergoes heart transplantation. The donor heart is obtained from a 17-year-old boy who died in a motor vehicle collision. Examination of the donor heart during the procedure shows a flat, yellow-white discoloration with an irregular border on the luminal surface of the aorta. A biopsy of this lesion is most likely to show which of the following?

	Answer	Image
A	Lipoprotein-laden macrophages	The diagram illustrates the progression of atherosclerosis over time, from asymptomatic stages to minutes before a thrombus forms. The stages are: Normal endothelium, Asymptomatic (Endothelial dysfunction, Infiltration of monocytes and LDL resulting in accumulation of macrophages and oxLDL), Decades (Foam cell formation resulting in fatty streaks, Proliferation and migration of SMCs and deposition of ECM), Symptomatic (Fibrous plaque formation, Plaque destabilization), and Minutes (Plaque rupture (e.g., in context of increased matrix degradation), Thrombus formation). A red arrow indicates the progression from left to right. A legend identifies various components: LDL, Monocyte, Macrophage, oxLDL, Foam cell, Platelets, Lymphocytes, Extracellular matrix (ECM), Smooth muscle cell (SMC), Cholesterol crystals, and Cellular debris. Key features include chronic stress (e.g., due to hypertension, inflammation), PDGF, FGF, Fibrous cap, Necrotic core, and SMC apoptosis/breakdown of the stabilizing ECM.
B	Proteoglycan accumulation	
C	Apoptotic smooth muscle cells	
D	Necrotic cell debris	
E	Dystrophic calcification	

	Answer	Image
F	Collagen deposition	


Hint

The morphology (flat, yellow-white intimal discoloration) and location (luminal surface) of the lesion corresponds with the fatty streak stage of atherogenesis.

Correct Answer

A - Lipoprotein-laden macrophages

Image



Explanation Why

Lipoprotein-laden macrophages (foam cells) within the intimal layer of vessels are seen in the early stage of atherosclerosis, which corresponds to the fatty streak phase. Fatty streaks appear macroscopically as flat yellow discolorations with irregular borders on the luminal surfaces of blood vessels, including the aorta. Starting in early adulthood, fatty streaks may be visible with the naked eye.

B - Proteoglycan accumulation

Explanation Why

[Proteoglycan](#) accumulation is seen in the [fibrous plaque](#) phase of atherogenesis. [Fibrous plaques](#) are the result of the accumulation and [proliferation](#) of lipid-filled [smooth muscle cells](#) and [macrophages](#), which induce deposition of [extracellular matrix](#) (e.g., [proteoglycans](#) and [elastin](#)). They are seen in patients with a long history of [atherosclerosis](#), which is unlikely in a 17-year-old donor.

C - Apoptotic smooth muscle cells

Explanation Why

[Apoptosis](#) of [smooth muscle cells](#) is seen in the late stages of [fatty streak](#) formation. As the [fatty streak](#) progresses, [smooth muscle](#) and [macrophages](#) accumulate in the [intimal](#) layer of vessels; [apoptosis](#) of the [smooth muscle cells](#) within the deep layer of the [fatty streak](#) can occur. These changes result in microvesicles that undergo calcification and contribute to the formation of [atherosclerotic plaques](#). On macroscopic examination, [atherosclerotic plaques](#) are visible as raised pale yellow lesions that impede and obstruct [arteries](#), rather than the flat lesions seen here.

D - Necrotic cell debris

Explanation Why

[Necrotic](#) cell debris can be seen in advanced [atherosclerosis](#). Advanced atherosclerotic lesions usually expand beyond the [intimal](#) layer of vessels and contain a [necrotic](#) lipid-rich core with calcifications. Macroscopically, advanced [atherosclerosis](#) appears as calcified, raised, pale, yellow lesions with a [necrotic](#) lipid-rich core and [cholesterol](#) crystallization, rather than the flat yellow lesions seen here. Moreover, advanced [atherosclerosis](#) would not be expected in a 17-year-old donor.

E - Dystrophic calcification

Explanation Why

[Dystrophic calcification](#) occurs in advanced atherosclerotic lesions that expand beyond the [intimal](#) layer of vessels. These lesions are most commonly seen in patients with a long history of [atherosclerosis](#). [Dystrophic](#) calcifications appear as raised pale yellow lesions with a [necrotic](#) lipid-rich core, rather than the flat yellow lesions seen here. Moreover, advanced [atherosclerosis](#) would not be expected in a 17-year-old donor.

F - Collagen deposition

Explanation Why

[Collagen](#) deposition is seen in the [fibrous plaque](#) phase of atherogenesis. [Fibrous plaques](#) are the result of the accumulation and [proliferation](#) of lipid-filled [smooth muscle cells](#) and deposition of [extracellular matrix](#) (e.g., [collagen](#) and [elastin](#)). They are seen in patients with a long history of [atherosclerosis](#), which is unlikely in a 17-year-old donor.

Question # 29

A 62-year-old woman comes to the physician because of a 2-month history of exertional shortness of breath and fatigue. She sometimes wakes up at night coughing and gasping for air. Cardiac examination shows a grade 3/6 holosystolic murmur best heard at the apex. Which of the following physical exam findings would be consistent with an exacerbation of this patient's condition?

	Answer	Image
A	Paradoxical splitting of A2 and P2	
B	Early diastolic extra heart sound	
C	Prominent V wave	
D	Head bobbing	
E	Absence of A2 heart sound	
F	Radiation of the murmur to the axilla	

Hint

A holosystolic murmur at the cardiac apex suggests mitral regurgitation (MR).

Correct Answer


A - Paradoxical splitting of A2 and P2

Explanation Why

A paradoxical [splitting of the second heart sound](#) (A2 and P2) is seen in conditions that delay [aortic valve](#) closure such as [aortic regurgitation](#) (AR) and [aortic stenosis](#) (AS). Both these conditions can present with exertional dyspnea, [paroxysmal nocturnal dyspnea](#), fatigue, and a cardiac [murmur](#), as seen here. However, the [murmur](#) of AR is a decrescendo, early [diastolic murmur](#), and the [murmur](#) of AS is a crescendo-decrescendo [systolic ejection](#). Also, both these [murmurs](#) would be best heard at the right upper sternal border ([aortic area](#)) rather than the cardiac apex. This patient's auscultatory finding is suggestive of [mitral regurgitation](#), which is not associated with a paradoxical [splitting of the second heart sound](#).

B - Early diastolic extra heart sound

Image



Explanation Why

The presence of a [diastolic](#) extra [heart sound](#) ([S3](#)) is associated with an exacerbation of a pre-existing [MR](#). Patients with chronic [mitral valve](#) insufficiency have an elevated ventricular volume and pressure during [diastole](#) as a result of insufficient valve closure, which leads to backflow of blood through the mitral valves. This higher blood volume entering the [left ventricle](#) during [diastole](#) results in an [S3 heart](#) sound on auscultation ([ventricular filling](#) sound). The presence of an [S3 heart](#) sound is associated with advanced-stage or a severe [MR](#).

C - Prominent V wave

Explanation Why

A prominent [V wave](#) on jugular venous examination (Lancisi sign) due to the merging of the [V wave](#) with the [C wave](#) is a sign of exacerbation in [tricuspid regurgitation](#) (TR). Severe TR can manifest with exertional dyspnea, [paroxysmal nocturnal dyspnea](#), fatigue, and a [holosystolic murmur](#) but the [murmur](#) would be best heard at the left lower sternal border ([tricuspid area](#)), not at the cardiac apex.

D - Head bobbing

Explanation Why

Head bobbing is a characteristic [physical exam](#) finding in patients with [aortic regurgitation](#) (AR). Although AR may present with exertional dyspnea, [paroxysmal nocturnal dyspnea](#), and fatigue, the [murmur](#) is classically early [diastolic](#), decrescendo, and best heard at the right upper sternal border, unlike the [holosystolic murmur](#) best heard at the apex here.

E - Absence of A2 heart sound

Explanation Why

The absence of an A2 heart sound can be seen in severe [aortic stenosis](#) (AS). The [second heart sound](#) (S2) is composed of two parts, the A2 (closure of aortic valves) and the P2 (closure of pulmonic

valves) sound. The A2 heart sound typically occurs prior to the P2 and is a very soft [heart sound](#). In severe AS, the [aortic valve](#) may remain open until later in [diastole](#), causing the P2 heart sound to drown out the A2. Patients with AS often have exertional dyspnea, [paroxysmal nocturnal dyspnea](#), and fatigue. However, AS presents with a harsh crescendo-decrescendo [murmur](#) best heard at the right upper sternal border, not a [holosystolic murmur](#) best heard at the apex.

F - Radiation of the murmur to the axilla

Explanation Why

Radiation of the [murmur](#) to the [axilla](#) is commonly seen in patients with [MR](#). However, radiation of the [murmur](#) to the [axilla](#) is not suggestive of [MR](#) exacerbation.

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

Question # 30

A 2-year-old boy is brought to the physician by his father for a well-child examination. He recently emigrated from Mexico with his family and has not seen a physician since birth. Vital signs are within normal limits. Cardiac examination shows a harsh, grade 3/6 holosystolic murmur heard best at the left lower sternal border. During deep inspiration, the second heart sound is split. If left untreated, irreversible changes would most likely be seen in which of the following structures?

	Answer	Image
A	Right atrium	
B	Ascending aorta	
C	Superior vena cava	
D	Pulmonary artery	
E	Mitral valve	

Hint

A harsh holosystolic murmur heard along the left lower sternal border with normal splitting is suggestive of a ventricular septal defect (VSD), the most common congenital heart defect.

Correct Answer

A - Right atrium

Explanation Why

Irreversible enlargement of the [right atrium](#) can be caused by an untreated [tricuspid regurgitation](#) (e.g., due to [Ebstein anomaly](#)), tricuspid stenosis/atresia, or [atrial septal defect](#) (ASD). [Tricuspid regurgitation](#) would cause a [holosystolic murmur](#) at the left lower sternal border (LLSB), as seen here, but the patient would also have [features of right heart failure](#) (e.g., increased [JVP](#), [hepatomegaly](#), [ascites](#)), unlike this asymptomatic patient. The [murmur](#) due to tricuspid stenosis would also be heard at the LLSB but the [murmur](#) would be a [diastolic](#), not [systolic](#). The [murmur](#) due to an ASD would be a [systolic ejection murmur](#) that is best heard at the upper (not lower!) left sternal border and it would be associated with a fixed [split in S₂](#), unlike this patient with a normal inspiratory split of S₂. This patient most likely has an isolated [VSD](#), which is associated with irreversible changes in a different structure.

B - Ascending aorta

Explanation Why

Irreversible coarctation or atresia of the [ascending aorta](#) is associated with certain congenital disorders such as [bicuspid aortic valve](#) and [hypoplastic left heart syndrome \(HLHS\)](#). A [bicuspid aortic valve](#) can cause an asymptomatic [systolic murmur](#) due to [aortic stenosis](#) but such a [murmur](#) would be a mid-to-late [systolic murmur](#) and it would be best heard at the right upper sternal border ([aortic area](#)). Patients with [HLHS](#) can have a [systolic murmur](#) due to the [atrial septal defect](#) associated with [HLHS](#) but such a [murmur](#) would be a [systolic ejection murmur](#) that is best heard at the upper (not lower!) left sternal border and it would be associated with a fixed [split in S₂](#), unlike this patient with a normal inspiratory split of S₂. Moreover, patients with HLS would manifest with [cyanosis](#) and features of [heart failure](#) shortly after [birth](#), unlike this asymptomatic patient. This patient most likely has an isolated [VSD](#), which does not cause irreversible changes in the [ascending aorta](#).

C - Superior vena cava

Explanation Why

Irreversible stenosis of the [superior vena cava \(SVC\)](#) is a rare congenital condition that manifests with features of [superior vena cava syndrome](#) (e.g., facial [plethora](#), [edema](#) of the face and upper extremities) at [birth](#), not an asymptomatic [murmur](#), as seen here. This patient most likely has a [VSD](#), which does not cause irreversible changes in the [SVC](#).

D - Pulmonary artery


Explanation Why

A [VSD](#) is characterized by a [left-to-right shunt](#), leading to reduced [cardiac output](#) despite increased [left ventricular](#) load and pulmonary blood flow. Oxygenated blood from the [left ventricle](#) re-enters the pulmonary system, increasing [pulmonary artery](#) and [right ventricular](#) pressure, which leads to [hypertrophy](#) in those two areas. The irreversible thickening and narrowing of the pulmonary vasculature cause [pulmonary hypertension](#). In a late stage of the disease, due to irreversible [pulmonary hypertension](#), the [right ventricle](#) pressure increases until it exceeds the [left ventricular](#) pressure, causing a shunt reversal ([Eisenmenger reaction](#)).

E - Mitral valve

Explanation Why

Irreversible [fibrosis](#) and stenosis of the [mitral valve](#) are seen in [rheumatic heart disease](#), which is a chronic complication of [rheumatic fever](#). [Mitral regurgitation](#), which manifests with a [holosystolic murmur](#), is the most common valve defect associated with [rheumatic fever](#) and this patient's migrant background is a [risk factor](#) for [rheumatic fever](#). However, [murmurs](#) due to [mitral valve](#) disease are best heard at the cardiac apex, not at the left lower sternal border, and this child has no features to suggest [rheumatic fever](#) (e.g., [fever](#), history of recent [pharyngitis](#)). Moreover, [rheumatic fever](#) is uncommon in children below the age of 5 years. This patient most likely has a [VSD](#), which is not associated with irreversible changes in the [mitral valve](#).



Question # 31

A 65-year-old woman comes to the physician because of a 1-month history of progressively worsening shortness of breath, dry cough, and fatigue. She has type 2 diabetes mellitus, hyperlipidemia, and coronary artery disease. Current medications include aspirin, metformin, and atorvastatin. Her temperature is 36°C (96.8°F), heart rate is 97/min, and blood pressure is 102/61 mm Hg. Cardiopulmonary examination shows an S₃ gallop and coarse crackles at the lung bases. There is 1+ ankle edema bilaterally. Which of the following sets of hemodynamic changes is most likely in this patient?

	Cardiac output	Pulmonary capillary wedge pressure	Total peripheral resistance	Mixed venous oxygen content
A	Decreased	decreased	decreased	decreased
B	Decreased	increased	increased	decreased
C	Decreased	normal	increased	decreased
D	Decreased	decreased	increased	decreased
E	Increased	decreased	decreased	increased

	Answer	Image
A	A	
B	B	
C	C	
D	D	

	Answer	Image
E	E	

Hint

This patient's dyspnea, peripheral edema, signs of pulmonary edema (coarse pulmonary crackles), and S₃ gallop suggest systolic heart failure. Given her risk factors (type 2 diabetes mellitus, hyperlipidemia, and coronary artery disease), she most likely has underlying ischemic cardiomyopathy.

Correct Answer

A - A

Explanation Why

These parameters are seen in the late stages of untreated [distributive shock](#) (e.g., due to [sepsis](#)). Inflammatory mediators lead to systemic [vasodilation](#) and decreased [peripheral vascular resistance](#), both of which decrease the effective circulating volume. This subsequently results in low [left ventricular end-diastolic volume](#) and decreased [PCWP](#). In the later stages of [shock](#), the [heart](#) can no longer compensate; [cardiac output](#) and [mixed venous oxygen content](#) begin to fall, resulting in combined [shock](#) (e.g., [distributive shock](#) and [cardiogenic shock](#)). However, patients with this set of hemodynamic parameters would be critically ill and present with abnormal vital parameters such as [hypotension](#), [tachycardia](#) or [bradycardia](#), and [fever](#) or [hypothermia](#), none of which is seen here.

B - B

Explanation Why

In [systolic heart failure](#), poor contractility leads to decreased [stroke volume](#) and [cardiac output](#). Because the low [cardiac output](#) does not allow oxygen demand in peripheral tissues to be met, a higher [proportion](#) of oxygen than normal is extracted from the blood, resulting in a lower [mixed venous oxygen content](#). With increased [left ventricular end-diastolic volume](#) and poor forward flow, [left ventricular](#) pressure increases and is transmitted to the [left atrium](#), resulting in increased [PCWP](#). Decreased [cardiac output](#) also results in activation of the [RAAS](#) and [sympathetic nervous system](#), leading to increased [peripheral vascular resistance](#) and enhanced sodium and water retention. Although these changes compensate for low [cardiac output](#) initially, persistent neurohumoral activation further promotes sodium and water retention, cardiac dysfunction, and [myocyte hypertrophy](#).

C - C

Explanation Why

These parameters are consistent with [obstructive shock](#), e.g., in a patient with [massive pulmonary embolism](#) or [pericardial tamponade](#). In [obstructive shock](#), [venous return](#) remains the same but cannot be transmitted through the [heart chambers](#) effectively. This results in decreased [left ventricular end-diastolic volume](#) and [cardiac output](#). Decreased [cardiac output](#) and effective circulating volume activate the [RAAS](#) and [sympathetic nervous system](#), resulting in increased [peripheral vascular resistance](#). In addition, low [cardiac output](#) leads to decreased [mixed venous oxygen content](#). Depending on the extent of obstruction, [PCWP](#) may be decreased or unchanged. Patients with [obstructive shock](#) are most often hypotensive with compensatory [tachycardia](#), neither of which is seen in this patient.

D - D

Explanation Why

These parameters are consistent with [hypovolemic shock](#) (e.g., due to hemorrhage). With decreased circulating blood volume, there is reduced [venous return](#) to the [heart](#), which results in decreased [PCWP](#), [left ventricular end-diastolic volume](#), and [cardiac output](#). Decreased [cardiac output](#) and effective circulating volume activate the [RAAS](#) and [sympathetic nervous system](#), resulting in increased [peripheral vascular resistance](#). In addition, low [cardiac output](#) leads to decreased [mixed venous oxygen content](#). Patients with [hypovolemic shock](#) are most often hypotensive with compensatory [tachycardia](#), neither of which is observed in this patient. Moreover, these hemodynamic changes are unlikely to be seen in this patient, as there is no reported volume depletion.

E - E

Explanation Why

These parameters are seen in the early stages of untreated [septic shock](#). Inflammatory mediators lead to systemic [vasodilation](#) and decreased [peripheral vascular resistance](#), both of which result in

decreased effective circulating volume. This subsequently leads to low [left ventricular end-diastolic volume](#) and decreased [PCWP](#). In the early stages of [sepsis](#), the [heart](#) compensates with hyperdynamic function, resulting in increased [cardiac output](#). Increased blood flow to peripheral tissues decreases the [proportion](#) of O₂ that is extracted from the blood, causing an increase in [mixed venous oxygen content](#). However, patients in the hyperdynamic phase of [septic shock](#) classically present with flushed, warm [skin](#), [fever](#), [hypotension](#), and [bounding pulses](#), none of which is seen here.

Question # 32

A 51-year-old man comes to the physician because of progressive shortness of breath, exercise intolerance, and cough for the past 6 months. He is no longer able to climb a full flight of stairs without resting and uses 3 pillows to sleep at night. He has a history of using cocaine in his 30s but has not used any illicit drugs for the past 20 years. His pulse is 99/min, respiratory rate is 21/min, and blood pressure is 95/60 mm Hg. Crackles are heard in both lower lung fields. An x-ray of the chest shows an enlarged cardiac silhouette with bilateral fluffy infiltrates and thickening of the interlobar fissures. Which of the following findings is most likely in this patient?

	Answer	Image
A	Decreased pulmonary vascular resistance	
B	Decreased lung compliance	
C	Decreased forced expiratory volume	
D	Increased carbon dioxide production	
E	Increased residual volume	

Hint

This patient's history of reduced exercise tolerance and orthopnea, physical examination showing crackles and hypotension, and imaging showing cardiomegaly and pulmonary edema are highly suggestive of congestive heart failure (CHF) due to left ventricular dysfunction.

Correct Answer

A - Decreased pulmonary vascular resistance

Explanation Why

Decreased [pulmonary vascular resistance](#) would not be expected in a patient with [congestive heart failure](#). CHF is characterized by reduced [cardiac output](#) that results in a backup of blood in the pulmonary vasculature. The ensuing pulmonary venous congestion causes an increase in [pulmonary vascular resistance](#).

B - Decreased lung compliance

Explanation Why

Poor [left ventricular](#) function impairs the forward flow of blood and subsequently results in excess blood within the pulmonary vasculature. This excess of blood leads to vascular engorgement, extravasation due to increased [hydrostatic pressure](#), and [pulmonary edema](#). Increased fluid within the [lungs](#) makes the [lungs](#) stiffer, reducing [lung compliance](#).

C - Decreased forced expiratory volume

Explanation Why

[Forced expiratory volume](#) (FEV) is most commonly reduced by [obstructive lung disease](#) (e.g., [asthma](#), [COPD](#)), which can present with [dyspnea](#), exercise intolerance, [cough](#), and an enlarged [cardiac silhouette](#) (due to [right ventricular hypertrophy](#)). However, auscultation would reveal bilateral [wheezes](#) and a chest [x-ray](#) would show features of [pulmonary hyperinflation](#) (e.g., increased [radiolucency](#), flattening of the diaphragm). [Crackles](#) and fluffy infiltrates would not be expected. This patient's presentation is more consistent with [CHF](#), which would not significantly affect [FEV](#).

D - Increased carbon dioxide production

Explanation Why

Increased carbon dioxide production is one of the causes of hypercapnia and may be due to physiologic states (e.g., exercise) or pathologic conditions (e.g., [sepsis](#)). This patient's presentation is consistent with [CHF](#), which would be unlikely to cause an increase in carbon dioxide production.

E - Increased residual volume

Explanation Why

Increased [residual volume](#) (RV) due to [air trapping](#) is seen in [obstructive lung disease](#) (e.g., [asthma](#), [COPD](#)), which can present with [dyspnea](#), exercise intolerance, [cough](#), and an enlarged [cardiac silhouette](#) (due to [right ventricular hypertrophy](#)). However, auscultation would reveal bilateral [wheezes](#) and a chest x-ray would show features of [pulmonary hyperinflation](#) (e.g., increased [radiolucency](#), flattening of the diaphragm). [Crackles](#) and fluffy infiltrates would not be expected. This patient's presentation is more consistent with [CHF](#), which would be associated with a normal or reduced [residual volume](#) as a result of cardiogenic pulmonary edema.

Question # 33

A 62-year-old man comes to the physician for an annual health maintenance examination. He has a history of stable angina, gout, and hypertension. His medications include lisinopril and aspirin. He has smoked a pack of cigarettes daily for 20 years. He drinks 5–6 beers on the weekends. His blood pressure is 150/85 mm Hg. Laboratory studies show a total cholesterol of 276 mg/dL with an elevated low-density lipoprotein (LDL) concentration and low high-density lipoprotein (HDL) concentration. Administration of which of the following agents is the most appropriate next step in treatment?

	Answer	Image
A	Vitamin B ₃	
B	HMG-CoA reductase inhibitor	
C	Peroxisome proliferator-activated receptor alpha activator	
D	Cholesterol absorption inhibitor	
E	Proprotein convertase subtilisin kexin 9 inhibitor	
F	Bile acid resin	

Hint

This patient has hyperlipidemia and a history that suggests coronary artery disease (stable angina).

Correct Answer

A - Vitamin B₃

Explanation Why

[Vitamin B₃ \(niacin\)](#) is a lipid-lowering agent synthesized from [tryptophan](#) that decreases hepatic [VLDL](#) synthesis and inhibits the activity of [hormone-sensitive lipase](#). It increases [HDL](#) levels and, to a lesser degree, reduces [triglyceride](#) and [LDL](#) synthesis. However, it is not considered a first-line monotherapy in patients with [hyperlipidemia](#) and it is not recommended in patients with [gout](#) because of the associated risk of [hyperuricemia](#).

B - HMG-CoA reductase inhibitor

Explanation Why

[HMG-CoA reductase inhibitors](#), more commonly referred to as [statins](#), are the first-line treatment for [hyperlipidemia](#). [Statins](#) inhibit [cholesterol synthesis](#) in the [liver](#) by inhibiting the conversion of [HMG-CoA](#) to [mevalonate](#). [Statins](#) are the most effective drug for reducing [LDL](#) levels, and they also increase [HDL](#) and lower [triglyceride](#) levels. Common side effects of [HMG-CoA reductase inhibitors](#) include elevated [LFTs](#) and myalgias (especially when used concomitantly with [fibrates](#)).

C - Peroxisome proliferator-activated receptor alpha activator

Explanation Why

Peroxisome proliferator-activated receptor alpha (PPAR- α) activators, which are more commonly referred to as [fibrates](#) (e.g., [bezafibrate](#), [fenofibrate](#)), are considered second-line treatment for [hyperlipidemia](#). The activation of PPAR- α increases [lipoprotein lipase](#) production, [fatty acid](#) uptake, and oxidation as well as [HDL](#) synthesis. [Fibrates](#) are most effective at decreasing [triglyceride](#) levels and increasing [HDL](#) levels and only have a mild effect on [LDL](#) levels. For this reason, [fibrates](#) would not be recommended as monotherapy in this patient.

D - Cholesterol absorption inhibitor

Explanation Why

[Cholesterol](#) absorption inhibitors like [ezetimibe](#) decrease [LDL](#) by inhibiting [cholesterol](#) absorption at the brush borders of the [small intestine](#). However, [ezetimibe](#) is considered a second-line lipid-lowering agent. It is reserved for patients who do not tolerate the first-line drug and would not be the first choice for this patient. Side effects of [ezetimibe](#) include [diarrhea](#) and increased [LFTs](#).

E - Proprotein convertase subtilisin kexin 9 inhibitor

Explanation Why

[PCSK9 inhibitors](#) such as [alirocumab](#) and [evolocumab](#) work by inhibiting the degradation of [LDL receptors](#), which decreases circulating [LDL](#) levels. However, they are currently only recommended as an adjunct treatment in patients in whom first-line therapy has failed. Side effects include myalgias, [delirium](#), and cognitive impairment.

F - Bile acid resin

Explanation But

[Bile acid resins](#) are also indicated for the treatment of [cholestasis](#)-associated [pruritus](#).

Explanation Why

[Bile acid resins](#) such as [cholestyramine](#) lower plasma [LDL cholesterol](#) levels by binding to [bile acid](#) and preventing its reabsorption in the [gastrointestinal tract](#), thereby increasing intestinal [LDL](#) losses (which is bound to [bile salts](#)). [Bile acid resins](#) are not considered a first-line therapy for [hyperlipidemia](#) because of their gastrointestinal side effects (e.g., [diarrhea](#)) and the fact they lower [LDL](#) only minimally.

Question # 34

A 51-year-old man comes to the physician because of progressively worsening dyspnea on exertion and fatigue for the past 2 months. Cardiac examination shows no murmurs or bruits. Coarse crackles are heard at the lung bases bilaterally. An ECG shows an irregularly irregular rhythm with absent P waves. An x-ray of the chest shows globular enlargement of the cardiac shadow with prominent hila and bilateral fluffy infiltrates. Transthoracic echocardiography shows a dilated left ventricle with an increased ventricular cavity size; left ventricular ejection fraction is 40%. Which of the following is the most likely cause of this patient's condition?

	Answer	Image
A	Uncontrolled essential hypertension	
B	Chronic supraventricular tachycardia	
C	Deposition of misfolded proteins	
D	Inherited β -myosin heavy chain mutation	
E	Acute psychological stress	

Hint

This patient presents with symptoms of heart failure (pulmonary edema, dyspnea, fatigue) and echocardiography findings that suggest dilated cardiomyopathy (globular enlargement, increased ventricular cavity size).

Correct Answer

A - Uncontrolled essential hypertension

Explanation Why

Uncontrolled [essential hypertension](#) can lead to concentric [left ventricular hypertrophy \(LVH\)](#) due to chronically increased [afterload](#). Individuals with [LVH](#) from [hypertension](#) may eventually also develop [dyspnea](#), fatigue, and [pulmonary edema](#), as well as a decreased [ejection fraction](#). However, this patient's chest [x-ray](#) and [transthoracic echocardiography](#) show [cardiomegaly](#) and an increased ventricular cavity size, which indicates eccentric ventricular dilation.

B - Chronic supraventricular tachycardia

Explanation But

[Tachycardia](#)-induced [dilated cardiomyopathy](#) usually resolves with treatment of the underlying [arrhythmia](#) (typically within 3 months).

Explanation Why

This patient's irregularly irregular rhythm on [ECG](#) suggests chronic [supraventricular tachycardia](#) as the etiology for his current presentation with [heart failure](#). [Supraventricular tachycardia](#) reduces [stroke volume](#), which leads to increased end-[systolic](#) volume, increased wall stress, and subsequent [eccentric hypertrophy](#). [Tachycardia](#)-induced [cardiomyopathy](#) is a form of [dilated cardiomyopathy](#), which leads to reduced [myocardial contractility](#) that results in [heart failure](#), as seen here. Other causes of [dilated cardiomyopathy](#) include infection (e.g., [Coxsackie B myocarditis](#), [Chagas disease](#)), toxic substances (e.g., alcohol, [doxorubicin](#)), and [malnutrition](#) (e.g., [thiamine deficiency](#)).

C - Deposition of misfolded proteins

Explanation Why

Abnormal deposition of misfolded [proteins](#) is the underlying cause of [amyloid cardiomyopathy](#). Patients with [amyloid cardiomyopathy](#) typically present with signs of [diastolic heart failure](#), such as [dyspnea](#) and [pulmonary edema](#), which are also seen here. However, [myocardial amyloid](#) deposition usually causes [restrictive cardiomyopathy](#), whereas this patient's imaging findings are consistent with [dilated cardiomyopathy](#).

D - Inherited β -myosin heavy chain mutation

Explanation Why

Inherited β -[myosin heavy chain](#) mutation is one of the many genetic mutations that lead to primary [hypertrophic cardiomyopathy](#). Individuals with this genetic mutation may eventually develop [dyspnea](#), fatigue, and [pulmonary edema](#), as well as a decreased [ejection fraction](#). However, this patient's chest [x-ray](#) and [transthoracic echocardiography](#) show signs of ventricular dilation, which is not explained by primary [hypertrophic cardiomyopathy](#).


E - Acute psychological stress

Explanation Why

Severe acute psychological stress can lead to transient stress-induced [cardiomyopathy](#) ([Takotsubo cardiomyopathy](#)), which is thought to be caused by a [catecholamine](#) surge. Patients with stress-induced [cardiomyopathy](#) typically present with acute [dyspnea](#) and [chest pain](#) due to [myocardial](#) motion abnormalities. However, in addition to LV dilation, [transthoracic echocardiography](#) would show apical ballooning, which is not seen here. Moreover, this patient's symptoms are progressive and chronic, and there is no history of intense emotional or physical stress.

Question # 35

A 32-year-old man who recently emigrated from Colombia comes to the physician because of a 3-month history of shortness of breath and fatigue. Physical examination shows jugular venous distention and an additional late diastolic heart sound. Crackles are heard at the lung bases bilaterally. Cardiac catheterization is performed and left ventricular pressures are obtained. The left ventricular pressure-volume relationship compared to that of a healthy patient is shown. Which of the following is the most likely cause of this patient's heart failure?



	Answer	Image
A	Chagas heart disease	
B	Viral myocarditis	

	Answer	Image
C	Cardiac sarcoidosis	Chronic sarcoidosis Etiology Multi-system disorder with T cell dysfunction and noncaseating granuloma formation Epidemiology ♀ > ♂; peak age: 20–40 years More common in African American women. More prevalent in Northern than Southern Europe Serology Typically negative for autoantibodies Laboratory studies Soluble interleukin-2 receptor ACE Bronchoalveolar lavage: ↑ CD4/CD8 ratio (often > 5) Complications Bronchiectasis, pulmonary fibrosis, chronic renal failure Prognosis Spontaneous remission in 20%–70% of cases General symptoms Low-grade fever Fatigue Weight loss
D	Alcohol use disorder	
E	Thiamine deficiency	
F	Muscular dystrophy	

Hint

The slope of the pressure-volume curve in the graph for this patient compared to that of a healthy patient shows decreased compliance of the left ventricle (i.e., a decreased ability to expand in volume in response to pressure), indicating diastolic dysfunction.

Correct Answer

A - Chagas heart disease

Explanation Why

Chagas [heart](#) disease is a late complication of infection with [Trypanosoma cruzi](#). It typically occurs 10–20 years after primary disease and manifests as [dilated cardiomyopathy \(DCM\)](#). [DCM](#) manifests with symptoms of [heart failure](#) (e.g., [shortness of breath](#), fatigue), which are seen here. Since [Chagas disease](#) is [endemic](#) to Central and South America, this patient from Colombia is at increased risk for infection. [DCM](#) is characterized by [systolic dysfunction](#) due to decreased ventricular contractility secondary to dilation. Hence, increased [left ventricular](#) volumes but lower pressures compared to a healthy individual would be expected in the pressure-volume curve of a patient with [DCM](#). This patient's graph, however, indicates [diastolic dysfunction](#).


B - Viral myocarditis

Explanation Why

Viral [myocarditis](#) (e.g., caused by [parvovirus B19](#), [human herpesvirus 6](#), [coxsackievirus](#), [echovirus](#), or [HIV](#)) is the most common form of [myocarditis](#). Long-term complications include [dilated cardiomyopathy \(DCM\)](#), which manifests with symptoms of [heart failure](#) such as the [shortness of breath](#) and fatigue seen here. [DCM](#) is characterized by [systolic dysfunction](#) due to decreased ventricular contractility secondary to dilation. Hence, increased [left ventricular](#) volumes but lower pressures compared to a healthy individual would be expected in the pressure-volume curve of a patient with [DCM](#). This patient's graph, however, indicates [diastolic dysfunction](#).

C - Cardiac sarcoidosis

Image



Explanation But

Further manifestations of cardiac [sarcoidosis](#) include [dilated cardiomyopathy](#), conduction abnormalities, ventricular and [supraventricular arrhythmias](#), and valvular dysfunction.

Explanation Why

[Restrictive cardiomyopathy](#) leading to [diastolic dysfunction](#) of the [left ventricle](#) is a manifestation of cardiac [sarcoidosis](#). In this condition, stiffening of the ventricles leads to a defective [diastolic](#) filling while [ejection fraction](#) (i.e., [systolic](#) function) is (nearly) preserved. Patients present with signs and symptoms of heart failure (e.g., [shortness of breath](#), [jugular venous distention](#)). Other causes of [restrictive cardiomyopathy](#) include [amyloidosis](#), [hemochromatosis](#), and postradiation [fibrosis](#).

D - Alcohol use disorder

Explanation Why

Long-term heavy alcohol consumption has a toxic effect on cardiac [myocytes](#) and, therefore, may lead to [dilated cardiomyopathy \(DCM\)](#). This condition manifests with symptoms of [heart failure](#) (e.g., [shortness of breath](#), fatigue), which are seen here. [DCM](#) is characterized by [systolic dysfunction](#) due to decreased ventricular contractility secondary to dilation. Hence, increased [left ventricular](#) volumes but lower pressures compared to a healthy individual would be expected in the pressure-volume curve of a patient with [DCM](#). This patient's graph, however, indicates [diastolic dysfunction](#).

E - Thiamine deficiency

Explanation Why

[Thiamine deficiency](#) (e.g., due to chronic alcohol consumption) can manifest as [wet beriberi](#), a condition characterized by [dilated cardiomyopathy \(DCM\)](#) and [high-output heart failure](#). This patient presents with signs and symptoms of heart failure (e.g., [shortness of breath](#), [jugular venous distention](#)). [DCM](#) is characterized by [systolic dysfunction](#) due to decreased ventricular contractility secondary to dilation. Hence, increased [left ventricular](#) volumes but lower pressures compared to a healthy individual would be expected in the pressure-volume curve of a patient with [DCM](#). This patient's graph, however, indicates [diastolic dysfunction](#).

F - Muscular dystrophy

Explanation Why

Cardiac involvement in [muscular dystrophies](#), such as [Duchenne muscular dystrophy](#) and [Becker muscular dystrophy](#), typically manifests as [dilated cardiomyopathy \(DCM\)](#). [DCM](#) manifests with symptoms of [heart failure](#) (e.g., [shortness of breath](#), fatigue), which are seen here. [DCM](#) is characterized by [systolic dysfunction](#) due to decreased ventricular contractility secondary to dilation. Hence, increased [left ventricular](#) volumes but lower pressures compared to a healthy individual would be expected in the pressure-volume curve of a patient with [DCM](#). This patient's graph,

however, indicates [diastolic dysfunction](#). Moreover, [skeletal muscle](#) weakness would be expected in a patient with [muscular dystrophy](#).

Question # 36

A 75-year-old man comes to the physician because of a 4-month history of progressive shortness of breath and chest pressure with exertion. Cardiac examination shows a crescendo-decrescendo systolic murmur that is heard best in the second right intercostal space. Radial pulses are decreased and delayed bilaterally. Transesophageal echocardiography shows hypertrophy of the left ventricle and a thick, calcified aortic valve. The area of the left ventricular outflow tract is 30.6 mm^2 . Using continuous-wave Doppler measurements, the left ventricular outflow tract velocity is 1.0 m/s , and the peak aortic valve velocity is 3.0 m/s . Which of the following values most closely represents the area of the stenotic aortic valve?

	Answer	Image
A	16.0 mm^2	
B	10.2 mm^2	
C	23 mm^2	
D	6.2 mm^2	
E	2.0 mm^2	

Hint

According to the law of conservation of mass, the volume of fluid flowing into the left ventricular outflow tract per unit time must be equal to the volume of fluid flowing out of the aortic valve per unit time.

Correct Answer

A - 16.0 mm²

Explanation Why

An aortic valve area of 16.0 mm² would not appropriately respect the law of conservation of mass in this patient.

B - 10.2 mm²

Explanation Why

The area of the stenotic [aortic valve](#) is 10.2 mm². Conceptually, the flow on one side of a valve (measured as volume per unit time) must be equal to the flow on the other side of the valve. This equal flow is a representation of the law of conservation of mass; all the blood entering the valve must also be accounted for on the other side of the valve. This concept is expressed mathematically via the continuity equation: $A_1V_1 = A_2V_2$ where subscript 1 denotes blood flow through the left ventricular outflow tract and subscript 2 denotes blood flow through the [aortic valve](#). The velocity in the [left ventricle](#) is 1.0 m/s, while its area is 30.6 mm². As the velocity at the [aortic valve](#) was measured to be 3.0 m/s, the aortic valve area must be $(1 \text{ m/s} \times 30.6 \text{ mm}^2)/3 \text{ m/s} = 10.2 \text{ mm}^2$.

C - 23 mm²

Explanation Why

An aortic valve area of 23.0 mm² would not appropriately respect the law of conservation of mass in this patient.

D - 6.2 mm^2

Explanation Why

An aortic valve area of 6.2 mm^2 would not appropriately respect the law of conservation of mass in this patient.


E - 2.0 mm^2

Explanation Why

An aortic valve area of 2.0 mm^2 would not appropriately respect the law of conservation of mass in this patient.

Question # 37

A 31-year-old man comes to the physician because of a 5-day history of fever, chills, and dyspnea. His temperature is 38.9°C (102°F) and pulse is 90/min. Cardiac examination shows a murmur. In addition to other measures, cardiac catheterization is performed. A graph showing the results of the catheterization is shown. This patient most likely has which of the following valvular heart defects?



	Answer	Image
A	Mitral stenosis	

	Answer	Image
B	Mitral regurgitation	Mitral regurgitation Pressure (mm Hg) Systole Diastole Aorta Left ventricle Left atrium Time
C	Pulmonary regurgitation	
D	Aortic regurgitation	Aortic regurgitation Pressure (mm Hg) Systole Diastole Aorta Left ventricle Left atrium Time
E	Aortic stenosis	Aortic stenosis Pressure (mm Hg) Systole Diastole Aorta Left ventricle Left atrium Time


Hint

The pressure-time graph shows low aortic diastolic pressure and high systolic pressure. This indicates elevated pulse pressure.

Correct Answer

A - Mitral stenosis

Image




Explanation Why

[Mitral stenosis](#) (MS) is often accompanied by a rumbling, delayed [diastolic murmur](#) that is characterized by an [opening snap](#). In contrast to the findings in this patient, [cardiac catheterization](#) would reveal a left atrial pressure that is significantly higher than the [left ventricular](#) pressure during the [ventricular filling](#) phase (i.e., [diastole](#)). This increase in pressure occurs as the [left atrium](#) contracts to fill the [left ventricle](#) across the stenotic [mitral valve](#).

B - Mitral regurgitation

Image



Explanation Why

[Mitral regurgitation \(MR\)](#) is often accompanied by a high-pitched, [holosystolic murmur](#). The characteristic pressure tracing in [MR](#) is a rapid increase in left atrial pressure during ventricular [systole](#) due to the regurgitation of blood from the [left ventricle](#) into the [left atrium](#). However, [cardiac catheterization](#) in this patient does not show an increase in left atrial pressure during [systole](#).


C - Pulmonary regurgitation

Explanation Why

On [cardiac catheterization](#), pulmonary regurgitation is associated with increased end-[diastolic right ventricular](#) pressure due to the increased backflow of blood across the regurgitant [pulmonary valve](#) during [diastole](#). However, this graph shows pressure changes in the left rather than the right [heart](#).

D - Aortic regurgitation

Image




Explanation Why

[Aortic regurgitation](#) (AR) is often a feature of [infectious endocarditis](#), which can manifest with the [fever](#), chills, and [dyspnea](#) seen here. Valvular bacterial vegetations can cause destruction and incompetency of the [aortic valve](#). In AR, regurgitation of blood from the aorta into the [left ventricle](#) (LV) leads to higher end-[diastolic](#) volume and therefore higher [preload](#) and [stroke volume](#), which increases aortic [systolic](#) pressure. [Diastolic](#) aortic pressure is classically decreased because of rapid return of blood to the LV through the incompetent [aortic valve](#). Consequently, there would be an increase in [pulse pressure](#), which can manifest with [water hammer pulse](#) and [pulse](#)-synchronous head nodding ([de Musset sign](#)).

E - Aortic stenosis

Image




Explanation Why


[Aortic stenosis](#) (AS) is often accompanied by a harsh [systolic murmur](#) with a crescendo-decrescendo pattern. [Cardiac catheterization](#) in AS typically shows a peak ventricular pressure that is significantly higher than the peak aortic pressure. This finding is a result of the elevated ventricular pressure required to pump blood through the stenotic [aortic valve](#) before the return to normal [systolic](#) pressure in the aorta. However, [cardiac catheterization](#) in this patient shows a peak ventricular pressure that is identical to the peak aortic pressure.

Question # 38

A 62-year-old man comes to the physician for a 2-month history of progressive shortness of breath during exercise and episodic chest pain. He has no history of serious illness and takes no medications. Vital signs are within normal limits. Cardiac auscultation shows a murmur. Left-heart catheterization is performed. A graph representing the results of the catheterization is shown. The intensity of this patient's cardiac murmur is most likely to be greatest at which of the following labeled points along the timeline?



	Answer	Image
A	A	

	Answer	Image
B	B	 The graph, titled "Aortic stenosis", plots Pressure (mm Hg) on the y-axis (0 to 120) against Time on the x-axis. It shows three pressure curves: Left ventricle (blue), Aorta (orange), and Left atrium (green). The Left ventricle curve shows a normal systolic peak of approximately 120 mm Hg. The Aorta curve shows a lower systolic peak of approximately 100 mm Hg and a prolonged, slow diastolic decay, characteristic of aortic stenosis. The Left atrium curve shows a normal low-pressure waveform. The graph is divided into Systole and Diastole phases.
C	C	
D	D	
E	E	

Hint

The graph shows a peak ventricular pressure that is significantly higher than the peak aortic pressure. This finding is classically seen in aortic stenosis (AS) and can be explained by an elevated ventricular pressure required to pump blood through the stenotic aortic valve.

Correct Answer


A - A

Explanation Why

Point A marks a point in the timeline during the [isovolumetric contraction](#) phase (early [systole](#)) shortly before the [aortic valve](#) opens. However, the characteristic crescendo-decrescendo [murmur](#) in [aortic stenosis](#) will be loudest during the [systolic ejection](#) phase, when there is blood flow through the [aortic valve](#).

B - B

Image



Explanation Why

This patient most likely presents with [aortic stenosis](#) (AS), which is associated with an increased

pressure gradient across the [aortic valve](#) during [ventricular ejection](#). The [ventricular ejection](#) phase starts when the [left ventricular](#) pressure exceeds the aortic pressure and forces the [aortic valve](#) to open. In [aortic stenosis](#), the characteristic harsh crescendo-decrescendo [murmur](#) is caused by the [turbulent blood flow](#) through the narrowed [aortic opening](#). The intensity of this [murmur](#) will be loudest during the [ventricular ejection](#) phase when the aorto-ventricular pressure gradient is highest, which is the point in the timeline marked with B.

C - C

Explanation Why

Point C marks the point in the timeline during which [aortic valve](#) closure occurs, producing the [second heart sound](#) (closure of the pulmonic valve also contributes to this sound). In patients with [aortic stenosis](#) (AS), the [second heart sound](#) is typically soft because of delayed closure and reduced mobility of the [aortic valve](#). However, the characteristic crescendo-decrescendo [murmur](#) in AS will be loudest during the [systolic ejection](#) phase, when there is blood flow through the [aortic valve](#).

D - D

Explanation Why

Point D marks a point in the timeline during the [ventricular filling](#) phase, when blood flows through the [mitral valve](#) into the [left ventricle](#). However, the characteristic crescendo-decrescendo [murmur](#) in AS will be loudest during the [systolic ejection](#) phase, when there is blood flow through the [aortic valve](#).

E - E


Explanation Why

Point E marks the point in the timeline during which the contraction of the [left atrium](#) (i.e., left atrial [systole](#)) occurs and blood is ejected from the [left atrium](#) through the [mitral valve](#) into the [left ventricle](#). Longstanding [aortic stenosis](#) (AS) is associated with [left ventricular hypertrophy](#), which decreases ventricle compliance. As a result, patients with AS may present with a [fourth heart sound](#),

which is caused by late [diastolic](#) contraction of the [left atrium](#) against a noncompliant [left ventricle](#). However, the characteristic crescendo-decrescendo [murmur](#) in AS will be loudest during [systole](#), when there is blood flow through the [aortic valve](#).

Question # 39

A 61-year-old man comes to the physician because of a 3-month history of fatigue and progressively worsening shortness of breath that is worse when lying down. Recently, he started using two pillows to avoid waking up short of breath at night. Examination shows a heart murmur. A graph with the results of cardiac catheterization is shown. Given this patient's valvular condition, which of the following murmurs is most likely to be heard on cardiac auscultation?



	Answer	Image
A	High-pitched, holosystolic murmur that radiates to the axilla	

	Answer	Image
B	Rumbling, delayed diastolic murmur heard best at the cardiac apex	Mitral stenosis The graph for Mitral stenosis shows three pressure curves over time. The y-axis is Pressure (mmHg) from 0 to 120. The x-axis is Time, divided into Systole and Diastole. The Aorta curve (orange) starts at ~90 mmHg, rises to ~120 mmHg at systole, and falls to ~80 mmHg at diastole. The Left atrium curve (green) shows a small 'M' shape with a peak of ~35 mmHg during diastole. The Left ventricle curve (blue) shows a normal systolic peak of ~120 mmHg and a normal diastolic drop to ~0 mmHg.
C	Blowing, early diastolic murmur heard best at the Erb point	Aortic regurgitation The graph for Aortic regurgitation shows three pressure curves over time. The y-axis is Pressure (mm Hg) from 0 to 120. The x-axis is Time, divided into Systole and Diastole. The Aorta curve (orange) starts at ~90 mmHg, rises to ~120 mmHg at systole, and falls to ~60 mmHg at diastole. The Left atrium curve (green) shows a small 'M' shape with a peak of ~20 mmHg during diastole. The Left ventricle curve (blue) shows a normal systolic peak of ~120 mmHg and a normal diastolic drop to ~0 mmHg.
D	High-frequency, diastolic murmur heard best at the 2 nd left intercostal space	
E	Harsh, late systolic murmur that radiates to the carotids	Aortic stenosis The graph for Aortic stenosis shows three pressure curves over time. The y-axis is Pressure (mm Hg) from 0 to 120. The x-axis is Time, divided into Systole and Diastole. The Aorta curve (orange) starts at ~60 mmHg, rises to ~110 mmHg at systole, and falls to ~60 mmHg at diastole. The Left atrium curve (green) shows a small 'M' shape with a peak of ~20 mmHg during diastole. The Left ventricle curve (blue) shows a normal systolic peak of ~120 mmHg and a normal diastolic drop to ~0 mmHg.


Hint

The murmur can be intensified by a handgrip maneuver.

Correct Answer

A - High-pitched, holosystolic murmur that radiates to the axilla

Image




Explanation Why

A high-pitched, [holosystolic murmur](#) that radiates to the [axilla](#) is typically heard in severe cases of [mitral regurgitation](#) (MR). MR can manifest with [dyspnea](#), [cough](#), and other [symptoms of left heart failure](#) (e.g., fatigue, orthopnea, [paroxysmal nocturnal dyspnea](#)). This patient's [cardiac catheterization](#) shows a rapid increase in left atrial (LA) pressure during ventricular [systole](#), a finding that is characteristic of MR. The LA pressure increases as blood enters the [atrium](#) both antegrade through the [pulmonary veins](#) and retrograde from the [left ventricle](#) (LV) through the regurgitant [mitral valve](#).

B - Rumbling, delayed diastolic murmur heard best at the cardiac apex

Image




Explanation Why

A rumbling, delayed [diastolic murmur](#) heard best at the cardiac apex is characteristic of [mitral stenosis](#) (MS). In contrast to the findings in this patient, [cardiac catheterization](#) would reveal a left atrial pressure that is significantly higher than the [left ventricular](#) pressure during the [ventricular filling](#) phase (i.e., [diastole](#)). This increase in pressure occurs as the [left atrium](#) contracts to fill the [left ventricle](#) across the stenotic [mitral valve](#).

C - Blowing, early diastolic murmur heard best at the Erb point

Image



Explanation Why

A blowing, early [diastolic murmur](#) heard best at the [Erb point](#) is a characteristic of [aortic regurgitation](#) (AR). Unlike the pressure changes seen here, [cardiac catheterization](#) in a patient with AR would typically show increased aortic [systolic](#) pressure (regurgitation of blood from the aorta into the LV leads to higher [preload](#) and [stroke volume](#)) and decreased [diastolic](#) aortic pressure (due to rapid return of blood to the LV through the insufficient [aortic valve](#)). Consequently, there would be an increase in [pulse pressure](#), which can manifest with [water hammer pulse](#) and the [de Musset sign](#).

D -


High-frequency, diastolic murmur heard best at the 2nd left intercostal space

Explanation Why

A high-frequency, [diastolic murmur](#) heard best at the second left [intercostal space](#) indicates pulmonary regurgitation. On [cardiac catheterization](#), pulmonary regurgitation is associated with increased end-[diastolic right ventricular](#) pressure due to the increased backflow of blood across the regurgitant [pulmonary valve](#) during [diastole](#). However, this graph shows the pressure changes in the left rather than the right [heart](#).

E - Harsh, late systolic murmur that radiates to the carotids

Image




Explanation Why

A harsh, late [systolic murmur](#) that radiates to the carotids is characteristic of [aortic stenosis](#). Unlike the pressure changes seen here, [cardiac catheterization](#) in AS typically shows a peak ventricular pressure that is significantly higher than the peak aortic pressure. This finding is a result of the elevated ventricular pressure required to pump blood through the stenotic [aortic valve](#) before the return to normal [systolic](#) pressure in the aorta. However, [cardiac catheterization](#) in this patient shows a peak ventricular pressure that is identical to the peak aortic pressure.

Question # 40

An otherwise healthy 27-year-old woman is referred to a cardiologist because of a murmur detected by her general practitioner during a routine health maintenance examination. She has a history of recurrent episodes of sore throat and fever during childhood. She emigrated from Kenya 6 years ago. Cardiac examination shows an opening snap followed by a late diastolic rumble, which is best heard at the fifth intercostal space in the left midclavicular line. Which of the following pressure-volume loop graphs is most likely to correspond to this patient's cardiac condition?



	Answer	Image
A	A	
B	B	
C	C	

	Answer	Image
D	D	
E	E	

Hint

This patient's murmur is characteristic of mitral stenosis. Her history of recurrent fever and sore throat before emigrating from Kenya indicates that she may have had rheumatic fever, a known cause of mitral stenosis.

Correct Answer

A - A

Explanation Why

Graph A indicates [aortic stenosis](#) (AS). In AS, the left-ventricular end-[systolic](#) volume (top left) is greater than normal because outflow is impeded by the narrowed [aortic valve](#), and the end-[diastolic](#) volume (bottom right) is slightly greater than normal due to the increase in [residual volume](#) from the previous [systole](#). The [stroke volume](#) (the width of the pressure-volume loop) is less than normal because the mild increase in end-[diastolic](#) volume does not lead to sufficient [preload](#) to overcome the increased opening resistance of the stenotic valve. However, in mild [aortic stenosis](#), the increased [preload](#) (increased end-[diastolic](#) volume) can result in a more forceful contraction (due to the [Frank-Starling mechanism](#)), such that the [stroke volume](#) is not decreased. The peak [systolic](#) pressure also rises far above normal in order to overcome the high [vascular resistance](#) and [turbulent blood flow](#) created by the stenosed [aortic valve](#). In contrast to this patient's auscultation findings, AS would classically manifest with a harsh crescendo-decrescendo late [systolic murmur](#) best heard over the second right [intercostal space](#).

B - B

Explanation Why

Graph B indicates [mitral regurgitation](#) (MR). The left-ventricular end-[diastolic](#) volume (bottom right) is increased because each subsequent [diastolic](#) filling includes the normal left atrial volume of blood as well as regurgitated blood from the previous cycle. This ventricular volume overload results in increased [preload](#), which increases the [stroke volume](#) (via the [Frank-Starling](#) mechanism). This is apparent in the widened [pressure-volume loop](#). There is no true [isovolumetric contraction](#) phase in the [mitral regurgitation pressure-volume loop](#), as blood flows back to the [left atrium](#) during contraction of the ventricle before the [aortic valve](#) opens (top right). The [stroke volume](#) is divided between the output through the [aortic valve](#) and through the abnormal regurgitant [mitral valve](#), causing a reduced [afterload](#) (↓ outflow resistance) and smaller end-[systolic](#) volume (top left). [MR](#) would present with a high-pitched [holosystolic murmur](#) which is best heard over the apex and radiates to the left [axilla](#), which is not consistent with this patient's auscultation findings.

C - C

Explanation Why

Graph C indicates [aortic regurgitation](#) (AR). The end-[diastolic](#) volume (bottom right) is greater than normal because blood flows into the [left ventricle](#) from both the [left atrium](#) and from the aorta. This ventricular volume overload results in an increased [preload](#), which increases the [stroke volume](#) (via the [Frank-Starling](#) mechanism). This is apparent in the widened [pressure-volume loop](#). There is no true [isovolumetric relaxation](#) phase because, when the [left ventricle](#) begins to relax following [aortic valve](#) closure (top left), blood flows backwards from the aorta into the [left ventricle](#). This, in turn, increases [left ventricular](#) volume prior to the opening of the [mitral valve](#) (bottom left). AR would manifest with an early [diastolic](#) high-pitched decrescendo [murmur](#) best heard along the left sternal border, which is not consistent with this patient's auscultation findings.

D - D

Explanation Why

The pressure-volume loop in graph D indicates [mitral stenosis](#). It shows decreased volume at end-[diastole](#) (bottom right), a direct result of the narrowed [mitral valve](#) opening. According to the [Frank-Starling law](#), a lowered end-[diastolic](#) volume leads to reduced [left ventricular preload](#), resulting in decreased [stroke volume](#) and [cardiac output](#). This is apparent in the narrowed [pressure-volume loop](#). A decrease in [stroke volume](#), in turn, reduces the [afterload](#) imposed on the [left ventricle](#) (via reduction of aortic pressure). This can result in a slightly decreased left-ventricular volume at end-[systole](#) (top left).

E - E


Explanation Why

Graph E indicates ventricular [systolic failure](#). The end-[systolic](#) volume (top left) is greater than normal due to a decreased contractility of the [left ventricle](#). A compensatory increase in end-[diastolic](#) volume (bottom right) elevates [preload](#) and functions to help maintain [stroke volume](#) via the [Frank-Starling](#) mechanism. Although this elevates the [stroke volume](#) so that it is almost normal, [ejection](#)

[fraction](#) is still significantly reduced. Patients with [systolic heart failure](#) typically present with [dyspnea](#), fatigue and edemas, none of which is seen in this patient. Moreover, unlike this patient's auscultatory findings, [heart failure](#) would manifest with an [S3/S4 gallop](#).

Question # 1

A 51-year-old man comes to the physician for 2 months of intermittent low-grade fever, malaise, and joint pain. He has a history of recurrent dental abscesses requiring drainage but has otherwise been healthy. His temperature is 38.3°C (100.9°F) and pulse is 112/min. Physical examination shows a new holosystolic murmur in the left midclavicular line that radiates to the axilla. There are linear reddish-brown lesions underneath the nail beds and tender violaceous nodules on the bilateral thenar eminences. Two sets of blood cultures grow *Streptococcus mutans*. A transthoracic echocardiogram shows moderate regurgitation of the mitral valve. Which of the following mechanisms is most likely directly involved in the pathogenesis of this patient's valvular condition?

	Answer	Image
A	Myxomatous degeneration	
B	Antibody cross-reaction	
C	Sterile platelet thrombi deposition	
D	Leaflet calcification and fibrosis	
E	Coagulative necrosis	
F	Fibrin clot formation	

Hint

The patient meets two of the major (modified) Duke Criteria for infective endocarditis: two blood cultures growing a typical pathogen and evidence of a new valvular regurgitation on transthoracic echocardiogram. The 2-month history of intermittent illness, recurrent dental abscesses, and isolation of *S. mutans* on blood cultures make subacute infective endocarditis the most likely diagnosis.

Correct Answer

A - Myxomatous degeneration

Explanation Why

Myxomatous degeneration of the [mitral valve](#) is the underlying pathology of [mitral valve prolapse \(MVP\)](#). [MVP](#) is a [risk factor](#) for [infective endocarditis](#), which this patient has, but it is not directly involved in the pathogenesis of the disease. It is most often an incidental finding on auscultation in asymptomatic patients. This patient presents with a new [murmur](#) on auscultation, making a pre-existing [MVP](#) unlikely.

B - Antibody cross-reaction

Explanation Why

[Antibody](#) cross-reaction due to [molecular mimicry](#) is the presumed pathogenic mechanism of [rheumatic fever](#) (RF). Acute RF may manifest with [mitral regurgitation](#), [joint pain](#), [skin](#) manifestations, and [fever](#), all of which this patient has. However, RF also manifests with typical [skin](#) findings, such as [erythema marginatum](#) and subcutaneous nodules, which are not seen here. Moreover, this patient has no history of [sore throat](#) or [group A streptococcal infection](#), which typically precede RF. Furthermore, blood cultures are negative in RF as it is an immunologic, non-infectious condition.

C - Sterile platelet thrombi deposition

Explanation Why

[Nonbacterial thrombotic endocarditis \(NBTE\)](#) results from sterile (i.e., noninfectious) [platelet thrombi](#) deposition on the cardiac valves. [NBTE](#) most commonly affects the [mitral valve](#); however, it typically manifests with signs and symptoms of systemic embolization rather than [mitral valve](#) dysfunction, which is seen here. This patient presents with [fever](#) and [bacteremia](#), which is highly suggestive of [infectious endocarditis](#). Moreover, [NBTE](#) usually affects patients with advanced [malignancy](#) or [systemic lupus erythematosus](#), neither of which this patient has.

D - Leaflet calcification and fibrosis

Explanation Why

Leaflet calcification and [fibrosis](#) are pathologic changes seen in [mitral stenosis](#) (MS) due to [rheumatic fever](#) (RF). While MS does produce a [murmur](#) in the left [midclavicular line](#), it is a [diastolic murmur](#) that follows an [opening snap](#). Moreover, this patient has a [TTE](#) with evidence of [mitral regurgitation](#), not stenosis. While acute RF can also manifest with [joint pain](#), [fever](#), and malaise, MS would not be expected until 10 to 20 years after the original illness, and blood cultures would be negative. Finally, this patient presents with [subacute infective endocarditis](#), which individuals with MS are at an increased risk of developing, but calcification and [fibrosis](#) play no direct part in the pathogenesis of this condition.


E - Coagulative necrosis

Explanation Why

[Coagulative necrosis](#) is the mechanism that causes [cell death](#) in [myocardial infarction](#) ([MI](#)). Acute [mitral regurgitation](#), as seen in this patient, can be a complication of [MI](#). However, further symptoms such as left-sided [chest pain](#), [dyspnea](#), nausea, and [diaphoresis](#) would be expected. This patient presents instead with [fever](#), [bacteremia](#), and [skin](#) findings that are highly suggestive of [infectious endocarditis](#).

F - Fibrin clot formation

Image



Explanation But

[Infective endocarditis](#) should always be considered in a patient with a [fever of unknown origin](#).

Explanation Why

[Fibrin](#) clot formation is a key step in the pathogenesis of [subacute infective endocarditis](#). First, [endothelial](#) damage occurs (in native valves this is typically due to [turbulent blood flow](#)), followed by adherence of [platelets](#) and [fibrin](#) to the damaged valve areas. These initially sterile [platelet-fibrin](#) aggregates can become infected by bacteria circulating in the blood, forming vegetations. This patient is at increased risk for [streptococcal bacteremia](#) due to recurrent dental [abscesses](#). Microbial adherence to the [platelet-fibrin](#) nidus is mediated by interaction of [bacterial cell wall](#) components (e.g., [dextrans](#)) and the valvular [endothelium](#). [Fibrin](#) clots encase this vegetation and subsequent destruction of the affected valves leads to valvular insufficiency, as seen here. This patient's [physical examination](#) findings of [splinter hemorrhages](#) and [Osler nodes](#) represent areas of microembolization of vegetation material.

Question # 2

A previously healthy 26-year-old man is brought to the emergency department 30 minutes after collapsing during soccer practice. The patient appears well. His pulse is 73/min and blood pressure is 125/78 mm Hg. Cardiac examination is shown. Rapid squatting decreases the intensity of the patient's auscultation finding. Which of the following is the most likely cause of this patient's condition?

[Video](#)

	Answer	Image
A	Asymmetric hypertrophy of the septum	
B	Fibrinoid necrosis of the mitral valve	
C	Eccentric dilatation of the left ventricle	
D	Calcification of the aortic valve	
E	Dilation of the aortic root	

Hint

This young man has just experienced exertional syncope. He has a harsh crescendo-decrescendo systolic murmur heard best at the left sternal border as well as a fourth heart sound (S4).

Correct Answer

A - Asymmetric hypertrophy of the septum

Explanation Why

[Hypertrophic cardiomyopathy \(HCM\)](#) manifests with an area of increased [left ventricular](#) wall thickness that is typically asymmetrical and involves the septum or anterolateral LV wall. Along with [mitral valve](#) and [papillary muscle](#) abnormalities, septal [hypertrophy](#) can lead to dynamic left ventricular outflow tract obstruction, resulting in exertional [syncope](#) and [systolic ejection murmur](#), both of which are seen in this patient. The intensity of the [murmur](#) associated with [HCM](#) decreases with maneuvers that increase [venous return](#) (e.g., rapid squatting) because they increase [preload](#), which in turn decreases the pressure gradient across the obstruction. This change in [murmur](#) intensity can help differentiate [HCM](#) from other causes of [systolic murmurs](#), such as [ventricular septal defect](#) and [mitral regurgitation](#).

B - Fibrinoid necrosis of the mitral valve

Explanation Why

[Fibrinoid necrosis](#) of the [mitral valve](#) occurs in acute [rheumatic fever](#) (RF), which is typically preceded by a [group A streptococcal pharyngitis](#). Acute RF could lead to significant [mitral regurgitation](#), manifesting as exercise intolerance. However, [mitral regurgitation](#) would cause a pansystolic [murmur](#) heard best at the apex rather than a [systolic ejection murmur](#). Rapid squatting would result in an increased, not decreased, intensity of the [murmur](#) associated with [mitral regurgitation](#). Furthermore, this patient has not had a recent [GAS infection](#) and has had no other features of RF.

C - Eccentric dilatation of the left ventricle

Explanation Why

Significant dilatation of the [left ventricle](#) is characteristic of [dilated cardiomyopathy](#). While patients with [dilated cardiomyopathy](#) may present with exertional dyspnea from [heart failure](#), they are less

likely to experience exertional [syncope](#). Examination may reveal an [S3](#) and/or a pansystolic [murmur](#) of [secondary mitral regurgitation](#) heard best at the apex due to LV dilatation. A prominent [systolic ejection murmur](#), as seen in this patient, would not be expected. Furthermore, rapid squatting would result in an increased, not decreased, intensity of the [murmur](#) associated with [mitral regurgitation](#).

D - Calcification of the aortic valve

Explanation Why

Calcification of the [aortic valve](#) occurs secondary to [aortic valve stenosis](#) (AS). Patients can present with exertional [syncope](#) due to an inability to increase [cardiac output](#), and examination may reveal a harsh [systolic ejection murmur](#), as seen here. However, an AS [murmur](#) is heard best at the right upper sternal border, not the left sternal border. In addition, calcific AS is a disease usually seen in advanced age. Even though individuals with an anatomically abnormal (unicuspid or bicuspid) valve can develop calcific AS one or two decades earlier than those with a normal (tricuspid) valve, this 26-year-old patient is most likely too young to have developed calcific AS.

E - Dilation of the aortic root



Explanation Why

Dilation of the aortic root occurs in individuals with [connective tissue](#) syndromes (e.g., [Marfan syndrome](#)), a [bicuspid aortic valve](#), and [aortic dissection](#). Significant dilatation of the aortic annulus can lead to [aortic regurgitation](#), which results in a decrescendo [diastolic murmur](#) heard best at the left sternal border, in contrast to the [systolic ejection murmur](#) seen here. Furthermore, this patient's condition is not associated with [aortic root dilation](#).



Question # 3

A 48-year-old man comes to the physician because of worsening shortness of breath and nocturnal cough for the past 2 weeks. On two occasions, his cough was bloody. He had a heart condition as a child that was treated with antibiotics. He emigrated to the US from Kazakhstan 15 years ago. Pulmonary examination shows crackles at both lung bases. Cardiac examination is shown. Which of the following is the most likely diagnosis?

[Video](#)

	Answer	Image
A	Aortic valve regurgitation	 The image displays two cardiac monitoring traces over two cardiac cycles. The top trace is an Electrocardiogram (ECG) in blue, showing a regular rhythm with labeled 'Diastole' and 'Systole' phases. The bottom trace is a Phonocardiogram (PCG) in orange, showing heart sounds. In the first cycle, the PCG shows a normal S1 (labeled 'S1') followed by a normal S2 (labeled 'S2'). In the second cycle, the PCG shows a normal S1 (labeled 'S1') followed by a very loud, prolonged, and decrescendo diastolic murmur (labeled 'S2') that extends into the diastole, characteristic of aortic valve regurgitation.
B	Mitral valve prolapse	 The image displays two cardiac monitoring traces over two cardiac cycles. The top trace is an Electrocardiogram (ECG) in blue, showing a regular rhythm with labeled 'Diastole' and 'Systole' phases. The bottom trace is a Phonocardiogram (PCG) in orange, showing heart sounds. In the first cycle, the PCG shows a normal S1 (labeled 'S1') followed by a normal S2 (labeled 'S2'). In the second cycle, the PCG shows a normal S1 (labeled 'S1') followed by a normal S2 (labeled 'S2') with a distinct 'Mid-systolic click' (labeled 'Mid-systolic click') occurring mid-systole, characteristic of mitral valve prolapse.

	Answer	Image
C	Pulmonary valve stenosis	The image displays two cardiac waveforms over three cycles of Diastole and Systole. The top waveform is the Electrocardiogram (ECG), showing a normal QRS complex. The bottom waveform is the Phonocardiogram, showing the first heart sound (S1) and second heart sound (S2). In pulmonary valve stenosis, the S2 is significantly reduced in intensity, and there is a prominent, prolonged, and decelerating diastolic murmur (S2 to S1). The background is shaded to indicate Diastole (light blue) and Systole (light orange).
D	Mitral valve stenosis	The image displays two cardiac waveforms over three cycles of Diastole and Systole. The top waveform is the Electrocardiogram (ECG), showing a normal QRS complex. The bottom waveform is the Phonocardiogram, showing the first heart sound (S1) and second heart sound (S2). In mitral valve stenosis, the S2 is normal in intensity, and there is a prominent, prolonged, and decelerating diastolic murmur (S2 to S1). The background is shaded to indicate Diastole (light blue) and Systole (light orange).
E	Mitral valve regurgitation	The image displays two cardiac waveforms over three cycles of Diastole and Systole. The top waveform is the Electrocardiogram (ECG), showing a normal QRS complex. The bottom waveform is the Phonocardiogram, showing the first heart sound (S1) and second heart sound (S2). In mitral valve regurgitation, the S2 is normal in intensity, and there is a prominent, prolonged, and decelerating diastolic murmur (S2 to S1). The background is shaded to indicate Diastole (light blue) and Systole (light orange).

	Answer	Image
F	Aortic valve stenosis	 The image displays two cardiac waveforms for Aortic valve stenosis. The top waveform is an Electrocardiogram (ECG) in blue, and the bottom is a Phonocardiogram (PCG) in orange. The background is divided into Diastole (grey) and Systole (white) phases. The ECG shows a narrow QRS complex. The PCG shows a high-pitched, late-systolic murmur that begins after the S2 heart sound. The heart sounds S1 and S2 are labeled at the bottom of the PCG trace.
G	Tricuspid valve stenosis	 The image displays two cardiac waveforms for Tricuspid valve stenosis. The top waveform is an Electrocardiogram (ECG) in blue, and the bottom is a Phonocardiogram (PCG) in orange. The background is divided into Diastole (grey) and Systole (white) phases. The ECG shows a narrow QRS complex. The PCG shows a high-pitched, early-diastolic murmur that begins after the S2 heart sound. The heart sounds S1 and S2 are labeled at the bottom of the PCG trace.


Hint

This patient's heart problem is consistent with a history of rheumatic fever. An x-ray of the chest in this patient could show left atrial enlargement as a result of this patient's condition.

Correct Answer

A - Aortic valve regurgitation

Image




Explanation Why

Although [aortic regurgitation](#) may cause signs of [pulmonary edema](#) (orthopnea, [hemoptysis](#), [crackles](#)), it causes a [diastolic](#) decrescendo [murmur](#) heard best parasternally at the second [intercostal space](#), and may sometimes be accompanied by an [Austin Flint murmur](#).

B - Mitral valve prolapse

Image




Explanation Why

[Mitral valve prolapse](#) is the most common valvular abnormality in developed countries and is often caused by [rheumatic fever](#). However, it is usually asymptomatic and is best heard at the fifth [intercostal space](#) at the [midclavicular line](#) as a mid-[systolic](#) high-frequency click.

C - Pulmonary valve stenosis

Image




Explanation Why

[Pulmonary valve stenosis](#) is usually congenital and causes a [systolic murmur](#) best heard at the second left [intercostal space](#) parasternally. It can radiate to the back and also be present as a widely [split second heart sound](#), but this is not consistent with this patient's findings.

D - Mitral valve stenosis

Image



Explanation Why

This patient's [diastolic murmur](#) at the fifth [intercostal space](#) and [opening snap](#) indicate [mitral stenosis](#), which causes an increase in left atrial pressure (and $\uparrow$ [PCWP](#)) and a backup of blood into [lungs](#). The volume overload eventually causes symptoms of cardiogenic pulmonary edema, which explain this patient's orthopnea, [hemoptysis](#), and [crackles](#). This patient most likely had [rheumatic fever](#) as a child, the most common cause of [mitral stenosis](#).

E - Mitral valve regurgitation

Image




Explanation Why

[Mitral regurgitation](#) may cause signs of [pulmonary edema](#) (orthopnea, [hemoptysis](#), [crackles](#)). However, it presents as a [holosystolic murmur](#) best heard at the apex of the fifth left [intercostal space](#), in contrast to this patient's [diastolic murmur](#).

F - Aortic valve stenosis

Image




Explanation Why

Although severe [aortic valve stenosis](#) may eventually cause signs of [pulmonary edema](#) (orthopnea, [hemoptysis](#), [crackles](#)), patients would most likely first present with [syncope](#) or [angina](#). Furthermore, [aortic stenosis](#) causes a harsh [systolic](#) crescendo-decrescendo [murmur](#) best heard at the second right [intercostal space](#), which can radiate to the carotids.

G - Tricuspid valve stenosis

Image



Explanation Why

Tricuspid valve stenosis is extremely rare and can be caused by [rheumatic fever](#). However, it causes a [diastolic murmur](#) best heard at the left fourth [intercostal space](#) parasternally and is usually associated with signs of [right heart failure](#) (e.g., peripheral [edema](#), [jugular venous distention](#)).

Question # 4

A 33-year-old pilot is transported to the emergency department after she was involved in a cargo plane crash during a military training exercise in South Korea. She is conscious but confused. She has no history of serious illness and takes no medications. Physical examination shows numerous lacerations and ecchymoses over the face, trunk, and upper extremities. The lower extremities are cool to the touch. There is continued bleeding despite the application of firm pressure to the sites of injury. The first physiologic response to develop in this patient was most likely which of the following?

	Answer	Image
A	Increased heart rate	
B	Increased production of lactic acid	
C	Decreased urine output	
D	Increased firing of chemoreceptors	
E	Increased capillary refill time	
F	Decreased systolic blood pressure	
G	Increased respiratory rate	

Hint

This patient's history of trauma, numerous lacerations and ecchymoses, cool extremities, and continued bleeding, despite the application of pressure, are concerning for hemorrhagic shock.

Correct Answer

A - Increased heart rate

Explanation Why

An increase in [heart rate](#) is the first compensatory mechanism for [hypovolemic shock](#) and may be seen as early as class I hemorrhagic shock. In this patient, blood loss has resulted in [hypovolemia](#) and decreased [stroke volume](#). The [heart rate](#) increases (as a result of intracardiac and arterial [baroreceptor reflexes](#)) to help maintain [cardiac output](#). Initially, this mechanism may be sufficient to compensate for volume loss; patients in class I hemorrhagic shock may be otherwise asymptomatic. As [shock](#) progresses, compensatory mechanisms become overwhelmed by volume loss, resulting in further derangements in vital signs (e.g., [tachycardia](#), [hypotension](#), [tachypnea](#), [hypoxemia](#)) and damage to multiple organ systems.

B - Increased production of lactic acid

Explanation Why

Increased production of [lactic acid](#) occurs as a result of the body's inability to meet tissue demand for oxygen. This occurs in the later phase of [shock](#); it is not the first physiologic response. Blood loss in hemorrhagic shock results in poor tissue perfusion. Consequently, cells are unable to maintain [aerobic metabolism](#) and revert to [anaerobic metabolism](#), a less efficient mechanism that causes increased production of [lactic acid](#). Blood is simultaneously shunted away from the extremities to more vital organs, such as the [heart](#) and brain, further decreasing the supply of oxygen to peripheral tissues.

C - Decreased urine output

Explanation Why

In response to the [hypovolemic](#) state of hemorrhagic shock, [urine](#) output decreases in an attempt to preserve the remaining circulatory volume. However, this physiologic response does not occur until class II of hemorrhagic shock and, therefore, would not be the first one to occur in this patient.

D - Increased firing of chemoreceptors

Explanation Why

Increased firing of [chemoreceptors](#) occurs in patients with hemorrhagic shock in response to [hypoxia](#) of the carotid body and [metabolic acidosis](#) (caused by decreased tissue perfusion and increased [lactic acid](#) production). The chemoreceptor reflex augments the [baroreceptor reflex](#) in activating the [sympathetic nervous system](#). However, this occurs in the later phases of [shock](#); it is not the first physiologic response.

E - Increased capillary refill time

Explanation Why

Peripheral vasoconstriction is an early response to [hypovolemia](#), which causes decreased tissue perfusion and leads to an increased [capillary refill time](#). However, the peripheral [capillaries](#) of the [skin](#) and the [nails](#) are not typically involved until class II of hemorrhagic shock. Therefore, increased [capillary refill time](#) is not the first physiologic response to occur in this patient.

F - Decreased systolic blood pressure

Explanation Why

Hemorrhagic shock results in the depletion of intravascular volume. However, because of several compensatory mechanisms, the body is initially able to maintain blood pressure and tissue perfusion via peripheral vasoconstriction, which may lead to a narrowed [pulse pressure](#); a decreased [systolic](#) blood pressure is not the first physiologic response. As [shock](#) progresses (class III hemorrhagic shock), the compensatory mechanisms are overwhelmed by [hypovolemia](#), and the [systolic](#) blood pressure decreases.


G - Increased respiratory rate

Explanation Why

The [respiratory rate](#) increases in patients with hemorrhagic shock because [tachypnea](#) helps to alleviate the [metabolic acidosis](#) caused by decreased tissue perfusion and increased [lactic acid](#) production. However, this physiologic response does not occur until class II of hemorrhagic shock and, therefore, would not be the first one to occur in this patient.

Question # 5

A 54-year-old man is brought to the emergency department after having been hit by a car while riding his bicycle. He was not wearing a helmet. Despite appropriate life-saving measures, he dies 2 hours later because of a severe intracranial hemorrhage. Autopsy of the heart shows general thickening of the left ventricular wall with narrowing of the ventricular chamber. Which of the following conditions is the most likely underlying cause of the described cardiac changes?

	Answer	Image
A	Aortic regurgitation	
B	Chronic heavy drinking	
C	Myocardial ischemia	
D	Chronic hypertension	
E	Hemochromatosis	
F	Sarcoidosis	

Hint

Thickening of the ventricular wall with narrowing of the ventricular chamber is caused by concentric ventricular hypertrophy.

Correct Answer

A - Aortic regurgitation

Explanation Why

[Aortic regurgitation](#) causes backflow of blood into the [left ventricle](#) during [diastole](#), resulting in an increased [preload](#). The increased blood volume in the [left ventricle](#) triggers myocardial remodeling, specifically the addition of [sarcomeres](#) in series. This results in eccentric (not concentric) [hypertrophy](#), eventually leading to [dilated cardiomyopathy](#).

B - Chronic heavy drinking

Explanation Why

Chronic heavy drinking can result in [dilated cardiomyopathy](#), which is caused by eccentric ventricular [hypertrophy](#), not the concentric ventricular [hypertrophy](#) seen in this patient. The pathogenesis of alcohol-induced [cardiomyopathy](#) is not entirely understood, but it occurs in part due to the direct toxic effects of alcohol and its metabolites (e.g., acetaldehyde) on [myocytes](#).


C - Myocardial ischemia

Explanation Why

[Myocardial ischemia](#) is the cause of [ischemic cardiomyopathy](#), which results in the replacement of the [myocytes](#) with [scar](#) tissue. The lack of contractile tissue in [myocardial fibrosis](#) causes a [systolic](#) impairment, which results in a reduced [ejection fraction](#) (EF). Cardiac [preload](#) is increased to compensate for the reduced EF (via the [Frank Starling mechanism](#)), which results in [sarcomeres](#) being duplicated in series and eccentric ventricular [hypertrophy](#).

D - Chronic hypertension

Image



Explanation But

Other causes of concentric [left ventricular hypertrophy](#) are [aortic valve stenosis](#) and [hypertrophic cardiomyopathy](#).

Explanation Why

Chronic [hypertension](#) is the most common cause of concentric [left ventricular hypertrophy](#). [Hypertension](#) causes increased [afterload](#), resulting in elevated [left ventricular](#) wall stress. When this increased wall stress is sustained, e.g., in chronic [hypertension](#), [gene expression](#) is altered in muscle cells, leading to an increased number of [sarcomeres](#). When responding to increased [afterload](#), [sarcomeres](#) will duplicate in parallel, resulting in [concentric hypertrophy](#) and the thickened ventricular wall seen in this patient. A sustained increase in [preload](#), on the other hand, will result in [eccentric hypertrophy](#) with an associated dilation of the ventricular chamber.

E - Hemochromatosis

Explanation Why

[Hemochromatosis](#) is associated with both dilated and [restrictive cardiomyopathy](#). During the early stages of cardiac [hemochromatosis](#), a restrictive pattern of [left ventricular](#) filling is seen due to [iron](#) deposition in [cardiac muscle](#). As the disease progresses, [systolic dysfunction](#) occurs, which leads to [dilated cardiomyopathy](#) (due to [eccentric hypertrophy](#)). However, cardiac [hemochromatosis](#) does not result in concentric ventricular [hypertrophy](#).

F - Sarcoidosis

Explanation Why

[Sarcoidosis](#) can manifest with either dilated or [restrictive cardiomyopathy](#), both of which are due to granulomatous infiltration of the cardiac [myocardium](#). Clinical features of cardiac [sarcoidosis](#) depend on the degree of [myocardial](#) granulomatous infiltration. Cardiac [sarcoidosis](#) is most commonly asymptomatic but can also result in [arrhythmias](#) and conduction defects. Cardiac [sarcoidosis](#) does not cause concentric ventricular [hypertrophy](#).

Question # 6

A 34-year-old woman, gravida 4, para 3, comes to the physician because of left ankle swelling for 2 months. She notes that the swelling is present throughout the day and decreases when she goes to sleep. One year ago, she has had an episode of deep venous thrombosis after the delivery of her third child. Her prepregnancy BMI was 34 kg/m^2 . Examination shows distended, tortuous veins in the legs bilaterally and pitting edema of the left ankle. There are erythematous scaling patches on the medial side of the left ankle. Duplex ultrasonography is performed. Which of the following directions of blood flow would most likely confirm the diagnosis?

	Answer	Image
A	Anterior tibial vein to popliteal vein	
B	Popliteal vein to small saphenous vein	
C	Femoral vein to external iliac vein	
D	Great saphenous vein to femoral vein	
E	Dorsal venous arch to great saphenous vein	

Hint

Distended and tortuous veins in both legs, ankle edema, and skin changes on the medial ankle indicate chronic venous disease.

Correct Answer


A - Anterior tibial vein to popliteal vein

Explanation Why

Venous blood flowing from the [anterior](#) tibial [vein](#) to the [popliteal vein](#) indicates normal blood flow from a peripheral deep vein to a deep vein. [Chronic venous disease](#), however, is characterized by retrograde blood flow, or venous reflux from the deep veins into the superficial veins.

B - Popliteal vein to small saphenous vein

Image



Explanation Why

Venous blood flowing from the deep [popliteal vein](#) to the [superficial small saphenous vein](#) indicates retrograde blood flow and confirms the diagnosis of [chronic venous disease](#). Normally, the [small](#)

[saphenous vein](#) would arise from the [dorsal](#) venous arch and drain into the [popliteal vein](#) at the [saphenopopliteal junction](#). In a patient with [chronic venous disease](#), however, incompetence of the venous valves caused by elevated venous pressure (e.g., due to [pregnancy](#), increased [BMI](#), prior thrombosis) leads to blood reflux from deep into superficial veins. This results in the formation of [varicose veins](#), and ultimately [edema](#) and [skin](#) changes (e.g., [stasis dermatitis](#)), as seen in this patient. First-line treatment consists of conservative measures such as leg elevation, compression therapy, and application of topical dermatologic agents.

C - Femoral vein to external iliac vein

Explanation Why

Venous blood flowing from the [femoral vein](#) to the external iliac vein indicates normal blood flow from a deep vein to a [proximal](#) deep vein. [Chronic venous disease](#), however, is characterized by retrograde blood flow, or venous reflux from the deep veins into the superficial veins.

D - Great saphenous vein to femoral vein

Explanation Why

Venous blood flowing from the [great saphenous vein](#) to the [femoral vein](#) indicates normal blood flow from a [superficial](#) to a deep vein. [Chronic venous disease](#), however, is characterized by retrograde blood flow, or venous reflux from the deep veins into the superficial veins.

E - Dorsal venous arch to great saphenous vein

Explanation Why

The [dorsal](#) venous arch is a superficial vein that gives rise to the [great saphenous vein](#) and [small saphenous vein](#), which are superficial veins that originate from the [medial](#) and [lateral](#) sides of the [dorsal](#) venous arch, respectively. Therefore, blood normally flows from the [dorsal](#) venous arch to the [great saphenous vein](#). [Chronic venous disease](#), however, is characterized by retrograde blood flow, or venous reflux from the deep veins into the superficial veins.

Question # 7

A 52-year-old woman comes to the physician because of a 1-month history of mild fever, fatigue, and shortness of breath. She has no history of serious medical illness and takes no medications. Cardiopulmonary examination shows a mid-diastolic plopping sound heard best at the apex and bilateral rales at the base of the lungs. Echocardiography shows a pedunculated, heterogeneous mass in the left atrium. A biopsy of the mass shows clusters of mesenchymal cells surrounded by gelatinous material. Further evaluation of this patient is most likely to show which of the following?

	Answer	Image
A	Increased IL-6 serum concentration	
B	Increased S100 protein serum concentration	
C	Axillary lymphadenopathy	
D	Malignant pleural effusion	
E	Ash-leaf skin lesions	

Hint

This patient's echocardiography and biopsy findings are characteristic of cardiac myxoma.

Correct Answer

A - Increased IL-6 serum concentration

Explanation But

[Cardiac myxomas](#) are the most common primary [tumor](#) of the [heart](#).

Explanation Why

This patient's [fever](#) is most likely caused by [IL-6](#), a [cytokine](#) that stimulates the production of [acute phase reactants](#) and is secreted by [cardiac myxomas](#). This patient's cardiopulmonary symptoms (fatigue, [dyspnea](#)) and examination findings (mid-[diastolic murmur](#), bilateral [rales](#)) are due to [left atrium](#) obstruction and resultant [pulmonary edema](#). The treatment for [cardiac myxoma](#) is surgical resection.

B - Increased S100 protein serum concentration

Explanation Why

Increased [S100 protein](#) serum concentration is found in [malignant melanoma](#). While [melanomas](#) are likely to [metastasize](#) to the [heart](#), the pedunculated appearance seen in this patient's [echocardiography](#) and biopsy findings indicates [cardiac myxoma](#), a primary cardiac [tumor](#).

C - Axillary lymphadenopathy

Explanation Why

Axillary [lymphadenopathy](#) is seen in several conditions, including [malignancy](#). [Malignancy](#) and subsequent cardiac [metastases](#), which are the most common cardiac tumors, may explain this patient's [echocardiography](#) findings and symptoms of left atrial obstruction. However, the pedunculated appearance of the [tumor](#) on [echocardiography](#) and biopsy findings indicate [cardiac myxoma](#), which is typically benign and would not cause axillary [lymphadenopathy](#).

D - Malignant pleural effusion

Explanation Why

[Malignant pleural effusion](#) occurs with several cancers (e.g., [lung cancer](#), [breast cancer](#)), some of which may [metastasize](#) to the [heart](#). However, this patient's [echocardiography](#) and biopsy findings are consistent with [cardiac myxoma](#), which is typically benign. Furthermore, malignant pulmonary effusion typically manifests unilaterally; this patient's bilateral pulmonary [rales](#) are more likely due to [left atrium](#) obstruction and resultant [pulmonary edema](#).


E - Ash-leaf skin lesions

Explanation Why

Ash-leaf [skin](#) lesions are a feature of [tuberous sclerosis](#), which may also manifest with cardiac [rhabdomyoma](#). While this patient's cardiopulmonary symptoms and [echocardiography](#) findings are consistent with a cardiac [tumor](#), the biopsy findings are characteristic of [cardiac myxoma](#) rather than [rhabdomyoma](#). Furthermore, [tuberous sclerosis](#) typically manifests during childhood, often with additional neurocognitive symptoms (e.g., [intellectual disability](#), [seizures](#)) due to tumors in the [central nervous system](#).

Question # 8

A 42-year-old woman comes to the physician because of frequent episodes of headaches and tinnitus over the past 3 months. One week ago, she had a brief episode of left arm weakness and numbness that lasted for 2 minutes before spontaneously resolving. She is otherwise healthy and takes no medications. She has smoked one-half pack of cigarettes daily for 22 years. Her pulse is 84/min and blood pressure is 155/105 mm Hg. Abdominal examination shows no masses or tenderness. A bruit is heard on auscultation of the abdomen. Abdominal ultrasonography shows a small right kidney. CT angiography shows stenosis of the distal right renal artery. Which of the following is the most likely underlying cause of the patient's condition?

	Answer	Image
A	Fibromuscular dysplasia	
B	Polyarteritis nodosa	
C	Atherosclerotic plaques	
D	Congenital renal hypoplasia	
E	Systemic lupus erythematosus	

Hint

A characteristic “string of beads” appearance of the renal artery on digital subtraction angiography confirms the diagnosis.

Correct Answer

A - Fibromuscular dysplasia

Image



Explanation But

[FMD](#) is treated with [antihypertensives](#) (e.g., ACE inhibitors), [antiplatelet drugs](#) for [stroke](#) prevention, and revascularization (e.g., percutaneous transluminal angioplasty, [bypass surgery](#)).

Explanation Why

[Fibromuscular dysplasia \(FMD\)](#) is an idiopathic, noninflammatory, nonatherosclerotic developmental defect of the blood vessel wall that causes both narrowing (stenosis) and bulging ([aneurysms](#)) of medium-sized [arteries](#) (leading to a “string of beads” appearance on angiography), both of which can lead to tissue [ischemia](#). The most frequently affected [arteries](#) are the [internal carotid](#) and (mid or [distal](#)) [renal arteries](#), leading to features of cerebrovascular FMD (e.g., [headaches](#), pulsatile [tinnitus](#), [transient ischemic attacks](#)) and/or renal [FMD](#) (e.g., [renal hypertension](#), abdominal [bruit](#), renal [atrophy](#)), as seen in this patient.

B - Polyarteritis nodosa

Explanation Why

[Polyarteritis nodosa \(PAN\)](#) is a small and [medium-sized vessel vasculitis](#) that results in tissue [ischemia](#) and can involve both the [kidneys](#) and the nervous system. [PAN](#) may also manifest with symptoms such as [hypertension](#), renal [atrophy](#), [headaches](#), and [transient ischemic attacks](#), as seen in this patient. However, [PAN](#) usually manifests with features of systemic disease (e.g., [fever](#), fatigue, weight loss, arthralgia) as well as additional symptoms depending on the organs involved (e.g., [polyneuropathy](#), [skin](#) lesions, abdominal [pain](#)), none of which are present here. Moreover, [PAN](#) most commonly affects men and is often associated with [hepatitis B](#) or [hepatitis C](#) infection. Renal angiography characteristically shows numerous microaneurysms and spasms within the affected vessels.

C - Atherosclerotic plaques

Explanation Why

[Atherosclerosis](#) is the most common cause of both renal and [carotid artery stenosis](#), which can also manifest with [renal hypertension](#), abdominal [bruit](#), renal [atrophy](#), [headaches](#), [tinnitus](#), and [transient ischemic attacks](#). This patient's 11-[pack-year](#) smoking history increases her risk of developing [atherosclerosis](#). However, she lacks further known [risk factors](#), such as advanced age, [obesity](#), [dyslipidemia](#), [diabetes mellitus](#), and chronic [hypertension](#). Moreover, [atherosclerotic plaques](#) resulting in stenosis are typically found in the [proximal](#) $\frac{1}{3}$ of the [renal artery](#) as opposed to the [distal](#) involvement seen here.

D - Congenital renal hypoplasia

Explanation Why

[Hypertension](#) and a small right [kidney](#) raise suspicion for congenital [renal hypoplasia](#). However, unilateral [renal hypoplasia](#) is usually asymptomatic and would not explain this patient's neurological symptoms. Moreover, angiography would show a proportionally small [renal artery](#) and arterial branches instead of the [distal renal artery stenosis](#) seen here.

E - Systemic lupus erythematosus

Explanation Why

Neurological complaints, renal, and vascular disease in a woman of childbearing age raise suspicion for [systemic lupus erythematosus \(SLE\)](#). However, in the absence of [antiphospholipid syndrome](#), [SLE](#) does not (typically) manifest with [renal artery stenosis](#). Moreover, this patient lacks other characteristic findings of [SLE](#) such as fatigue, [malar rash](#), polyarthritis/[polyarthralgia](#), symptoms of serositis (e.g., [chest pain](#)), or hematologic disorders (e.g., [anemia](#), [thrombocytopenia](#)).

Question # 9

A 46-year-old woman comes to the emergency department because of a 1-hour history of severe pain, numbness, and weakness in her right lower leg. One week ago, she had an episode of syncope that was preceded by severe palpitations. She has no history of major medical illness and takes no medications. Pulse is 110/min and irregular. Blood pressure is 124/82 mm Hg. The right leg below the knee is pale and cold to touch. Popliteal and pedal pulses are absent on the right. Which of the following is the most likely cause of the patient's symptoms?

	Answer	Image
A	Atherosclerotic narrowing of the artery	
B	Arterial vasospasm	
C	Atheroembolism	
D	Arterial embolism	
E	Popliteal artery aneurysm	
F	Venous thrombosis	

Hint

This patient presents with features of acute lower limb ischemia: pain, pallor, pulselessness, paresthesia (numbness), and poikilothermia.

Correct Answer

A - Atherosclerotic narrowing of the artery

Explanation Why

Atherosclerotic narrowing of the peripheral [arteries](#), the most common cause for [peripheral artery disease](#), would manifest with [intermittent claudication](#) and/or chronic [rest pain](#), not [acute limb ischemia](#). [Acute limb ischemia](#) would only occur if the [atherosclerotic plaque](#) ruptured, causing thrombosis within the vessel lumen. This patient does not have [risk factors for atherosclerosis](#) (e.g., [diabetes mellitus](#), [hypertension](#), [dyslipidemia](#), smoking, advanced age).

B - Arterial vasospasm

Explanation Why

Arterial vasospasm due to drugs such as ergots, [LSD](#), and [cocaine](#) can cause [acute limb ischemia \(ALI\)](#), but a different cause of [ALI](#) is much more likely in this patient with no history of drug use.

C - Atheroembolism

Explanation Why

[Atheroembolism](#) ([cholesterol embolization syndrome](#)), which is the embolization of [cholesterol](#) crystals from [atherosclerotic plaques](#), typically causes [microvascular ischemia](#), resulting in insidious organ damage (especially with renal involvement) and/or [skin](#) changes (e.g., [livedo reticularis](#), [blue toe syndrome](#)); [acute limb ischemia](#) due to [atheroembolism](#) is rare. Also, this patient does not have [risk factors for atherosclerosis](#) (e.g., advanced age, [diabetes mellitus](#), [hypertension](#), [dyslipidemia](#)). Moreover, [atheroembolism](#) is usually provoked by a vascular intervention (e.g., [cardiac catheterization](#), angiography) or [anticoagulant](#) therapy; this patient has no history of medical intervention.

D - Arterial embolism

Explanation Why

This patient presents with [acute limb ischemia \(ALI\)](#), which is commonly caused by arterial embolism. This patient's [syncope](#), [palpitations](#), and [tachycardia](#) with an irregular [pulse](#) suggest [atrial fibrillation](#). Blood stasis in the [atria](#) (in [atrial fibrillation](#)) or in the ventricles (in [ventricular aneurysm](#)) results in the formation of thrombi, which can embolize into the [systemic circulation](#), thereby occluding the peripheral [arteries](#). Other common causes of [acute limb ischemia](#) include thrombosis of a previously patent [artery](#), dissection, trauma, and stent or graft occlusion.

E - Popliteal artery aneurysm

Explanation Why

[Popliteal artery aneurysm](#) can manifest with [acute limb ischemia](#) if sudden thrombosis of the [aneurysm](#) or [distal](#) embolization occurs. This patient does not have [risk factors](#) for a [popliteal aneurysm](#) (e.g., male [gender](#), advanced age, smoking). Moreover, a patient with a [popliteal artery aneurysm](#) would have a pulsatile mass in the [popliteal fossa](#), whereas this patient has a nonpalpable popliteal [pulse](#).

F - Venous thrombosis

Explanation But

[DVT](#) can result in acute limb [ischemia](#), as seen here, as a result of [paradoxical embolism](#) (due to [patent foramen ovale](#)) or [compartment syndrome](#) (phlegmasia alba dolens). However, both of these scenarios are extremely rare.

Explanation Why

Acute [deep vein thrombosis \(DVT\)](#) of the lower limb can manifest with sudden unilateral leg [pain](#). However, it typically also manifests with swelling, [erythema](#), and warmth, not the pallor, muscle

weakness, numbness, absent pulses, or coldness to touch seen here. Furthermore, this patient does not have [risk factors](#) for a [DVT](#) (age > 60 years, [malignancy](#), [thrombophilia](#), and prolonged immobilization).

Question # 10

A 65-year-old man is brought to the emergency room 30 minutes after an acute onset of chest pain and shortness of breath. He appears anxious and diaphoretic. His temperature is 36.9°C (98.4°F), pulse is 110/min, respirations are 16/min, and blood pressure is 115/78 mm Hg. Physical examination shows an S4 gallop. ECG shows ST-segment elevation in leads I, aVL, and V₁-V₆. Emergent percutaneous coronary angiography shows complete occlusion of the left main coronary artery. The release of which of the following substances from platelets is most likely involved in the pathogenesis of this patient's angiography findings?

	Answer	Image
A	Prostaglandin E ₁ (PGE ₁)	
B	Leukotriene D ₄ (LTD ₄)	
C	Nitric oxide (NO)	
D	Prostaglandin E ₂ (PGE ₂)	
E	Adenosine	
F	Thromboxane A ₂ (TXA ₂)	The diagram illustrates the eicosanoid pathway. It begins with Linoleic acid (an essential fatty acid found in the cell membrane) being converted to Arachidonic acid (a C20 fatty acid) by Phospholipase A₂. Arachidonic acid is then converted to 5-HPETE by 5-Lipoxygenase. 5-HPETE is further converted to Leukotriene A₄ (LTA₄) by 5-Lipoxygenase. LTA₄ is then converted to various Leukotrienes (LTC₄, LTD₄, LTE₄, LTB₄) by Leukotriene synthase. Arachidonic acid is also converted to Prostaglandin H₂ by COX-1 and COX-2. Prostaglandin H₂ is then converted to various Prostaglandins (PGE₁, PGE₂, PGF_{2α}) by Prostaglandin synthase. Prostaglandin H₂ is also converted to Thromboxane A₂ (TXA₂) by Thromboxane synthase. The diagram also shows the effects of various drugs: Glucocorticoids (cortisone, prednisolone, triamcinolone, dexamethasone) inhibit Phospholipase A₂; Nonselective NSAIDs (aspirin (irreversible), diclofenac, ketorolac, ibuprofen, naproxen, indomethacin, meloxicam) and Selective NSAIDs (COX-2 inhibitors) (celecoxib) inhibit COX-1 and COX-2; 5-Lipoxygenase inhibitors (zileuton) inhibit 5-Lipoxygenase; Leukotriene receptor antagonists (montelukast, zafirlukast) inhibit Leukotriene receptors; and Thromboxane synthase inhibitors (epoprostenol, alprostadil, dinoprostone, carboprost) inhibit Thromboxane synthase.

	Answer	Image
G	Prostacyclin (PGI ₂)	

Hint

This patient presents with anterolateral myocardial infarction (MI) as a result of thrombotic occlusion of the left main coronary artery. The formation of the metabolite that is responsible for this finding is irreversibly inhibited by aspirin, which reduces the risk of MI.

Correct Answer

A - Prostaglandin E₁ (PGE₁)

Explanation Why

[Prostaglandin E₁ \(PGE₁\)](#) is produced by [platelets](#), but [PGE₁](#) normally causes [vasodilation](#) and inhibits thrombus formation. [PGE₁](#) does not cause thrombosis and vasoconstriction, which are responsible for this patient's [coronary angiography](#) finding.

B - Leukotriene D₄ (LTD₄)

Explanation Why

[Leukotrienes](#) such as [leukotriene D₄ \(LTD₄\)](#) are inflammatory mediators that are synthesized from [arachidonic acid](#) by [5-lipoxygenase](#) in [leukocytes](#), not [platelets](#). [LTD₄](#) increases [capillary](#) permeability, causes bronchoconstriction, and is implicated in the pathogenesis of [allergic rhinitis](#) and [asthma](#). [LTD₄](#) does not cause thrombosis and vasoconstriction, which are responsible for this patient's [coronary angiography](#) findings.

C - Nitric oxide (NO)

Explanation Why

[Nitric oxide](#) (NO) is synthesized primarily in [endothelial](#) cells; [platelets](#) only produce a minimal amount of NO. NO activates [guanylyl cyclase](#) to increase [cGMP](#) synthesis, which dilates blood vessels (due to vascular [smooth muscle](#) relaxation) and inhibits [platelet aggregation](#). NO does not cause thrombosis and vasoconstriction, which are responsible for this patient's [coronary angiography](#) findings.

D - Prostaglandin E₂ (PGE₂)

Explanation Why

[Prostaglandin E₂ \(PGE₂\)](#) can be produced by activated [platelets](#), but [PGE₂](#) has a variable effect on [platelet aggregation](#) (an inhibitory effect at high concentrations and a potentiating effect at low concentrations). In addition, [PGE₂](#) causes [vasodilation](#) (due to vascular [smooth muscle](#) relaxation). This patient's angiography findings are the result of thrombosis and constriction of a coronary vessel. A different substance that is released from activated [platelets](#) is primarily responsible for this finding.


E - Adenosine

Explanation Why

Infarcted cardiac [myocytes](#) release [ATP](#), which is converted to adenosine extracellularly by nucleotidases. Adenosine stimulates the adenosine A₁ receptors on cardiac afferent nerve endings and is responsible for [chest pain](#) in patients with [myocardial infarction](#). However, adenosine inhibits [platelet aggregation](#) and induces [vasodilation](#) by stimulating A_{2A} receptors; it would not cause thrombosis and vasoconstriction, which are responsible for this patient's angiography findings. Moreover, adenosine is not directly released from [platelets](#). [Adenosine diphosphate \(ADP\)](#), which is released from the dense granules of [platelets](#) following [platelet activation](#), would facilitate thrombus formation instead.

F - Thromboxane A₂ (TXA₂)

Image



Explanation Why

Thromboxane A₂ (TXA₂) is a key mediator of [platelet](#)-induced [coronary artery](#) stenosis and thrombosis. It is released primarily by activated [platelets](#) and promotes [platelet aggregation](#), vasoconstriction, and the activation of new [platelets](#). The [arachidonic acid](#) pathway is responsible for **TXA₂** synthesis. [Membrane phospholipid](#) (i.e., [linoleic acid](#)) is converted to [arachidonic acid](#) by [phospholipase A₂](#). In [platelets](#), [arachidonic acid](#) is converted by [cyclooxygenase-1 \(COX-1\)](#) to [prostaglandin H₂](#), which is then converted into **TXA₂** by thromboxane [synthetase](#). [Aspirin](#), which irreversibly inhibits [COX-1](#) and [COX-2](#), prevents the formation of [prostaglandin H₂](#) and, by extension, **TXA₂**. Since [platelets](#) do not have [DNA](#) and cannot synthesize new [COX-1](#) enzyme, the [platelet](#) inhibitory effects of [aspirin](#) last until new [platelets](#) are created.

G - Prostacyclin (PGI₂)

Explanation Why

[Prostacyclin](#) I₂ (PGI₂) is primarily produced by [endothelial](#) cells, not [platelets](#). PGI₂ inhibits [platelet aggregation](#) and promotes [vasodilation](#) in pulmonary and systemic vascular beds; it does not cause thrombosis and vasoconstriction, which are responsible for this patient's [coronary angiography](#) findings.

Question # 11

A 30-year-old woman comes to the physician because of worsening fatigue and shortness of breath for the past 7 months. She emigrated from China 2 years ago. Cardiac examination shows a loud S₁. Which of the following contributes most to this auscultation finding?

	Answer	Image
A	Tricuspid valve opening	
B	Mitral valve closure	
C	Aortic valve closure	

	Answer	Image
D	Atrial contraction	The image displays four stacked graphs over time, divided into Systole (Contraction, Ejection, Relaxation) and Diastole (Filling). 1. Pressure (mmHg): Shows atrial pressure (blue), ventricular pressure (green), and aortic pressure (yellow). Atrial pressure rises during contraction (1-2) and falls during relaxation (3-4). Ventricular pressure rises during ejection (2-3) and falls during relaxation (3-4). Aortic pressure rises during ejection (2-3) and falls during relaxation (3-4). 2. Volume (ml): Shows stroke volume (red), ventricular volume (green), and atrial volume (blue). Atrial volume rises during contraction (1-2) and falls during relaxation (3-4). Ventricular volume rises during ejection (2-3) and falls during relaxation (3-4). Stroke volume is the difference between ventricular and atrial volume. 3. Propagation of excitation: Shows the P wave (atrial contraction) and the QRS complex (ventricular contraction). 4. Heart sounds: Shows the S1 (closure of AV valve) and S2 (closure of AV valve) sounds.
E	Pulmonary valve opening	The image displays two stacked graphs over time, divided into Diastole and Systole. 1. Electrocardiogram (ECG): Shows the P wave, QRS complex, and T wave. The P wave is labeled 'Ejection click' and the QRS complex is labeled 'A2 P2'. 2. Phonocardiogram (PCG): Shows the S1 (closure of AV valve) and S2 (closure of AV valve) sounds. The S1 is labeled 'Ejection click' and the S2 is labeled 'A2 P2'.
F	Ventricular filling	The image displays a graph of ventricular pressure (mmHg) and volume (ml) over time. The pressure curve shows isovolumetric contraction (1-2), ejection (2-3), and isovolumetric relaxation (3-4). The volume curve shows rapid filling (1-2), reduced filling (2-3), and isovolumetric relaxation (3-4). The stroke volume (EDV - ESV) is indicated. Below the graph are three diagrams showing the effect of changes in contractility, preload, and afterload on the pressure-volume loop.

Hint

Worsening fatigue and dyspnea with a loud S₁ in a young individual who recently immigrated from a developing country raises concern for a sequela of rheumatic heart disease.

Correct Answer


A - Tricuspid valve opening

Explanation Why

[Tricuspid valve](#) opening can be affected in patients with [rheumatic heart disease](#) as a result of tricuspid stenosis (TS). However, in the case of TS, [tricuspid valve](#) opening would be associated with an [opening snap](#), which is distinct from the S₁ heard during early [diastole](#) (i.e., immediately after S₂). The closure rather than the opening of the [tricuspid valve](#) contributes to S₁. [Tricuspid valve](#) closure in patients with TS can cause a loud S₁, as seen here, but TS would cause [features of right heart failure](#) (e.g., peripheral [edema](#), [ascites](#)) rather than those of [left heart failure](#) (e.g., [dyspnea](#)).

B - Mitral valve closure

Image



Explanation But


MS is associated with other auscultatory findings such as an [opening snap](#) followed by a low-pitched [diastolic murmur](#) with presystolic accentuation. However, as MS progresses, left atrial pressure increases and causes the [opening snap](#) to occur earlier in the [cardiac cycle](#). If [atrial fibrillation](#) develops in a patient with MS, the presystolic accentuation of the [murmur](#) will be absent.

Explanation Why

The [first heart sound](#) (S₁) is a physiological sound created by the closure of the [mitral valve](#) and [tricuspid valve](#) that occurs at the onset of isovolumetric ventricular contraction, when ventricular pressure exceeds atrial pressure. S₁ is best heard in the 5th left [intercostal space](#) in the [midclavicular line](#) (cardiac apex). In patients with [mitral stenosis](#) (MS) and mobile leaflets (i.e., in mild to moderate MS), a loud S₁ is often present because the increased left atrial pressure moves the mitral leaflets further apart during [diastole](#) so that the leaflets snap shut with a greater velocity at the start of ventricular [systole](#). If the leaflets are immobile (e.g., in severe MS) or if the leaflets fail to close completely (e.g., in [mitral regurgitation](#)), S₁ would be soft. In the US, [rheumatic heart disease](#) is the most common cause of [mitral stenosis](#) in patients that have recently migrated from a developing country.

C - Aortic valve closure

Image




Explanation Why

[Aortic valve](#) closure can be affected in patients with [rheumatic heart disease](#) as a result of [aortic stenosis](#), which would also cause fatigue and [dyspnea](#), as seen here. However, [aortic valve](#) closure generates the A₂ component of S₂, and patients with [aortic stenosis](#) would typically have a soft S₂ (due to immobile atrial valve leaflets) and/or [paradoxical splitting](#) of S₂. [Aortic valve](#) closure does not contribute towards the generation of S₁.

D - Atrial contraction

Image



Explanation Why

Late [diastolic](#) contraction of the [atria](#) (“atrial kick”) against a stiff, noncompliant ventricle can cause S4. This auscultatory finding can be seen in patients with [hypertrophic cardiomyopathy](#), which can present in young adults with worsening fatigue and [dyspnea](#), as seen here. However, atrial contraction is not responsible for the generation of S1.

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>


USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>

E - Pulmonary valve opening

Image




Explanation Why

The opening of a stiff [pulmonary valve](#) is responsible for a [pulmonary valve ejection click](#), which can be heard in patients with mild or moderate [pulmonary stenosis](#) (PS). However, a [pulmonary valve ejection click](#) is heard during early [systole](#) (i.e., immediately after S₁); [pulmonary valve](#) closure does not contribute to the generation of S₁. Moreover, PS is typically encountered among pediatric patients with other congenital [heart](#) diseases (e.g., [tetralogy of Fallot](#)).

F - Ventricular filling

Image



Explanation Why

Rapid [ventricular filling](#), which occurs during early [diastole](#), can generate [S₃](#) if the [ventricular filling](#) pressure is increased (e.g., [mitral regurgitation](#), [heart failure](#)) or if the ventricles are dilated (e.g., [dilated cardiomyopathy](#)). However, [ventricular filling](#) itself does not significantly increase ventricular pressure, so it does not contribute to the generation of S₁. Rather, isovolumetric ventricular contraction, which causes ventricular pressure to surpass atrial pressure, leads to the event that generates S₁.

Question # 12



A 38-year-old man comes to the physician because of recurrent palpitations over the past 2 months. He says that he was diagnosed with a minor heart defect during childhood but does not remember the details and has not had regular medical follow-up examinations since. Physical examination shows a heart murmur. Cardiac catheterization is performed and hemodynamic results show:

	Oxygen saturation (%)	Pressure (mm Hg)
Right atrium	75	3
Right ventricle	75	25/4
Pulmonary trunk	75	22/10
Left atrium	99	10
Left ventricle	99	165/45
Ascending Aorta	99	155/70

Which of the following is the most likely diagnosis?

	Answer	Image
A	Mitral stenosis	

	Answer	Image
B	Mitral regurgitation	A pressure-volume (PV) loop for the left ventricle in mitral regurgitation. The y-axis represents LV pressure in mmHg (0 to 100), and the x-axis represents LV volume in mL (50 to 200). The loop is shaded light blue. The systolic pressure is approximately 100 mmHg. The aortic valve closure occurs at a volume of about 120 mL. The mitral valve opens at approximately 100 mL, and the regurgitant volume is the area between the aortic and mitral closure points, which is large. The diastolic pressure is low, around 10 mmHg.
C	Aortic valve regurgitation	A pressure-volume (PV) loop for the left ventricle in aortic valve regurgitation. The y-axis represents LV pressure in mmHg (0 to 200), and the x-axis represents LV volume in mL (50 to 200). The loop is shaded light blue. The systolic pressure is approximately 120 mmHg. The aortic valve closure occurs at a volume of about 120 mL. The mitral valve opens at approximately 100 mL, and the regurgitant volume is the area between the aortic and mitral closure points, which is large. The diastolic pressure is low, around 10 mmHg.
D	Ventricular septal defect	Three diagrams (1, 2, 3) illustrating the path of blood flow in a ventricular septal defect (VSD). The diagrams show the heart in cross-section with the VSD. Diagram 1 shows the normal flow of blood from the left ventricle to the aorta. Diagram 2 shows the flow of blood from the left ventricle to the right ventricle through the VSD. Diagram 3 shows the flow of blood from the right ventricle to the pulmonary artery.

	Answer	Image
E	Aortic stenosis	 A pressure-volume (PV) loop graph for the left ventricle. The y-axis is labeled 'LV pressure [mmHg]' with a scale from 0 to 200. The x-axis is labeled 'LV volume [mL]' with a scale from 0 to 150. The loop shows a normal diastolic pressure-volume relationship (dashed line) and a normal passive pressure-volume relationship (dotted line). The systolic pressure is significantly elevated, reaching approximately 200 mmHg, which is shaded in light blue. The area under the curve is shaded in light blue, indicating increased stroke work. The stroke volume is approximately 120 mL.
F	Atrial septal defect	 Three diagrams of the heart showing the progression of an atrial septal defect (ASD). Diagram 1 shows a small ASD with minimal shunting. Diagram 2 shows a moderate ASD with increased shunting. Diagram 3 shows a large ASD with significant shunting and pulmonary hypertension. The diagrams are labeled 1, 2, and 3 in the top left corner of each panel.


Hint

The patient has a widened pulse pressure in the ascending aorta.

Correct Answer

A - Mitral stenosis

Image




Explanation Why

[Mitral stenosis](#) can cause a [heart murmur](#) (decrecendo [diastolic murmur](#)), [palpitations](#), and normal [oxygen saturation](#), as seen in this patient. However, [mitral stenosis](#) would increase left atrial pressure (N = 2–12 mm Hg) and decrease [left ventricular diastolic](#) pressure (N = 5–12 mm Hg) due to a decreased amount of blood reaching the [left ventricle](#) during [diastole](#), unlike the normal left atrial pressure and increased [left ventricular diastolic](#) pressure seen here. Moreover, this patient's increased [systolic](#) pressures in the aorta and [left ventricle](#) are not consistent with [mitral stenosis](#).

B - Mitral regurgitation

Image




Explanation Why

[Mitral regurgitation](#) can cause a [murmur](#) ([holosystolic murmur](#)), [palpitations](#), and normal [oxygen saturation](#), as seen here. However, chronic [mitral regurgitation](#) increases left atrial pressure (N = 2–12 mm Hg), unlike the normal left atrial pressure seen here. Moreover, [mitral regurgitation](#) does not explain this patient's increased aortic or [left ventricular](#) pressure.

C - Aortic valve regurgitation

Image



Explanation But


This patient's underlying [heart](#) defect is most likely a [bicuspid aortic valve](#), which is one of the most common causes of [aortic insufficiency](#) and can lead to symptoms such as [palpitations](#).

Explanation Why

[Aortic valve regurgitation](#) causes a backward flow from the aorta during [diastole](#). This backflow decreases [diastolic](#) pressure in the aorta (N = 60–90 mm Hg) and increases [diastolic](#) pressure in the [left ventricle](#) (N = 5–12 mm Hg). This regurgitant blood combines with blood from the [left atrium](#) to fill the [left ventricle](#), which results in an increased [preload](#) and a subsequent increase in [stroke volume](#). As a result, the [systolic](#) pressure in both the [ascending aorta](#) (N = 90–140 mm Hg) and in the [left ventricle](#) (N = 90–140 mm Hg) is increased. [Oxygen saturation](#) and right [heart](#) pressures are normal in [aortic regurgitation](#). If left untreated, this condition leads to [eccentric hypertrophy](#) of the [left ventricle](#) and [left heart failure](#).

D - Ventricular septal defect

Image




Explanation Why

[Ventricular septal defects \(VSD\)](#) can cause a [heart murmur \(holosystolic murmur\)](#) and [palpitations](#). However, [VSDs](#) usually increase [right ventricular](#) pressure (N = 15–30/1–7 mm Hg) and [right ventricular oxygen saturation](#) (N = 60–80%) due to the shunting of oxygen-rich blood from the [left ventricle](#) to the [right ventricle](#), unlike the normal [right ventricular](#) pressure and [oxygen saturation](#) seen here. Moreover, [VSD](#) would not explain this patient's increased aortic and [left ventricular](#) pressures.

E - Aortic stenosis

Image




Explanation Why

[Aortic stenosis](#) can cause a [heart murmur](#) (crescendo-decrescendo [systolic murmur](#)), [palpitations](#), normal [oxygen saturation](#), and increased [left ventricular systolic](#) pressure (due to higher ejection resistance), as seen here. However, aortic [systolic](#) pressure would be decreased in these patients, not increased. Additionally, a decreased [diastolic](#) pressure in the aorta is inconsistent with [aortic stenosis](#).

F - Atrial septal defect

Image



Explanation Why




[Atrial septal defects](#) (ASD) can cause a [heart murmur](#) ([systolic](#) pulmonary ejection [murmur](#)) and [palpitations](#), both of which are seen in this patient. However, in the case of an ASD, pressures in the [right atrium](#) (N = 2–7 mm Hg), [right ventricle](#) (N = 15–30/1–7 mm Hg), and [pulmonary trunk](#) (N = 15–30/4–12 mm Hg), and the [oxygen saturation](#) in the [right atrium](#) (N = 60–80%) would be increased as a result of a left-right shunt, unlike the findings in this patient. Moreover, this patient's increased [systolic](#) pressures in the aorta and [left ventricle](#) are not consistent with ASD.

Question # 13

Twelve hours after being admitted to the hospital for an inferior wall myocardial infarction, a 65-year-old woman begins to present with dyspnea, dizziness, and syncopal episodes that last a few seconds. Her pulse is 40/min. An ECG is shown. Which of the following is the most likely diagnosis?



	Answer	Image
A	First-degree atrioventricular block	

	Answer	Image
B	Atrial fibrillation	
C	Mobitz type I block	
D	Sinus bradycardia	
E	Atrial flutter	

	Answer	Image
F	Mobitz type II block	
G	Third-degree atrioventricular block	


Hint

This patient's ECG shows a ventricular rate of 40/min and an atrial rate of 100/min. The placement of a temporary transcutaneous or transvenous pacemaker is indicated.

Correct Answer

A - First-degree atrioventricular block

Image




Explanation Why

First-degree [atrioventricular blocks](#) can develop after an acute inferior [myocardial infarction](#) (the [right coronary artery](#) also supplies the [atrioventricular node](#), which is where this block occurs). However, patients with [first-degree AV block](#) are typically asymptomatic. Moreover, the condition is characterized by a prolonged [PR interval](#) (> 200 ms) with a [QRS complex](#) following every [P wave](#) (i.e., all [P waves](#) are conducted through to the ventricles), which is not the case here.

B - Atrial fibrillation

Image




Explanation Why

[Atrial fibrillation](#) (AF) is a common complication of [acute myocardial infarction](#) that can cause [dyspnea](#), [dizziness](#), and [syncope](#), as seen here. However, AF is characterized by an irregularly irregular rhythm with varying R-R intervals and absent [P waves](#). Moreover, AF is commonly associated with a rapid ventricular response (ventricular rate > 100 bpm), which is not consistent with this patient's [bradycardia](#).

C - Mobitz type I block

Image



Explanation Why

[Dyspnea](#), [dizziness](#), syncopal episodes, and [bradycardia](#) may rarely be seen in [Mobitz type I](#) block, a potential complication of an inferior [myocardial infarction](#) (the [right coronary artery](#) also supplies the [atrioventricular node](#), which is where this block occurs). However, [Mobitz type I](#) blocks are characterized by progressive lengthening of the [PR interval](#) until a beat is dropped (i.e., a nonconducted [P wave](#) that is not followed by a [QRS complex](#)), which is not the case here.

D - Sinus bradycardia


Explanation Why

[Dyspnea](#), [dizziness](#), and [syncope](#) may rarely be seen in sinus [bradycardia](#), a potential complication of an inferior [myocardial infarction](#) (because the [right coronary artery](#) also supplies the [sinoatrial node](#)).

However, the [ECG](#) in patients with sinus [bradycardia](#) shows [P waves](#) of normal morphology that are always followed by [QRS complexes](#), which is not the case here.

E - Atrial flutter

Image



Explanation Why

[Atrial flutter](#) can complicate an [acute myocardial infarction](#) and cause [dyspnea](#), [dizziness](#), and [syncope](#) (even though most patients with [atrial flutter](#) are asymptomatic). However [atrial flutter](#) is a type of [supraventricular tachycardia](#) that is typically associated with a fast [heart rate](#), which is not consistent with this patient's [bradycardia](#). Moreover, flutter waves (i.e., sawtooth appearance of [P waves](#) that occur in sequence at a rate of ~ 300 bpm), which are the characteristic [ECG](#) finding in [atrial flutter](#), are not seen here.

F - Mobitz type II block

Image

Explanation Why

[Dyspnea](#), [dizziness](#), syncopal episodes, [bradycardia](#), and [ECG](#) showing atrioventricular dissociation after a recent [myocardial infarction](#) may raise suspicion for a [Mobitz type II](#) block (due to damage to the His-Purkinje system, which is where this block most commonly occurs). However, [Mobitz type II](#) blocks are characterized by intermittent dropped beats (i.e., nonconducted [P waves](#) that are not followed by [QRS complexes](#)) and a constant [PR interval](#), which is not the case here.

G - Third-degree atrioventricular block

Explanation But

[Third-degree AV block](#) occurs due to impaired conduction at the level of the [AV node](#) (~20%), [bundle of His](#), or (most commonly) below the [His bundle](#) (~60%). The escape [QRS complex](#) may be

narrow or wide depending on the location of the block. Narrow QRS complexes (< 120 ms), as seen here, indicate a block that is above the [His bundle](#); a wide QRS complex is associated with a block at or below the [His bundle](#).

Explanation Why

This patient's [dyspnea](#), [dizziness](#), syncopal episodes ([Stokes-Adams attacks](#)), [bradycardia](#), recent inferior [myocardial infarction](#) ([MI](#)), and [ECG](#) findings are consistent with a [third-degree atrioventricular block](#) (i.e., complete [AV block](#)). In [third-degree AV blocks](#), there is no rhythmic relationship between the [atria](#) ([P waves](#)) and ventricles ([QRS complexes](#)), which beat independently of one another. Additionally, the atrial rate is greater than the ventricular rate (i.e., more [P waves](#) than [QRS complexes](#)) and both rates are constant, as seen in this patient. [Third-degree AV block](#) may complicate an inferior [MI](#) (because the [AV node](#) and [His bundle](#) are primarily supplied by the [right coronary artery](#)) or an [anterior MI](#) (because the [His bundle](#) may receive blood from the septal branches of the [left coronary artery](#)). Other causes of [third-degree AV block](#) include infections (e.g., [Lyme disease](#), bacterial [endocarditis](#)), [cardiomyopathy](#), [hyperkalemia](#), and [iatrogenic](#) effects (e.g., cardiac interventions, drugs such as [beta blockers](#)).

USMLE Telegram Channels

MedQuest Videos: <https://t.me/USMLEMedQuest>

Kaplan Classroom Videos: <https://t.me/USMLEKaplan>

USMLE Rx Videos Step 1: <https://t.me/USMLERxStep1>

Becker Videos: <https://t.me/USMLEBecker>

Boards & Beyond Videos Step 1: <https://t.me/USMLEBoardsAndBeyond>

Boards & Beyond Videos Step CK: <https://t.me/USMLEBoardsAndBeyondStep2CK>

Physeio Videos: <https://t.me/USMLEPhyseio>

Pixorize Videos: <https://t.me/USMLEPixorize>

Pathoma Videos: <https://t.me/USMLEPathoma>

Goljan Audios High Quality: <https://t.me/USMLEGoljanAudios>

Amboss Group: <https://t.me/USMLEAmboss>

Lecturio QBank: <https://t.me/USMLELecturio>

Endpoint PDFs: <https://t.me/USMLEEndpoint>

USMLE NBMEs: <https://t.me/USMLENBME>

ONLINE MedED: <https://t.me/USMLEOnlineMedED>

USMLE World Step 1: <https://t.me/USMLEWorldStep1>

USMLE World Step 2 CK: <https://t.me/USMLEWorldStep2CK>

USMLE World Step 3 Qbank: <https://t.me/USMLEWorldStep3>

USMLE Anki: <https://t.me/USMLEAnki>

KissPrep: <https://t.me/USMLEKissPrep>