

QUIZ TIME

oral physiology

Done by : Lemar Mustafa

Lec: 22



لجنة
طب الأسنان
جامعة مؤتة

1. The Maxillary sinus is also known as:

- A) Frontal sinus
- B) Antrum of Highmore
- C) Sphenoidal sinus
- D) Ethmoidal sinus

Answer: B) Antrum of Highmore

2. At what stage does the Maxillary sinus begin to develop?

- A) At birth
- B) 3-4 months of intrauterine life (MIU)
- C) 7 years of age
- D) During puberty

Answer: B) 3-4 months of intrauterine life (MIU)

3. Which of the following is the largest of all paranasal air sinuses?


- A) Frontal sinus
- B) Ethmoidal sinus
- C) Sphenoidal sinus
- D) Maxillary sinus

Answer: D) Maxillary sinus

4. The Maxillary sinus communicates with the nasal cavity through the:

- A) Superior meatus
- B) Middle meatus
- C) Inferior meatus
- D) Sphenoethmoidal recess

Answer: B) Middle meatus



5-The Maxillary sinus is lined by which type of epithelium?

- A) Stratified squamous epithelium
- B) Simple cuboidal epithelium
- C) Pseudostratified ciliated columnar epithelium
- D) Transitional epithelium

Answer: C) Pseudostratified ciliated columnar epithelium

6-At birth, the size of the maxillary sinus is approximately:

- A) 15 x 15 x 10 mm
- B) 30 x 25 x 20 mm
- C) 8 x 6 x 4 mm
- D) 2 x 2 x 2 mm

Answer: C) 8 x 6 x 4 mm

7-Which of the following is the correct order of teeth proximity to the sinus floor (from most close to least)?

- A) 8, 7, 6, 5, 4
- B) 6, 7, 5, 4, 8, 3
- C) 3, 4, 5, 6, 7
- D) 1, 2, 3, 4, 5

Answer: B) 6, 7, 5, 4, 8, 3

