




# Action potential

BY  
Dr. Nour A. Mohammed  
Mut'ah school  
of medicine



# Definition

it is a **transient** change in the resting membrane potential as a result of application of a **threshold stimulus**.



depolarisation moves along the axon


# Application of an adequate electric stimulus to the nerve fiber is followed by:

- 1) Stimulus artifact ✓
- 2) Latent period ✓
- 3) Spike potential
  - a) Depolarization
  - b) Repolarization
    - Rapid Repolarization
    - Slow Repolarization
    - Hyperpolarization

# Stages of action potential:

## (1) Stimulus artifact



- It is a short irregular deflection of the base line due to stimulus application.
- This is due to current leakage from the stimulating electrode to the recording electrode (indicates beginning of stimulation).



## (2) Latent period

- It represents the time that the nerve impulse (**response**) takes to travel from the stimulating to recording electrode.
- It indicates the rate of conduction in the axon.

Speed of conduction =  $\frac{\text{Distance between electrodes}}{\text{Latent period}}$




### (3) Depolarization

There is a rapid loss or (reversal) of polarity of the membrane.

It is recorded as a rise of membrane potential in the positive direction.


- At first, there is a slow depolarization of 25 mV (RMP changes from  $-90$  to  $-65$  mV)



depolarisation moves along the axon

## Depolarization steps:


- At -65 mV (*Firing level*), the depolarization occurs rapidly till the potential reaches zero mV (no difference between inside and outside the cell membrane).
- Then, the inside becomes positive to the outside of the membrane i.e. reversal of polarity (**overshoot**) till +35 mV.



So, the magnitude of the depolarization phase equals 125 mV (from  $-90$  to  $+35$  mV).


## Cause of depolarization:

- ✓ The stimulus opens some  $\text{Na}^+$  channels allowing  $\text{Na}^+$  to enter the cell.
- ✓ If the  $\text{Na}^+$  influx achieves threshold potential (the firing level) then additional  $\text{Na}^+$  gates open and depolarization will proceed rapidly.
- ✓ The flow of  $\text{Na}^+$  will cause more  $\text{Na}^+$  channels to open. (+ve feedback mechanism).




## Changes in voltage-gated fast Na<sup>+</sup> channels and Na<sup>+</sup> permeability during action potential:

- **Outer gate (activation gate):** opens at the start of depolarization causing Na<sup>+</sup> influx.
- **Inner gate (inactivation gate):** then closes, preventing further Na<sup>+</sup> influx and causing Na<sup>+</sup> channel inactivation.



- **During rest:** the activation gate is closed & the inactivation gate is opened → no  $\text{Na}^+$  permeability.
- **During activation:** change of membrane potential by 25 mV (from -90 to -65 mV) → the activation gate opens and  $\text{Na}^+$  permeability reaches maximum till the potential of +35 mV. Then the inactivation gate closes.




## (4) Repolarization


It is the return of the membrane potential to the resting state (from +35 to -90 mV).

It occurs in 3 steps:

1. Rapid repolarization
2. Slow repolarization
3. Hyperpolarization.



# 1. Rapid repolarization:




- During which the membrane restores 70% of its resting condition.
- *Cause:*
- a) Inactivation of voltage gated  $\text{Na}^+$  channels so,  $\text{Na}^+$  influx stopped.
- b) Activation of voltage gated  $\text{K}^+$  channels so,  $\text{K}^+$  outflux increased.

## Changes in voltage-gated $K^+$ channels during action potential:

$K^+$  channel has a single gate located on the inside of the membrane.

- ✦ **During rest:** the gate is closed.
- ✦ **During activation:** depolarization  $\rightarrow$  slow opening of  $K^+$  channels which coincides to the closure of  $Na^+$  gates  $\rightarrow$  repolarization.




Note that:

$K^+$  gates are stimulated by the same stimulus that stimulates  $Na^+$  channels but  $K^+$  channels are slow (take more time to open and close).

## (2) Slow repolarization

After 70% of repolarization, the rate of repolarization becomes slow.




### Cause:

Decrease in  $\text{K}^+$  gradient  $\rightarrow$  slow  $\text{K}^+$  efflux  $\rightarrow$  delayed repolarization.


## (3) hyperpolarization

After reaching the RMP, there is an overshooting of about **1-2 mV** hyperpolarization, then the membranes returns to normal **RMP**.



**Cause:** Delayed K<sup>+</sup> channels closure  
→ more K<sup>+</sup> efflux  
→ more hyperpolarization.

- Finally, *Na<sup>+</sup>-K<sup>+</sup> pump* helps to restore the normal ionic distribution of the RMP i.e., maintenance of Na<sup>+</sup> (extra cellular) and K<sup>+</sup> (intracellular )



## Properties of action potential:

- 1) Caused by **threshold** stimulus (or suprathreshold)
- 2) Caused by ionic changes.
- 3) Conducted (propagates) in both directions.
- 4) Constant duration.
- 5) Obeys All or none law  $\Rightarrow$  can't be graded (constant amplitude).
- 6) Has Absolute Refractory Period  $\Rightarrow$  can't be summated.

## Excitability changes:

At first, there is increase in excitability till the firing level then the following changes occur:

### 1- Absolute refractory period:

- No response to any stimulus (loss of excitability).
- Coincides with depolarization from the firing level till the first 1/3 of rapid repolarization.

## 2- Relative refractory period:

- Stronger stimulus → response (low excitability).
- Coincides with lower 2/3 of rapid repolarization.

### 3- Supernormal phase:


- Weak stimulus → response (high excitability).
- Coincides with the Slow repolarization
- During it, the membrane is partially depolarized and has low threshold for firing level.

## 4- Subnormal phase:

-Stronger stimulus → response  
(low excitability).

- Coincides with the hyperpolarization

- During it, the membrane is hyperpolarized with increase threshold for firing level and difficult stimulation.



Thank you

