

# **Gastrointrotology**

## **Lecture 1**



**By**

**Dr\ Nour A. Mohammed**

**Mutah school of medicine**

# Salivary secretion & Swallowing





# *Salivary Secretion*

- **Saliva** : 1.5 L/day, hypotonic, pH is 6.3-6.8
- **Salivary glands** : three pairs of salivary glands

| <b>Parotid</b>                                                                                                                                                    | <b>Sub maxillary</b>                                                                       | <b>sublingual</b>                                                                                                      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"><li>• 20 %.</li><li>• Serous acini for secretion (watery &amp; rich in enzy.).</li><li>• supplied by glossopharyngeal</li></ul> | <ul style="list-style-type: none"><li>• 75 %.</li><li>• Mixed.</li><li>• Facial.</li></ul> | <ul style="list-style-type: none"><li>• 5 %.</li><li>• Mucus acini (thick, rich in mucin).</li><li>• Facial.</li></ul> |


**Sublingual gland**

**Submandibular duct**

**Parotid duct**

**Submandibular gland**

**Parotid gland**



- 
- **Composition of saliva : a- 99.5 % water. b- 0.5 % solids.**
  - **0.3 % organic** : as enzymes (amylase, Lipase, Lysozymes) and mucus.
  - **0.2 % inorganic:**
    - ❖ **Buffers** as **phosphate** & **bicarbonate** buffering systems
    - ❖ **Soluble calcium salts:** which saturate saliva to prevent decalcification of teeth.
    - ❖ **Some electrolytes** as  $\text{Na}^+$  ,  $\text{Cl}^-$ ,  $\text{Hco}_3^-$ , and  $\text{K}^+$  ,they act as coenzymes for salivary enzyme amylase.



# Functions of saliva

1. **Facilitation of speech** and deglutition.
2. **Cleaning (hygiene)** of the mouth by washing and antibacterial effect of lysozymes & immunoglobulin A
3. **Buffering function** : by bicarbonate and phosphate systems to keep the PH at about 7.0 → the teeth do not loose their calcium.  
Also, saliva neutralizes gastric secretion in case of gastroesophageal reflux.
4. **Digestive function** :
  - **Ptyalin (salivary  $\alpha$ - amylase)** : digest starch to maltose in PH 6.9 so it is inhibited in the stomach.
  - **Lingual Lipase**: digest **30 % of lipids** and secreted from Ebner's gland of tongue.
5. **Excretory function** : of lead, mercury, fluoride and some drugs as morphine and alcohol.
6. **Facilitate taste sensation**
7. **Regulation of water balance** ( $\downarrow$  in dehydration and give thirst

# The Stages of salivary secretion

## ➤ I) Salivary acini (Primary):

→ saliva similar in composition to plasma  
isotonic

## ➤ II) Salivary duct (secondary):

due modification by the duct under effect of **aldosterone** hormone → active reabsorption of  $\text{Na}^+$ ,  $\text{Cl}^-$  &  $\text{HCO}_3^-$  and active secretion of  $\text{K}^+$ . So, saliva becomes hypo- tonic to plasma .

# Innervation of salivary glands:

## *A-Parasympathetic*

*It arises from superior salivatory nucleus in the **pons** → chorda tympani as a branch of **the facial nerve** → **submandibular ganglion** → submandibular and sublingual glands.*

Also, inferior salivatory nucleus in **medulla oblongata** → lesser superficial petrosal nerve as a branch of **glossopharyngeal nerve** → otic ganglion → parotid gland

→ **True secretion** : large in volume watery, rich in enzymes , Na<sup>+</sup>, CL<sup>-</sup>, Hco<sub>3</sub>.

***Parasympathetic causes V.D of blood vessels of salivary glands***

# Sympathetic Stimulation

- It arises from lateral horn cells of the upper two thoracic segments and relay in the superior cervical sympathetic ganglia→
- **Trophic secretion**: little in volume, viscous, and rich in mucin.
- **VC** of blood vessels of salivary glands.

## **N.B**

if the flow of salivary secretion increased → little time for modification  
→ ↑ Na<sup>+</sup>, Cl<sup>-</sup>, HCO<sub>3</sub><sup>-</sup> & ↓ K<sup>+</sup> concentration as in **parasympathetic stimulation**.

# Control of salivary secretion

➤ Nervous only via conditioned and unconditioned reflexes.

➤ **[I] Unconditioned reflex**

➤ Inborn reflex that needs no previous learning.

➤ **Stimuli** : direct contact of food , Chewing

➤ **Receptor** : taste receptors.& Receptors in GIT wall.

➤ **Afferent**

- Chorda tympani : from ant. 2/3 of tongue.
- Glossopharyngeal : from post. 1/3 of tongue
- Lingual nerve : movement of tongue.
- Vagus nerve : from epiglottis.

➤ **Center** : superior & inferior salivatory nuclei in brain stem

➤ **Efferent**: chordae tympani & glossopharyngeal.


➤ **Response**: ↑ salivary glands secretion.

## [II] Conditioned reflex :

- Acquired reflexes and need previous learning
- **Stimuli :**
  - Sight of food.
  - Smelling of food.
  - Hearing about food.
  - Thinking of food.
- **Receptors :** special sense receptors.
- **Afferent :** optic, olfactory & auditory nerves.
- **Center :** to cerebral cortex → salivatory nuclei.
- **Efferent & response** → as unconditioned reflex.

# Pharynx and oesophagus






**Pharynx**

# Pharynx

- It is a common pathway for respiratory and digestive system and has swallowing receptor area and the primary peristalsis waves start from it. It is separated from oesophagus by the upper oesophageal sphincter which is normally closed.

# Esophagus




# Oesophagus

- It is a muscular tube has outer longitudinal and inner circular muscle layers which are striated in the upper portion and smooth in the lower portion .So, the peristalsis in the upper portion depends on the vagovagal reflex, however in the lower portion it depends on the local enteric reflex.

# Swallowing (Deglutition)

- It is the propelling of food bolus from mouth to stomach.
- It is under control of the **swallowing center** in the **medulla**.
- *It can be divided into 3 phases:*





# Swallowing (Deglutition)

➤ *It can be divided into 3 phases:*

➤ **Buccal phase:**

(voluntary) elevation and retraction of tongue against the hard palate propels the bolus to the pharynx.

➤ **Pharyngeal phase (involuntary) :**

It is very rapid (1 second), occur reflexely via :

**Swallowing reflex**

# Swallowing reflex

- **Receptor:** in oropharynx (tonsillar pillars).
- **Afferent:** glossopharyngeal nerves.
- **Center:** medulla oblongata (swallowing center).
- **Efferent:** motor fibers of cranial nerves V, IX, X, XII .

➤ **Response:** series of reflexes to prevent entry of food into air passages:

Elevation of soft palate → closure of nasal cavity.


Approximation of palatopharyngeal folds → sagittal slit through which small food particles pass and prevent passage of large particles.

Closure of glottis (opening of larynx) by approximation of vocal cords & elevation of larynx and folding of epiglottis

Inhibition of breathing (**swallowing apnea**).

Relaxation of pharyngoesophageal sphincter and contraction of superior pharyngeal muscle → rapid pharyngeal peristalsis → forces the food into relaxed upper esophagus.


# Swallowing



# Esophageal phase (involuntary)

- *Upper esophageal sphincter : (UES)*
- The pharyngeo – esophageal junction is normally closed by striated muscle tone to prevent entry of inspired air into stomach. During swallowing the sphincter relaxes reflexely and then reclosed after swallowing.
- *Traveling along the esophagus*

# Smooth Muscle Peristalsis Moves Food Along Alimentary Canal



- Entry of food bolus into the esophagus initiate peristaltic waves of 2 types :

### *Primary peristaltic waves :*

- They start at the upper end of esophagus.
- They are continuation of the pharyngeal peristalsis.
- It travels at the rate of **2-4 cm/sec**. But gravity may increase velocity of food bolus .

# Secondary peristaltic waves

- Presence of bolus in the esophagus initiate peristaltic waves at site of bolus.
- These waves repeated until food bolus is driven down the stomach.
- Peristaltic movements in the **upper half** of esophagus is coordinated by **vago – vagal reflex** (striated ms.), while in **lower half** is coordinated by **local enteric reflex**.

# *Lower esophageal sphincter (LES)*

- It is called the cardiac sphincter.
- It is the lower 3-5 cm of the esophagus.
- It has high resting tone (**High – pressure zone**) and exert a pressure **15-30 cm H<sub>2</sub>O** above intra – abdominal pressure to prevent reflux of gastric content into esophagus.
- It is relaxed when food bolus reaches it **with some delay**, so this area is liable to damage or ulceration by cold, hot and spicey food.

# ***Lower esophageal sphincter (LES)***

## ➤ **Its tone is increased by : (contracted)**

Sympathetic alpha adrenergic.

Gastrin hormone (so, drugs which neutralize gastric acidity → ↑  
gastrin hormone release → contraction of the LES.

## ➤ **Its tone is decreased by : (Relaxed)**

Inhibitory vagal via VIP secretion.

Some food as fats, chocolate, alcohol & coffee.

# How gastric reflux into oesophogus is prevented ?

- High pressure zone sphincter.
- The intra abdominal small part of the oesoph. is squeezed by the increased intra abdominal pressure.
- The oesophagus enters the stomach in acute angle and act as a flap.
- Gastrin hormone increases the tone in the lower oesophagus.

# Gastro esophageal reflux

➤ is return of gastric contents to esophagus due to failure of anti reflux mechanisms as weak sphincter pressure.

➤ **-Increases in:**

pregnancy, smoking, ↑ coffee , alcohol & obesity.

➤ **-Leads to:**

Ulcer of lower esophagus.

Heat burn.

Stricture of cardiac sphincter.

Barrett's oesophagus due to prolonged effect on mucosa which are premalignant.



**Thank you**