



- Function of neuroenteric canal? Select one:
 - a. Maintains the closure between amniotic cavity and yolk sac.
 - b. Maintains the drainage of amniotic cavity.
 - C. Maintains the pressure between amniotic cavity and yolk sac.
 - d. Maintains the drainage of yolk sac.
 - e. Maintains the connection between amniotic cavity and yolk sac.

***C

- The epiblast of the male is ? Select one:
 - a. Endoderm
 - b. Mesoderm
 - C. Ectoderm
 - d. Exoderm
 - e. Hypoderm

***C

· Primitive streak is?

Selectone:

- a.Endoderm
- b.Mesoderm
- c.Ectoderm
- d.Smooth muscle
- e.Mucus membrane



***C

· The source of the notochord is?Selectone:

- a.Hansen's membrane
- b.Primitive node
- c.Amniotic cavity
- d.Primitive streak
- e.Prechordal plate

***b

· The notochord is stopped at? Selectone:

- a.Buccolaryngeal membrane.
- b.Prechordal sac
- c.Cloacalme mbrane
- d.Prechordal plate
- e.Heuser's membrane

***d

· Closure of notochordal canal by? Selectone:

- a.The cells of the medial of the notochordal canal

- b. The cells of the roof of the notochordal canal
- C. The cells of the lateral of the notochordal canal
- d. The cells of the median of the notochordal canal
- e. The cells of the floor of the notochordal canal

***b



· Primitive pit starts? Select one:

- a. Cranial
- b. Deep
- c. Ventral
- d. Caudal
- e. Dorsal

***d



