

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



السلام عليكم ورحمة الله وبركاته


# Biostatistics

LV

**PROF. DR. WAQAR AL-KUBAISY**

**18-7-2022**


# Interpreting Standard Deviation



For bell-shaped distributions, the following statements hold:

- Approximately 68% of the data fall between  $\bar{x} - 1s$  and  $\bar{x} + 1s$
- Approximately 95% of the data fall between  $\bar{x} - 2s$  and  $\bar{x} + 2s$
- Approximately 99.7% of the data fall between  $\bar{x} - 3s$  and  $\bar{x} + 3s$

For NORMAL distributions, the word 'approximately' may be removed from the above statements.



Example: Suppose the Hb levels of 150 women has a roughly bell-shaped distribution with a mean of 12 mg/dl. and standard deviation of 0.10 g/dl.


- a) Give the interval of the amount of Hb level that approximately 68% of the women will have

$$12 - 0.1 \text{ to } 12 + 0.1 = 11.9 \text{ to } 12.1 \text{ g/dl.}$$

- b) Give the interval of the amount of Hb level that approximately 95% of the women will have


$$12 - 2(0.1) \text{ to } 12 + 2(0.1) = 11.8 \text{ to } 12.2 \text{ g/dl.}$$

# **Important or Uses of SD**




# Normal Distribution Curve

Age (year)	Freq.	Commutative frequency	Relative frequency	% R.F.	Cumulative R.F.	%cum · Freq.
<b>20-29</b>	<b>1</b>	<b>1</b>	<b>0.02</b>	<b>2</b>	<b>0.02</b>	<b>2</b>
<b>30-39</b>	<b>2</b>	<b>3</b>	<b>0.04</b>	<b>4</b>	<b>0.06</b>	<b>6</b>
<b>40-49</b>	<b>2</b>	<b>5</b>	<b>0.04</b>	<b>4</b>	<b>0.1</b>	<b>10</b>
<b>50-59</b>	<b>3</b>	<b>8</b>	<b>0.06</b>	<b>6</b>	<b>0.16</b>	<b>16</b>
<b>60-69</b>	<b>12</b>	<b>20</b>	<b>0.24</b>	<b>24</b>	<b>0.4</b>	<b>40</b>
<b>70-79</b>	<b>14</b>	<b>34</b>	<b>0.28</b>	<b>28</b>	<b>0.68</b>	<b>68</b>
<b>80-89</b>	<b>12</b>	<b>46</b>	<b>0.24</b>	<b>24</b>	<b>0.92</b>	<b>92</b>
<b>90-99</b>	<b>4</b>	<b>50</b>	<b>0.08</b>	<b>8</b>	<b>1.00</b>	<b>100</b>
<b>total</b>	<b>50</b>	<b>---</b>	<b>1</b>	<b>100</b>	<b>---</b>	<b>---</b>



In large population  $\longrightarrow$  Graphically  $\longrightarrow$  Form of **Curve**  
**Normal Distribution curve,**  
**Gaussian Curve ,**  
**Bell Curve .**

In NDC



- ❖ All the **observation** are lying in area under the curve
- ❖ Average measures (mean Md , Mo) in the center of in the center of observation .
- ❖ Rest of observations distributed around the average measures .
- ❖ in a **homogenous** form
- ❖ *Half of them **higher** than the mean*  $\bar{X}$   $\bar{X}$
- ❖ Half of them **lesser** than the mean

So

- ❖ the distribution of observation in NDC is symmetrical .  $\longrightarrow$


under the NDC divided by

1- measures of C.T

$\bar{X}$

2-measures of variability

S.D




## By Measures of C.T

$\bar{X}$

Divided the area under the curve into two equal halves of observation,

**50 %** of observation their values less than  $\bar{X}$  value

and **50 %** of observation their values higher than  $\bar{X}$




## By Measures of variability (S.D)

S.D and its multiplicity ( one S.D, two S.D, three S.D divided the area under the NDC into small areas, each area containing certain and **fixed proportion** of observation

Within  $\pm 1$  S.D from the  $\bar{X}$

Within  $\pm 2$  S.D from the  $\bar{X}$


Within  $\pm 3$  S.D from the  $\bar{X}$



Within  $\pm 1$  S.D from the  $\bar{X}$   
68% of observations, (34% each side)  
68% of observation deviated from the  $\bar{X}$  by not more than  
 $\pm 1$  S.D                      ????????

Within  $\pm 2$  S.D from the  $\bar{X}$   
95% of observations lie,  
95% of observations deviated from the  $\bar{X}$  by not more than  
 $\pm 2$  S.D .    ????????

Within  $\pm 3$  S.D from the  $\bar{X}$   
99% of observations are located,  
99% of observations deviated from the  $\bar{X}$  by not more than  
 $\pm 3$  S.D .    ????????



??????????


# Characteristics of the NDC

1. Bell shape .
2. Symmetrical distribution of observations on both sides
3. Unimodal ??????????.
4. Curving downward on both sides from the mean toward the horizontal, **but never touch it** .
5. Mean, Median and Mode of distribution are identical or coincide .
6. All the Medical, Biological phenomenon following its distribution .

7- Area under curve divided by

Mean into two equal halves

8. Between  $\bar{X}$  and certain multiplicity of S.D on either side an area containing fixed proportion of observation





7/18/2022 **68% 99% 95% .**


## Importance

1-Most of the phenomenon in Medical field follow this distribution .

2-It is for **justification** and **calculation** of confidence interval .

3-It is form the **basis** of most of **significance testing** hypothesis .  
That is most test of significance depend on the theory of ND





Different samples  $\rightarrow$  different  $\bar{X}_s$  even if the sample size are equal

There is a **variation in the  $\bar{X}_s$**  of different samples  
This variation is **due to sampling variation.**

## Sampling Variability

$\bar{X}$

Mean  $\pm$  S.D of sample .


The **interest of sample** not in its own right **but**  
what it **tell us** about the **population** which this sample  
represent

The aim of Biostatistics is to have



a **sound** generalized **information** about the **population**  
**from** which the **sample** has been **drown**, depending on  
**evidence of this sample**


## Cont. ...Sampling Variability



Sample Statistics


true mean ,

Population Parameters



mean of  
universe

## Cont. ...Sampling Variability




**Different samples  $\rightarrow$  different  $\bar{X}_s$  even if the samples size are equal**

**There is a variation in the  $\bar{X}_s$  of different samples  
This variation is due to sampling variation.**

Cont. ...Sampling Variability

mean of universe  
true mean



Population mean  $\mu$   
usually unknown

the sample measurement ( mean  $\pm$  S.D) is  
**not exactly reflect** its population .

There is a **difference** between sample mean  $\bar{X}$  and  
population mean  $\mu$



## Cont. ...Sampling Variability

There is a difference between

**sample statistics** and **population parameters**,  
this variation is called **sampling error**


There is a **difference between** sample means and  
population mean.?????????

□ Deviation of the samples mean  $\bar{X}$  ) from the  
population mean ( $\mu$ )


✓ this **will be the S.D of sample mean** ( $\bar{X}$ .  
from the population mean ( $\mu$ )

▪ **Average** of S.D of sample means from  
population mean which is

□ known as **Standard Error**



## Cont. ...Sampling Variability



This mean that samples  $\bar{X}_s$  distributed **around population mean**, or Samples  $\bar{X}$ 's scatter around the  $\mu$ .

*The measurement of this scattering equal to  
S.D of the sample  $\bar{X}$*

7/18/2022

## Standard Error S.E

- It is the **average** deviation of the sample mean (  $\bar{X}$  ) from the **true** (population) mean ( $\mu$ ) of the population . So
- ❖ *it is equal to the **S.D** of sample mean  $\bar{X}$  divided by the square root of the **sample size (N)***

$$S.E = \frac{S.D}{\sqrt{N}}$$

- depend on
- ❖ sample size
  - ❖ S.D of sample

The larger the sample size (N) → smaller the S.E  
The smaller the S.D of sample → smaller the S.E

# Standard Error S.E

## Example

**8 plasma values of uric acid**

the mean ( $\bar{X}$ ) of uric acid is  **$3 \pm 0.31$**

$$S.E = \frac{0.31}{\sqrt{8}} = 0.11$$

$$S.E = \frac{S.D}{\sqrt{N}}$$

**16 plasma values of uric acid the mean ( $\bar{X}$ ) of uric acid is  $3 \pm 0.31$**

$$= \frac{0.31}{\sqrt{16}} = 0.0775$$

$$\frac{0.21}{\sqrt{16}} = 0.0525$$

$$\frac{0.41}{\sqrt{16}} = 0.1025$$

□ Distribution of samples mean ( $\bar{X}_S$ ) around the population mean ( $\mu$ ) in NDC area

❖ is similar to that

❖ of the distribution of X (values) around sample mean  $\bar{X}$


sample means  $\bar{X}_S$  deviated from  $\mu$  by

✓ S.E and its multiplicity, so

deviated from  $\mu$  by

1 S.E, 2 S.E and 3 S.E in proportion

68% 95% 99% .





## Remember that the


### ❖ SD is

➤ measure of spread of the data in a single **sample** .

### ❖ The S.E. is

➤ a measure of spread in **ALL** sample means from a **population**.


☐ **We notice** that the as sample size  $n$  increases the S.E decrease



???????

Thank you!

The image features the words "Thank you!" in a highly decorative, bubbly font. Each letter is a different color and has a textured, hand-drawn appearance. The 'T' is yellow, the 'h' is orange, the 'a' is green with a dotted border, the 'n' is purple, the 'k' is orange, the 'y' is green, the 'o' is red, and the 'u' is blue. The exclamation point is also blue. Several stylized flowers are scattered around the text: a large pink flower with a yellow center is in the foreground, and several smaller blue and light blue flowers are positioned above and to the sides of the letters. The entire graphic is set against a light blue background.



## Importance

**1-Most of the phenomenon in Medical field follow this distribution .**

**2-It is for justification and calculation of confidence interval .**

**3-It is form the basis of most of significance testing hypothesis .  
That is most test of significance depend on the theory of NDC .**